

THE PROBLEM OF NEUTRINO MASSES IN EXTENSIONS OF THE STANDARD MODEL ¹

ISABELLA MASINA

*Department of Physics G. Galilei, Univ. of Padova
Via Marzolo 8, I-35131 Padova, Italy ²*

We review the problem of neutrino masses and mixings in the context of Grand Unified Theories. After a brief summary of the present experimental status of neutrino physics, we describe how the see-saw mechanism can automatically account for the large atmospheric mixing angle. We provide two specific examples where this possibility is realized by means of a flavour symmetry. We then review in some detail the various severe problems which plague minimal GUT models (like the doublet-triplet splitting and proton-decay) and which force to investigate the possibility of constructing more elaborate but realistic models. We then show an example of a quasi-realistic SUSY $SU(5)$ model which, by exploiting the crucial presence of an abelian flavour symmetry, does not require any fine-tuning and predicts a satisfactory phenomenology with respect to coupling unification, fermion masses and mixings and bounds from proton decay.

¹Based on PhD thesis work, February 2001

²Now at Service de Physique Théorique, CEA-Saclay, F-91191 Gif-sur-Yvette, France
e-mail: masina@spht.saclay.cea.fr

Contents

Introduction	1
1 Neutrino Masses and Mixings	7
1.1 Present Experimental Status and Future Perspective	8
1.1.1 Atmospheric	8
1.1.2 Solar	10
1.1.3 LSND	12
1.1.4 Three Flavour Analysis	12
1.2 Beyond the Standard Model	13
1.2.1 Neutrino Masses and Lepton Number Violation	13
1.2.2 Smallness of Neutrino Masses	15
1.2.3 Approximate Mixing Matrix in the Neutrino Sector	17
1.3 How to Reproduce the Large Atmospheric Mixing	18
1.3.1 Mass Hierarchies and Approximate Zeroth Order Textures . .	18
1.3.2 Degenerate Neutrinos	20
1.3.3 Hierarchical Neutrinos	21
2 Models with Large Mixing	25
2.1 A Model with Large Mixing from m_D	25
2.1.1 The Choice of Horizontal Abelian Charges	26
2.1.2 Charged Fermion Mass Matrices	27
2.1.3 Neutrino Sector	28
2.1.4 Outlook	31
2.2 A Model with Large Mixing from Nothing	32
2.2.1 Renormalization Effects	36
2.2.2 An Explicit GUT Example with Broken Flavour Symmetry . .	38
2.2.3 Outlook	42

3	Problems of SUSY GUTs	43
3.1	The Doublet-Triplet Splitting Problem	44
3.1.1	The Doublet-Triplet Splitting Problem in Minimal $SU(5)$. .	44
3.2	Solutions to the Doublet-Triplet Splitting Problem	45
3.2.1	The Missing Doublet Mechanism	46
3.2.2	The Sliding Singlet Mechanism	48
3.2.3	The Missing VEV Mechanism	48
3.2.4	Higgses as pseudo-Goldstone Bosons	49
3.3	Proton Decay	50
3.3.1	Dimension Five Operators	52
3.3.2	Hunting Dominant Operators	53
3.3.3	Proton Decay from Non-Renormalizable Operators	56
3.4	How to Prolong Proton Lifetime	56
3.4.1	Altering Triplets Yukawa Couplings	57
3.4.2	Raising the Triplet Mass	57
3.4.3	Moving the Two CP Violating Phases	58
3.4.4	Playing with SUSY Particles	59
3.4.5	Numerical Examples	60
3.5	Fermion Masses and Mixings	63
3.5.1	Mass Relations	63
3.5.2	How to Achieve Realistic Spectra	64
3.6	Coupling Unification	66
3.6.1	Coupling Unification Beyond the Leading Order Approximation	70
4	A realistic SUSY $SU(5)$ GUT Model	77
4.1	The Model	78
4.1.1	$SU(5)$ Breaking	79
4.2	Doublet-Triplet Splitting	79
4.2.1	Limit of Unbroken SUSY	79
4.2.2	Broken Supersymmetry	81
4.3	Coupling Unification	85
4.3.1	Estimate of $\alpha_3(m_Z) _{\overline{MS}}$	85
4.3.2	Theoretical Uncertainties	89
4.4	Yukawa Couplings	90
4.4.1	A Numerical Example	92
4.5	Proton Decay	95
	Conclusions	99

Acknowledgements	101
Bibliography	105
A The Calculation of Proton Lifetime	115
A.1 d=5 B Violating Operators	115
A.2 The Relevant Operators	117
B Representative SUSY Spectrum	121
C Group Theory for the MDM	123
C.1 Basic Conventions for the Tensors in $SU(5)$	123
C.2 The 75 and 50 representations	125
C.3 The Method for Catching SM Multiplets	128
C.4 Basic Formulae for the MDM	133

Introduction

It is quite obvious that the Standard Model must be extended. Various arguments bring along the important conceptual implication that the Standard Model should be seen as *a low energy effective theory* valid up to some physical cut-off scale.

It is thus compelling to speculate about the underlying more fundamental theory. Various ideas have been proposed and developed to answer this question. For instance, recently a lot of interest and efforts have been devoted to the idea of extra-dimensions, where the different behaviour of the gauge and gravitational interactions may provide a solution to the hierarchy problem. But the attractive proposal of Grand Unification has not lost its charm.

The concept that all particle interactions unify at very high energies is indeed not very recent. The first paper on this subject was published 26 years ago and from then on this idea captured the interest of physicists. Many fundamental papers that explore the various aspects and consequences of Grand Unification have been written. What is then inside this proposal which renders it still so interesting to collect the general interest? The main point is that it received *phenomenological support* by:

- A) the success of gauge coupling unification;
- B) recent developments on neutrinos.

A) Gauge coupling unification works terribly well. The original proposal has however notably evolved because now we know that Grand Unification needs Supersymmetry. Only then gauge coupling unification successfully happens at the scale $M_{GUT} \sim 10^{16}$ GeV. In addition, in non supersymmetric Grand Unified theories the prediction of proton lifetime for the process $p \rightarrow e^+ \pi^0$ is in conflict with lower experimental limits, but including Supersymmetry it turns out to be close to the safe side.

B) Recent data coming from neutrino physics are much more robust and less fragmented then they were just few years ago. There is definitive evidence for a neutrino mass associated with atmospheric oscillations in the range $0.05 \div 1$ eV. There is no room in the Standard Model for neutrinos to have a mass, so that neutrinos represent the first true signal and actually give some hints on possible

extensions of the Standard Model. Baryon and lepton numbers are expected to be violated in Grand Unified Theories, leading to the fundamental consequences of proton instability and possibility of Majorana mass terms. The previously quoted mass range points in favour of lepton number violation at a scale $\sim 10^{(14\div 15)}$ GeV.

Thus both gauge coupling unification and neutrino masses independently strongly suggest that beyond the electroweak scale there is another crucial scale where some new physics really arises and which lies just two orders of magnitude below the Planck scale.

Neutrinos are curious also because, in contrast with the observed smallness of left handed quark mixings, it is now experimentally well established by Super-Kamiokande that they possess a large left handed mixing associated with the neutrino flavours involved in atmospheric oscillations. *To explain from the theoretical point of view the possible causes which determine the atmospheric mixing to be large and the possible consequences is the thread of the present review.*

Grand Unification represents an excellent framework for the study of neutrino masses and mixings, because it predicts fermion mass matrices to be related. Several Grand Unified models have been constructed which show that the presence of the large atmospheric mixing may naturally arise via the see-saw. These models can be distinguished into two classes, according to the basic mechanism exploited.

Most models are based on the possible presence of a large right handed down quark mixing, not directly observable in weak transitions, which is strictly correlated to a large left handed mixing for charged leptons. The latter can be transferred to the Dirac neutrino mass matrix and the desired large mixing is achieved if the right handed Majorana neutrinos are not widely split.

But there is also a particularly interesting alternative mechanism. This mechanism is based on the observation that for a sufficiently hierarchical right handed neutrino spectrum, a large atmospheric mixing automatically arises if the left handed ν_μ and ν_τ couple with comparable strength to the right handed ν_μ^c . Thus, remarkably, this situation does not require the presence of large mixings in the Dirac and/or Majorana mass matrices. As opposite to the first mechanism, this second one can then be realized even if all left and right mixings of quarks and leptons are very small. It is then a possibility for the large mixing to actually arise from nothing, that is only as an effect of the see-saw mechanism. This situation is particularly attractive from the theoretical point of view because it is compatible with left right symmetric scenarios.

The first two chapters of this review contain a summary of the present experimental status and future perspectives about neutrinos, show the two basic mechanisms providing an explanation of the large atmospheric mixing and present explicit Grand Unified models which realize these two possibilities.

The two independent supports in favour of the physical relevance of M_{GUT} , i.e. supersymmetric coupling unification and neutrino masses, can be hardly interpreted as just a coincidence. Armed with the results obtained from the study of the neutrino sector, we face the more ambitious task of *constructing a sufficiently complete and realistic Grand Unified model which could embed one of the two mechanisms for the generation of the large atmospheric mixing*. After 26 years, one could wonder if it is still possible and eventually what kind of informations could be added to further increase our knowledge of Grand Unification. There are mainly two new facts:

1. much stronger limits on proton decay rates, in particular on the channels which are expected to be dominant in supersymmetric Grand Unified theories. Super-Kamiokande gives a lower limit for proton lifetime of about $2 \cdot 10^{33}$ yr for the channel $p \rightarrow K^+ \bar{\nu}$;
2. the approach itself. Since Grand Unification seems so successful, it is desirable to use it as a framework where to address also the flavour problem, that is to explain the observed hierarchical spectra and mixings of all fermions, including neutrinos.

The flavour problem is independent of Grand Unification, in the sense that Grand Unification alone is not able to provide any light on this. However it is really an optimal framework in which to embed possible solutions of the flavour problem, because Grand Unification predicts more or less strict group theoretical relations between fermion mass matrices. One interesting way to address the flavour problem is by introducing an horizontal flavour symmetry, namely a symmetry under which the three copies of each fermion transform differently.

Thus the modern point of view is to construct not too complicated, sufficiently realistic and relatively complete models. These models have to be enough detailed to actually carry out the calculation of as much low energy observables as possible. We are not satisfied with too rough and incomplete models. It is clear that only those models which provide predictions in agreement with experiment survive.

In this respect, minimal versions of Grand Unified Theories (e.g. $SU(5)$ with only 24, 5 and $\bar{5}$ of Higgs fields) are not realistic. They suffer from serious problems:

- doublet-triplet splitting;
- predictions of proton decay rates;
- masses and mixings.

To address some of these points, an enormous fine-tuning is required, e.g. of about 14 orders of magnitude for the doublet-triplet splitting. But the serious fact is

that, even if one tolerates to do this unpleasant tunings, minimal models give wrong predictions. In fact, minimal models predict protons decaying at a too fast rate for the channel $p \rightarrow K^+ \bar{\nu}$, in conflict with experiment (point 1.) and give wrong relations between fermion masses (point 2.).

It is clear that something is lacking in minimal models. In particular it would be of crucial importance to extend the Higgs sector in order to:

- give an explanation to the doublet-triplet splitting;
- enhance the prediction of proton lifetime;
- adjust down quarks and charged lepton spectra and correctly describe neutrino masses.

However, the addition of new fields in the Higgs sector must be done in such a way not to destroy the only firm starting point, that is coupling unification. Here troubles come because a realistic model requires so many improvements over the minimal versions that one could think that the beautiful idea of Grand Unification is actually not realizable. Note that the effort does not go in the direction of making precise predictions on poorly measured observables, but rather of doing predictions which are in not rough conflict with experiment.

The problems and the technical difficulties which are typical of minimal models are reviewed in chapter 3, together with a review of the main ideas proposed to overcome them.

Many attempts of constructing realistic models have been done in the context of $SO(10)$ but in general they lack of predictivity on unification, which should then occur in an accidental way. This is due to the fact that despite $SO(10)$ is interesting because it contains right handed neutrinos, it requires a lot of assumptions on the Higgs content in order to reduce its rank, thus reaching $SU(5)$ or $SU(4) \otimes SU(2) \otimes SU(2)$.

We instead focus the attention directly on Supersymmetric $SU(5)$, which already possesses the same rank as the Standard Model, trying here to address as many problems as possible. We present a model which successfully overcomes fine-tunings problems and which gives predictions in agreement with experiments. This model exploits the crucial presence of an additional $U(1)$ flavour symmetry.

The fine-tunings linked to the doublet-triplet splitting are avoided by adopting the Missing Doublet Mechanism and by using the flavour symmetry to stabilize the Higgs doublet mass against large mass corrections due to non renormalizable operators. Fermion masses and mixings are reproduced, at the level of the correct order of magnitude, by the same flavour symmetry. In the neutrino sector the preferred solution is one with nearly maximal mixing both for atmospheric and

solar neutrinos, which turn out to be compatible with the LOW solution. The large atmospheric mixing follows from the presence of large left mixings both in the Dirac neutrino and charged lepton mass matrices. The latter is correlated to an unobservable large right mixing of the down quark mass matrix. A third effect of the flavour symmetry is to adequately suppress non renormalizable operators inducing proton decay at a too fast rate. Remarkably, the presence of the large representations demanded by the Missing Doublet Mechanism for the solution of the doublet-triplet splitting problem, makes possible a decrease of the predicted value of $\alpha_3(m_Z)$ assuming coupling unification thanks to an increase of the effective mass that mediates proton decay induced by dimension 5 operators. As a consequence the value of the strong coupling is in better agreement with the experimental value and the proton decay rate is still compatible with present bounds, even if it is difficult to avoid the conclusion that proton decay must occur with a rate at reach of current and next generation experiments. Chapter 4 contains a detailed description of this model. We think that this model is interesting because it proves that a Supersymmetric $SU(5)$ Grand Unification is not excluded and offers a benchmark for comparison with experiment.

Chapter 1

Neutrino Masses and Mixings

Recent data from Super-Kamiokande [1] have provided a more solid experimental basis for neutrino oscillations as an explanation of the atmospheric neutrino anomaly. Also the solar neutrino deficit, observed by several experiments, is probably an indication of a different sort of neutrino oscillations. Neutrino oscillations imply neutrino masses, so that it is compelling to look for viable extensions of the Standard Model. In order to fulfill such a program, it is important to recognize that the extreme smallness of neutrino masses in comparison with quark and charged lepton masses indicates that the formers have different nature, linked to lepton number violation.

Thus, neutrino masses provide some insight on the very large energy scale where lepton number is violated and on Grand Unified Theories (GUTs). Indeed, GUTs constitute a very natural framework for the generation of neutrino masses, because they generically predict lepton and baryon number violation. Neutrino masses could also give an important feedback on the problem of quark and charged lepton masses, as all these masses are related in GUTs. In particular the observation of a nearly maximal mixing angle for $\nu_\mu \rightarrow \nu_\tau$ is particularly interesting. At present, a large solar neutrino mixings seems to be slightly more preferable than a small one. Large mixings in the neutrino sector are very interesting because a first guess was in favour of small mixings, in analogy to what is observed for left mixings in the quark sector. If confirmed, single or double maximal mixings can provide an important hint on the mechanism that generate neutrino masses. With neutrino masses settled, observation of proton decay will be the next decisive challenge remained to support or eventually put in crisis GUTs.

After a concise summary of the present experimental status of neutrino masses and mixings and its future perspective, in this chapter we will review the most promising mechanisms which reproduce a large atmospheric mixing.

1.1 Neutrino Oscillations: Present Experimental Status and Future Perspective

Despite the remarkable progress reached during the last few years, the experimental status of neutrino oscillations is still very preliminary. While the evidence for the existence of neutrino oscillations from solar and atmospheric neutrino data is rather convincing by now, the values of the mass squared differences $|\Delta m^2|$ and mixing angles are not firmly established. However, there is a general confidence that future experiments will not only succeed in clarifying such a picture but hopefully will also find one of the CP violating phases of the neutrino sector and establish the sign of the Δm^2 .

1.1.1 Atmospheric

Present

Data with impressive high statistics are coming from Super-Kamiokande [1] which are in perfect agreement with the hypothesis of flavour oscillations [2, 3]. Such hypothesis is consistent with all Super-Kamiokande data and is also corroborated by independent atmospheric neutrino results from the Soudan-2 [4] and MACRO [5] experiments, as well as by the pioneering Kamiokande experiment [6].

Zenith-angle distributions, up-down asymmetries and so on provide plenty of smoking guns for the fact that something actually happens to atmospheric ν_μ as a function of LE^n with $n \approx -1$, where L is the baseline length and E the energy.

In the simplest 2ν case of $\nu_\mu \rightarrow \nu_\tau$, a Super-Kamiokande combined fit indicates that:

- (i) the preferred value of Δm_{atm}^2 lies in the range $(1.5 \div 5) \cdot 10^{-3} \text{ eV}^2$;
- (ii) a large mixing angle appears to be well established: $\sin^2 2\theta_{atm} > 0.88$ at 90% confidence level.

Interpreted in terms of 2ν oscillations, Super-Kamiokande favours $\nu_\mu \rightarrow \nu_\tau$ over $\nu_\mu \rightarrow \nu_s$ at 99% confidence level, based on separate analyses of the zenith-angle distributions. The same conclusions also come from the rate of π^0 production, but in this case systematic uncertainties are still large. Pure $\nu_\mu \rightarrow \nu_e$ transitions do not provide a good fit to the data, and are independently excluded by the negative ν_e disappearance searches in the CHOOZ [7] and Palo Verde [8] reactors.

However, these results have to be taken *cum grano salis*. Additional oscillation channels may be open, as naturally expected in 3ν and 4ν schemes accommodating the current phenomenology. In fact, it has been realized that the dominant $\nu_\mu \rightarrow \nu_\tau$ oscillations plus subdominant $\nu_\mu \rightarrow \nu_e$ oscillations are also consistent [9] with

combined Super-Kamiokande and CHOOZ data, leading to a much richer three-flavour phenomenology, as will be discussed in the following. On the other hand, when the data are analyzed [10, 11] for ν_μ oscillating into a linear combination of ν_τ and ν_s , one finds such a scenario to be still viable (and also compatible with solar neutrino data) for a sizeable region of parameter space. These analysis are very important to determine to what extent the 4ν scenario motivated by the LSND experiment can be accommodated.

It is clear that the definite evidence would be direct τ production, but this is not an easy task for Super-Kamiokande. Actually, there is not yet direct evidence for an oscillation pattern.

Thus, there is still room to speculate on other interpretations, like neutrino decay [12], decoherence scenarios [13] and ν_μ mixing with neutrino states propagating in large extra dimensions [14], which appear indistinguishable with present data with respect to the L/E distribution. Detailed analysis together with high statistics data could test these non standard explanations. For instance, the exotic scenario of neutrino decay seems to fail [9, 15] in reproducing the zenith-angle distributions of sub-GeV, multi-GeV and up-going muons observed by Super-Kamiokande.

Future

So, future experiments are highly needed, among which long-baseline laboratory experiments, which require a muon decay source for the neutrino beam, seem to be very promising in order to:

- (i) confirm the existence of atmospheric neutrino oscillations by using a controllable beam and measuring an oscillation pattern;
- (ii) precisely measure $|\Delta m_{atm}^2|$ and $\sin^2 2\theta_{atm}$;
- (iii) make new measurements possible.

The K2K experiment [16] ($L=250$ km, $\langle E_\nu \rangle = 1.4$ GeV, from KEK to Super-Kamiokande, in operation), finds an encouraging deficit of events with respect to the non oscillation hypothesis. The rate is perfectly compatible with $\Delta m_{atm}^2 = 3 \cdot 10^{-3}$ eV² and $\sin^2 2\theta_{atm} = 1$, but the statistics is still low and the significance only about 2σ .

Long-baseline investigations will be pursued by MINOS [17] ($L=732$ km, from Fermilab to Soudan, begin in 2002). In addition to (i) and (ii), it should be able to discriminate between $\nu_\mu \rightarrow \nu_\tau$ and $\nu_\mu \rightarrow \nu_s$ and to measure $|U_{e3}|$ from $\nu_e \rightarrow \nu_\mu$ appearance.

OPERA and ICANOE [18] ($L=743$ km, from CERN to Gran Sasso, approved), have the main purpose of looking for τ detection as the smoking gun for $\nu_\mu \rightarrow \nu_\tau$ oscillations. They also should be sensitive to $|U_{e3}|$.

With respect to long-term future, neutrino factories together with very long baselines ($L \sim 3000$ km) appear [19] to offer the possibility of determining the sign of Δm_{atm}^2 from matter effects on the Earth crust, measuring with an accuracy of tenths of degrees the angle θ_{13} , down to values of $\theta_{13} = 1^\circ$. The CP violating phase δ could also be measured if the solar neutrino parameters lie in the range of the MSW-LMA solution.

1.1.2 Solar

Present

For solar neutrinos the experimental situation and the interpretation in terms of frequency and mixing angles is less clear.

The solar neutrino experiments [20] sample different ν_e energy ranges and find different flux deficits compared to the Standard Solar Model (SSM) [21], as shown in Table 1.1, so that the ν_e survival probability is inferred to be energy dependent.

	$\text{flux}^{(\text{exp})} / \text{flux}^{(\text{SSM})}$
GALLEX	0.60 ± 0.06
SAGE	0.52 ± 0.06
Homestake	0.33 ± 0.03
Super-Kamiokande	0.47 ± 0.02

Table 1.1: For the energy range specific of each type of experiment, the ratio between the experimentally measured neutrino flux and the flux predicted by the SSM is shown.

Global oscillation fits have been made which include: total rates (the different suppression ratio for Homestake plays a vital role), day-night asymmetry, seasonal dependence beyond $1/r^2$. Typical candidate solar solutions are given in Table 1.2.

Interesting recent developments by Super-Kamiokande (with reduced background and threshold of 5 MeV) show that the flux deficit is confirmed but no other direct signal of oscillations emerges: the spectrum is compatible with flat, the day-night

solution	$\Delta m_{sol}^2 (eV^2)$	$\sin^2 2\theta_{sol}$
SMA-MSW	$\sim 5 \cdot 10^{-6}$	$\sim 5 \cdot 10^{-3}$
LMA-MSW	$\sim 2 \cdot 10^{-5}$	~ 1
LOW	$\sim 10^{-7}$	~ 1
VO	$\sim 10^{-(9 \div 10)}$	~ 1

Table 1.2: Typical solar solutions from global fits.

asymmetry is reduced at the 1.3σ level and the seasonal variation is compatible with the flux variation from the orbit eccentricity. These last two facts disfavour respectively the SMA-MSW and the VO at the 95% confidence level. We are left with the LMA-MSW and the LOW, both with large mixing. The pure transition $\nu_e \rightarrow \nu_s$ is disfavoured at 95%. It is important to keep in mind that the Δm_{sol}^2 values of the above solutions are determined by the experimental result that the flux suppression is energy dependent. If the Homestake indication were disregarded, new energy independent solutions would emerge, with much more extended ranges for Δm_{sol}^2 .

All these conclusions need to be confirmed by updated global analysis, without forgetting the region with $\theta_{sol} > \pi/4$, as well as matter and vacuum effects. It is also important to analyze the data in the 3ν and 4ν schemes, to understand to what extent a component $\nu_e \rightarrow \nu_s$ is tolerated [10].

Future

The experimental situation will be hopefully clarified with the results from future experiments like SNO [22], BOREXINO [23] and KamLAND [24].

In the present Phase I, SNO clearly sees the signal from charged current (CC) reactions $\nu_e + d \rightarrow p + p + e^-$, with an energy spectrum which seems to be proportional to that of the SSM. It also sees a clear signal from elastic scattering (ES) $\sum_i \nu_i + e \rightarrow \sum_i \nu_i + e$. The ratio CC/ES is expected to be measured in a sufficiently accurate way to reveal at 3σ if high energy neutrinos with $i \neq e$ are coming from the Sun at a rate

consistent with the observed deficit of ν_e with respect to the SSM expectation. In the following Phase II, the neutral current (NC) cross section $\sum_i \nu_i + d \rightarrow p + p + \sum_i \nu_i$ will be measured. With also the ratio of NC/CC, SNO should be able to discriminate between different solar neutrino scenarios.

BOREXINO can measure in real time the ${}^7\text{Be}$ flux and is sensitive to the VO solution.

KamLAND, starting in 2001, should be able to test definitively the LMA-MSW and LOW solutions, using neutrinos from nuclear power reactors. It should also measure the day-night and seasonal effects for ${}^7\text{Be}$ neutrinos.

Thus it seems possible that the solar neutrino problem will be better clarified in the near future.

1.1.3 LSND

The issue of the additional claim for $\nu_e \rightarrow \nu_\mu$ neutrino oscillations from the LSND experiment [25] is still open, because it is neither confirmed nor completely excluded by KARMEN-2 [26]. The range still surviving is at small mixing angle and with $\Delta m^2 = (0.2 \div 2) \text{ eV}^2$. This range is much above the solar and atmospheric neutrino frequencies, so that the need to accommodate a third frequency would imply the existence of a fourth light sterile neutrino, in addition to the three established weak interacting flavours, which should mix with solar and/or atmospheric neutrinos. MiniBOONE, starting in 2002, is an experiment designed to solve this problem. It will be possibly followed by BOONE, if the signal is confirmed.

1.1.4 Three Flavour Analysis

Given the present experimental uncertainties the theorist has to make some assumptions on how the data will finally look like in the future. Here and in all the following sections, we tentatively assume that the LSND evidence will disappear. If so then we only have two oscillations frequencies, which can be given in terms of the three known species of light neutrinos without additional sterile kinds.

As pointed out, Super-Kamiokande data are compatible with nearly maximal $\nu_\mu \rightarrow \nu_\tau$ two flavour oscillations, but, among the possible deviations from this standard picture, three flavour oscillations appear to be perfectly viable. In fact, Super-Kamiokande data do not exclude a non-vanishing U_{e3} element.

We then take for granted that the frequency of atmospheric neutrino oscillations will remain well separated from the solar neutrino frequency, even for the MSW solutions: $\Delta m_{atm}^2 > \Delta m_{sol}^2$. Assuming that two out of three active neutrinos are almost degenerate, say $m_1 \approx m_2$, one can show [9] that atmospheric neutrinos probe

only $\Delta m_{atm}^2 = m_3^2 - m_{1,2}^2$ and the mixing matrix elements $U_{\alpha 3}$, which satisfy the unitarity constraint $U_{e3}^2 + U_{\mu 3}^2 + U_{\tau 3}^2 = 1$. The unitarity constraint can be conveniently embedded in a triangle plot. The best fit for Super-Kamiokande data (70.5 kTy) is found at [27] $(\Delta m_{atm}^2, U_{e3}^2, U_{\mu 3}^2, U_{\tau 3}^2) = (3.5 \cdot 10^{-3} eV^2, 0.07, 0.57, 0.36)$. The tolerated $\nu_\mu \rightarrow \nu_e$ mixing is constrained by $U_{e3}^2 < 0.31$ at 90% confidence level.

The weak preference for $U_{e3}^2 \neq 0$ is suppressed by including CHOOZ data. The best fit for combined Super-Kamiokande and CHOOZ data gives $(\Delta m_{atm}^2, U_{e3}^2, U_{\mu 3}^2, U_{\tau 3}^2) = (3.0 \cdot 10^{-3} eV^2, 0., 0.5, 0.5)$, with the further constraint $U_{e3}^2 < 0.04$ at 90% confidence level.

It appears very difficult to probe, with present atmospheric data, values of U_{e3}^2 as small as a few %. Constraining U_{e3}^2 is one of the major tasks for future reactor and accelerator neutrino experiments.

1.2 Beyond the Standard Model

Assuming just three flavours, neutrino oscillations are due to a misalignment between the flavour basis $\nu' \equiv (\nu_e, \nu_\mu, \nu_\tau)$, where ν_e is the partner of the mass and flavour eigenstate e^- in a left-handed weak isospin $SU(2)$ doublet (similarly for ν_μ and ν_τ), and the neutrino mass eigenstates $\nu \equiv (\nu_1, \nu_2, \nu_3)$:

$$|\nu'\rangle = U|\nu\rangle, \quad (1.1)$$

where U is the MNS 3 by 3 mixing matrix [3]. The presence of mixing implies that neutrinos cannot be all massless and actually the presence of two different oscillation frequencies implies at least two different non zero masses. Neutrino oscillations are practically only sensitive to differences Δm^2 so that the absolute scale of squared masses is not fixed by the observed frequencies.

1.2.1 Neutrino Masses and Lepton Number Violation

Neutrino oscillations imply neutrino masses which in turn demand either the existence of right handed neutrinos (Dirac masses) or lepton number (L) violation (Majorana masses) or both. Anyway, one is forced to go beyond the Standard Model (SM). In fact, referring to three neutrino generations, in the SM Lagrangian density only the term $i\bar{\nu}_L^T \not{D} \nu_L$ appears, where D_μ is the $SU(2) \otimes U(1)$ gauge covariant derivative and $\nu_L = (\nu_e \ \nu_\mu \ \nu_\tau)^T$ is a column of three Weyl spinors in a four-component notation (a Weyl spinor has only two independent components) satisfying $\nu_L = 1/2 (1 + \gamma_5) \nu_L$. The standard convention is to assign $L(\nu_L) = 1$.

If no additional right handed neutrino field $N_R = (N_e \ N_\mu \ N_\tau)^T$ (satisfying $N_R = 1/2 (1 - \gamma_5)N_R$ and with $L(N_R) = 1$) is allowed, there isn't the possibility of including in the Lagrangian the Dirac mass term

$$\mathcal{L}_{Dir} \ni -\overline{\nu}_L^T m_D N_R + h.c. \quad (1.2)$$

which, by defining $\nu = \nu_L + N_R$, can be written as $\mathcal{L}_{Dir} \ni -\overline{\nu}^T m_D \nu$, m_D being a matrix in flavour space. The above term do not violate L, which is an additive quantum number.

For a massive charged particle of spin 1/2 one needs four states, while only two are enough for an intrinsically neutral particle. If L is violated, there is no conserved quantum number that really makes neutrinos and antineutrinos different and a new type of mass term is possible. We can have a Majorana mass term

$$\begin{aligned} \mathcal{L}_{Maj} &\ni -\frac{1}{2}(\overline{\nu^c})_R^T m_{LL} \nu_L + h.c. = -\frac{1}{2}\nu_L^t{}^T C m_{LL} \nu_L + h.c. \\ &\doteq -\frac{1}{2}\nu_L^T m_{LL} \nu_L + h.c. \end{aligned} \quad (1.3)$$

where $(\nu^c)_R = C\overline{\nu}_L^t$ and C is the 4 by 4 matrix in Dirac space that implements charge conjugation. We denote by t the transposition in Dirac space, while maintaining T for the transposition in flavour space. The Majorana mass term violates L by two units. Also, since in the SM ν_L is a weak isospin doublet, it transforms as a component of an isospin triplet. In the following, as we are only interested in flavour indices and not in Dirac indices, we will simply denote the Majorana term for left handed fields by $\nu_L^T m_{LL} \nu_L + h.c.$, omitting the Dirac matrix C and the transposition t . Note that if L is violated and N_R also exists, then a second type of Majorana mass is also possible which is

$$\mathcal{L}_{Maj} \ni -\frac{1}{2}N_R^T m_{RR} N_R + h.c. , \quad (1.4)$$

where we again omitted C and t . Clearly it also violates L by two units, but, since N_R is a gauge singlet, this term is invariant under the SM gauge group.

In conclusion, if N_R does not exist, we can only have a Majorana mass m_{LL} if L is violated. If N_R exists and L is violated, we can have both Dirac m_D and Majorana masses m_{LL} and m_{RR} .

If one wants to give masses to neutrinos but to avoid the conclusion that L is violated, then one must assume that N_R exists and that neutrinos acquire Dirac masses through the usual Higgs mechanism as quark and leptons do. Technically this is possible. But there are two arguments against this possibility.

1) The first argument is that neutrino masses are extremely small so that the corresponding Yukawa couplings would be enormously smaller than those of any other fermion. Note that within each generation the spread of masses is by no more than a factor $10^{3\div 5}$. But the spread between the t quark and the heaviest neutrino would exceed a factor of 10^{11} .

2) A second argument arises from the fact that once we introduce N_R in the theory, then the L violating term $N_R^T m_{RR} N_R$ is allowed in the Lagrangian density by the gauge symmetry. In the minimal SM, i.e. without N_R , L and B conservation are accidental global symmetries that hold because there is no operator term of dimension ≤ 4 that violates B and L but respects the gauge symmetry. On the other hand, in the presence of N_R , the dimension 3 operator $N_R^T m_{RR} N_R$ is gauge symmetric but violates L. We expect a mass factor in front of this operator in the Lagrangian density, and in the absence of a protective symmetry, we expect it to be of the order of the cut off, i.e. of order M_{GUT} or larger. Thus, L number violation is naturally induced by the presence of N_R , unless we enforce it by hand. Once we accept L violation we gain an elegant explanation for the smallness of neutrino masses as they turn out to be inversely proportional to the large scale where lepton number is violated.

1.2.2 Smallness of Neutrino Masses

Given that neutrino masses are extremely small, it is really difficult from the theory point of view to avoid the conclusion that L must be violated. In fact, it is only in terms of lepton number violation that the smallness of neutrino masses can be explained as inversely proportional to the very large scale where L is violated, of order M_{GUT} .

Assuming three neutrino flavours $\nu'_L = (\nu_e \ \nu_\mu \ \nu_\tau)^T$, consider the following two situations (in the neutrino interaction basis and after having diagonalized the charged lepton mass matrix).

- If L is not conserved, even in the absence of N_R , Majorana masses can be generated for neutrinos by allowing in the Lagrangian the presence of dimension five operators of the form

$$O_5 = -\frac{1}{2} \frac{(L^T \cdot \phi) \lambda (L \cdot \phi)}{M} \ni -\frac{1}{2} \frac{\nu'^T_L \lambda \nu'_L}{M} \langle \phi^0 \rangle \langle \phi^0 \rangle \quad (1.5)$$

with $\phi = (\phi^+ \ \phi^0)^T$ being the ordinary Higgs doublet, $L = (\nu'_L \ l_L)^T$ the weak lepton doublet, λ a matrix in flavour space and M a large scale ($\cdot$ is put to remember the weak isospin). Neutrino masses generated by O_5 are then of the order $m_\nu \sim v^2/M$ for $\lambda_{ij} \sim O(1)$, where $\langle \phi^0 \rangle = v \sim O(100 \text{ GeV})$ is the vacuum expectation value of the ordinary Higgs.

- The existence of a right handed neutrinos is quite plausible because all GUT groups larger than $SU(5)$ require them. In particular, the fact that a field with the same quantum numbers of a right handed neutrino completes the representation 16 of $SO(10)$: $16 = \bar{5} + 10 + 1$, so that all fermions of each family are contained in a single representation of the unifying group, is too impressive not to be significant. At least as a classification group $SO(10)$ must be of some relevance. Assuming that there are both $N_R = (N_e \ N_\mu \ N_\tau)^T$ and L violation, the see-saw mechanism [28] is possible. Consider the mass terms in the Lagrangian corresponding to Dirac and RR Majorana mass matrices (we consider LL Majorana mass terms as comparatively negligible) in the interaction basis in which charged leptons are diagonal:

$$\mathcal{L}_{seesaw} = -\overline{\nu'_L}^T m_D N_R + \frac{1}{2} N_R^T M_{RR} N_R + h.c. \quad (1.6)$$

The 3 by 3 matrices m_D and M_{RR} are the Dirac and Majorana mass matrices in flavour space (M_{RR} is symmetric, $M_{RR} = M_{RR}^T$, while m_D is, in general, non hermitian and non symmetric). One expects m_D to come from the Higgs mechanism: $\mathcal{L}_{Yuk} \ni -\overline{\nu'_L}^T y_D N_R \langle \phi^0 \rangle$. We expect the eigenvalues of M_{RR} to be of order M_{GUT} or more because RR Majorana masses are $SU(3) \times SU(2) \times U(1)$ invariant, hence unprotected and naturally of the order of the cut off of the low-energy theory. Since all N_R are very heavy we can integrate them away and the resulting effective neutrino mass matrix is:

$$\mathcal{L}_{seesaw} = -\frac{1}{2} \nu'^T_L m_\nu^{eff} \nu'_L + h.c. \quad (1.7)$$

with

$$m_\nu^{eff} = m_D M_{RR}^{-1} m_D^T. \quad (1.8)$$

This is the well known see-saw mechanism [28] result: the light neutrino masses are quadratic in the Dirac masses and inversely proportional to the large Majorana mass. If some N_R are massless or light they would not be integrated away but simply added to the light neutrinos.

Here we assumed that the additional non renormalizable terms from O_5 are comparatively negligible, otherwise they should simply be added. After elimination of the heavy right-handed fields, at the level of the effective low energy theory, the two types of terms are equivalent. In particular they have identical transformation properties under a chiral change of basis in flavour space. The difference is, however, that in the see-saw mechanism, the Dirac matrix m_D is presumably related to ordinary fermion masses because they are both generated by the Higgs mechanism

($m_D = y_D v$) and both must obey GUT-induced constraints. Thus, if we assume the see-saw mechanism more constraints are implied. In particular we are led to the natural hypothesis that m_D has a largely dominant third family eigenvalue in analogy to m_t , m_b and m_τ which are by far the largest masses among u quarks, d quarks and charged leptons. Once we accept that m_D is hierarchical it is very difficult to imagine that the effective light neutrino matrix, generated by the see-saw mechanism, could have eigenvalues very close in absolute value.

Then, assuming that one neutrino mass, presumably the third generation neutrino mass m_3 , is much larger than any other, we have $\Delta m_{atm}^2 \approx m_3^2$, which means $|m_3| \approx 0.05$ eV. If we apply the see-saw formula assuming that the Majorana RR masses lie around M , then, with $|m_3| \approx 0.05$ eV, $m_{D33} \sim 200$ GeV (like the top quark mass or the Higgs vev), we find $M \sim 0.8 \cdot 10^{15}$ GeV, a value amazingly close to M_{GUT} . This supports the initial assumption that indeed neutrino masses are related to the scale of L violation.

1.2.3 Approximate Mixing Matrix in the Neutrino Sector

Given the definition of the mixing matrix U in eq. (1.1) and the transformation properties of the effective light neutrino mass matrix $m_\nu^{eff} = m_\nu^{effT}$:

$$\begin{aligned} \nu_L'^T m_\nu^{eff} \nu_L' &= \nu_L^T (U^T m_\nu^{eff} U) \nu_L \\ U^T m_\nu^{eff} U &= \text{Diag}[e^{i\phi_1} m_1, e^{i\phi_2} m_2, m_3] \equiv m_{diag} \end{aligned} \quad (1.9)$$

where $m_{1,2,3}$ are real and positive, we obtain the general form of m_ν^{eff} :

$$m_\nu^{eff} = U m_{diag} U^T \quad . \quad (1.10)$$

The MNS matrix U can be parameterized in terms of three mixing angles and one phase, exactly as for the quark mixing matrix V_{CKM} .

$$U = \begin{bmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13} \\ -s_{12}c_{23}e^{i\delta} - c_{12}s_{13}s_{23} & c_{12}c_{23}e^{i\delta} - s_{12}s_{13}s_{23} & c_{13}s_{23} \\ s_{12}s_{23}e^{i\delta} - c_{12}s_{13}c_{23} & -c_{12}s_{23}e^{i\delta} - s_{12}s_{13}c_{23} & c_{13}c_{23} \end{bmatrix} \quad . \quad (1.11)$$

In addition we have the two phases ϕ_1 and ϕ_2 that are present because of the Majorana nature of neutrinos. Thus, in general, 9 parameters are added to the SM when non vanishing neutrino masses are included: 3 eigenvalues, 3 mixing angles and 3 CP violating phases [29, 30].

Maximal atmospheric neutrino mixing and the requirement that the electron neutrino does not participate in the atmospheric oscillations, as indicated by the

Super-Kamiokande and CHOOZ data, lead directly to consider as good approximation: $|U_{\mu 3}| = |U_{\tau 3}| = 1/\sqrt{2}$ and $U_{e3} = 0$. Hence, $c_{23} = s_{23} = 1/\sqrt{2}$ and $s_{13} = 0, c_{13} = 1$ (but different sign conventions are possible [31]) and the following structure of the U_{fi} ($f = e, \mu, \tau, i = 1, 2, 3$) mixing matrix is obtained:

$$U_{fi} = \begin{bmatrix} c_{12} & s_{12} & 0 \\ -s_{12}/\sqrt{2} & c_{12}/\sqrt{2} & 1/\sqrt{2} \\ s_{12}/\sqrt{2} & -c_{12}/\sqrt{2} & +1/\sqrt{2} \end{bmatrix} . \quad (1.12)$$

This result refers to the case of arbitrary solar mixing angle $s_{12} \equiv \sin \theta_{sol}$, so that $c_{12} = s_{12} = 1/\sqrt{2}$ holds for maximal solar mixing. In the limit $U_{e3} = 0$, all CP violating effects vanish and we can neglect the additional phase parameter δ generally present in U_{fi} : the matrix U is real and orthogonal and equal to the product of a rotation by $\pi/4$ in the 23 plane times a rotation in the 12 plane.

While the simple parameterization of the matrix U in eq. (1.12) is quite useful to guide the search for a realistic pattern of neutrino mass matrices, it should not be taken too literally. In particular, as stressed in section 1.1.4, the data do not exclude a non-vanishing U_{e3} element.

1.3 How to Reproduce the Large Atmospheric Mixing

1.3.1 Mass Hierarchies and Approximate Zeroth Order Textures

Note that since we are assuming only two frequencies, given by

$$\Delta_{sol} = m_2^2 - m_1^2, \quad \Delta_{atm} \cong m_3^2 - m_{1,2}^2 , \quad (1.13)$$

there are three possible hierarchies of mass eigenvalues [32, 31]:

$$\begin{aligned} \text{A} & : |m_3| \gg |m_{2,1}| \\ \text{B} & : |m_1| \sim |m_2| \gg |m_3| \\ \text{C} & : |m_1| \sim |m_2| \sim |m_3| \end{aligned} \quad (1.14)$$

(in case A there is no prejudice on the m_1, m_2 relation). For B and C different subcases are then generated according to the relative sign assignments for $m_{1,2,3}$.

Using eqs. (1.9) and (1.12), for each case, by setting to zero the small masses, it is possible to find [32, 33] the effective light neutrino matrices which automatically

lead to maximal atmospheric mixing ¹. Such zeroth order texture can be taken as guides in the construction of specific models for neutrino masses and mixings.

For example, in case A, $m_{diag} = \text{Diag}[0, 0, m_3]$ and we obtain

$$A : \quad m_{diag} = \text{Diag}[0, 0, 1] \, m_3 \quad \rightarrow \quad m_\nu^{eff} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & \frac{1}{2} & \frac{1}{2} \\ 0 & \frac{1}{2} & \frac{1}{2} \end{bmatrix} m_3 \quad . \quad (1.15)$$

In this particular case the results are the same for double and single maximal mixing. Note that the signs correspond to the phase convention adopted in eq. (1.12).

One example of case B follows by taking $m_{diag} = \text{Diag}[m_1, -m_1, 0]$ with double maximal mixing, so that:

$$B : \quad m_{diag} = \text{Diag}[1, -1, 0] \, m_1 \quad \rightarrow \quad m_\nu^{eff} = \begin{bmatrix} 0 & -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & 0 & 0 \\ \frac{1}{\sqrt{2}} & 0 & 0 \end{bmatrix} m_1 \quad . \quad (1.16)$$

A strong constraint arises in case C from the non observation of neutrino-less double beta decay which requires that the ee entry of m_ν^{eff} must obey $|(m_\nu)_{ee}| \leq 0.2$ eV [34]. As observed in ref. [35], if the average neutrino mass is around 1 eV, it can only be satisfied if bimixing is realized (that is double maximal mixing, with solar neutrinos explained by the VO solution or maybe by the large angle MSW solution). The reason is that, as results from eqs. (1.9) and (1.12), $(m_\nu)_{ee}^{eff} = m_1 c_{12}^2 + m_2 s_{12}^2$. For the required cancellation one needs opposite signs for m_1 and m_2 and comparable values of c_{12}^2 and s_{12}^2 , that is nearly maximal solar mixing. In this case, one example of case C is:

$$C : \quad m_{diag} = \text{Diag}[-1, 1, 1] \, m_1 \quad \rightarrow \quad m_\nu^{eff} = \begin{bmatrix} 0 & \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{2} & \frac{1}{2} \\ -\frac{1}{\sqrt{2}} & \frac{1}{2} & \frac{1}{2} \end{bmatrix} m_1 \quad . \quad (1.17)$$

It is possible to proceed similarly in all the other cases [33]. Once a solution to the solar neutrino problem is chosen and a set of suitable small perturbation terms is introduced, oscillation phenomena are unable to distinguish between the cases A, B and C. However, from the model building point of view, each texture represents an independent possibility in a zeroth order approximation.

¹Note that here we are working in the basis where the charged lepton masses are diagonal, and approximately given by $m_l = \text{Diag}[0, 0, m_\tau]$. For model building one has to arrange both the charged lepton and the neutrino mass matrices so that the neutrino results coincide with those given here after diagonalization of charged leptons.

1.3.2 Degenerate Neutrinos

Configurations B and C imply a very precise near degeneracy of squared masses. For instance, case C with $m_i \sim 1$ eV would require a relative splitting $|\Delta m/m| \sim \Delta m_{atm}^2/2m^2 \sim 10^{-(3\div 4)}$ and a much smaller one for solar neutrinos, especially if explained by the VO solution: $|\Delta m/m| \sim 10^{-(10\div 11)}$. Case C is the only one in which neutrinos could represent a significant component of hot dark matter. An intermediate possibility between the case of degenerate neutrino masses and the case of hierarchical ones is the situation *B* in eq. (1.14). With two almost degenerate heavier states and a nearly massless neutrino we cannot reach a mass range interesting for cosmological purposes. However, since the experimentally accessible quantities are the squared mass differences, case *B* remains an open possibility.

As mentioned above, it appears to be unplausible that starting from hierarchical Dirac matrices we end up via the see-saw mechanism into a nearly perfect degeneracy of squared masses. Thus models with degenerate neutrinos could only be natural if the dominant contributions directly arise from non renormalizable operators like O_5 in eq. (1.5) because they are *a priori* unrelated to other fermion mass terms. The degeneracy of neutrinos should be guaranteed by some slightly broken symmetry, which could also give the small terms required to go beyond the zeroth order approximation. Models based on discrete or continuous symmetries have been proposed [36].

Notice however that, for the VO scenario, even if arranged at the GUT scale, it is doubtful that such a precise degeneracy could be stable against renormalization group corrections when running down at low energy unless it is protected by a suitable symmetry [37, 38]. For instance, working only with the light neutrinos, the $U(1)_Q$ charge $Q \equiv (L_e - L_\mu - L_\tau)$ allows at leading order [39] for the texture of the example in (1.16). The non-vanishing entries $(m_\nu)_{12}$ and $(m_\nu)_{13}$ are expected to be of the same order, although not necessarily equal. The vanishing entries can be filled by the ratio of the vev $\langle \varphi \rangle$ of a scalar field carrying two units of Q and a mass scale M providing the cut off to the low-energy theory:

$$m_\nu = m \begin{bmatrix} \epsilon & 1 & 1 \\ 1 & \epsilon & \epsilon \\ 1 & \epsilon & \epsilon \end{bmatrix} \quad , \quad (1.18)$$

where $\epsilon = \langle \varphi \rangle / M$ and only the order-of-magnitudes are indicated. For values of ϵ smaller than 1 one obtains a small perturbation of the zeroth order texture. The mixing matrix has a large, not necessarily maximal, mixing angle in the 23 sector, a nearly maximal mixing angle in the 12 sector and a mixing angle of order ϵ in the 13 sector [40]. The mass parameter m should be close to $10^{-(1\div 2)}$ eV to provide

the frequency required by the atmospheric oscillations. Finally, $m_2^2 - m_1^2 \sim m^2 \epsilon$. In this model the VO solution to the solar neutrino deficit would require a very tiny breaking term, $\epsilon \sim \mathcal{O}(10^{-7})$. The symmetry protects the degeneracy of m_1 and m_2 against renormalization group corrections [38, 41].

This mechanism is a typical example of such models which are very attractive at first glance, but which, after a deeper analysis, turns out to be incomplete. In fact, the symmetry introduced is not able to account for the spectra of charged leptons - which, in this model, would be roughly degenerate - and of quarks. Thus, in order to construct a complete model of fermion masses, one has to do some further assumption. For instance one can introduce an ad hoc flavour symmetry which fulfills this task without damaging the effective neutrino mass matrix.

1.3.3 Hierarchical Neutrinos

We now discuss models of type A with large effective light neutrino mass splittings and large mixings. Reconciling large splittings with large mixing would seem difficult. Indeed, one could guess that, in analogy to what is observed for quarks, large splittings correspond to small mixings because only close-by states are strongly mixed. In particular it is of basilar importance to find a mechanism which leads to the large atmospheric mixing. Actually, there are mainly two ways in which this situation can be achieved.

Large Mixing from Stretching

The first one arises by stretching some parameters [42]. For instance, one could start with not too small 23 mixings both in the charged lepton and in the effective neutrino mass matrices. For instance, take $\theta_l \sim \theta_\nu \sim \theta_C \sim 13^\circ$ where θ_C is the Cabibbo angle. If the relative phases are such that the two angles just add, then from two not too small angles one can end up with a large one. Clearly the situation is not very satisfactory because the largeness of the mixing is completely accidental rather than descending as a consequence of some mechanism. In addition, one can expect at most the mixing to be large, but not at all to be exactly maximal.

Large Mixing from the See-Saw

Via the see-saw mechanism [28], there are other two particularly simple subcases in which a large atmospheric mixing can be realized. They are based on the observation that, in a 2 by 2 matrix context, the requirement of large splitting and large mixings

leads to a condition of vanishing determinant. For example the matrix

$$m \propto \begin{bmatrix} x^2 & x \\ x & 1 \end{bmatrix} \quad . \quad (1.19)$$

has eigenvalues 0 and $1 + x^2$ and for x of $O(1)$ the mixing is large. Thus, in the limit of neglecting small mass terms of order $m_{1,2}$, the demands of large atmospheric neutrino mixing and dominance of m_3 translate into the condition that the 2 by 2 subdeterminant 23 of the 3 by 3 mixing matrix vanishes. The problem is to show that this approximate vanishing can be arranged in a natural way without unacceptable fine tuning.

Without loss of generality, leaving aside for the moment the possible presence of flavour symmetries, one can go to the basis where both the charged lepton Dirac mass matrix m_l and the Majorana matrix M_{RR} of the right handed neutrinos are diagonal. For simplicity, we start assuming that the role of the first generation is not crucial in the mechanism for the generation of neutrino masses, so that one can, with good approximation, work in the 2 by 2 case (in the next section this condition will be relaxed). Writing m_D and M_{RR} , defined as in eq. (1.6), in the most general way

$$m_D = v \begin{bmatrix} a & b \\ c & 1 \end{bmatrix}, \quad M_{RR} = \begin{bmatrix} M_2 & 0 \\ 0 & M_3 \end{bmatrix} \quad , \quad (1.20)$$

where v is a vacuum expectation value, a, b and c are Yukawa couplings, then, via the see-saw, one obtains:

$$m_\nu^{eff} = \frac{v^2}{M_3} \begin{bmatrix} \frac{a^2}{M_2} + \frac{c^2}{M_3} & \frac{a b}{M_2} + \frac{c}{M_3} \\ \frac{a b}{M_2} + \frac{c}{M_3} & \frac{b^2}{M_2} + \frac{1}{M_3} \end{bmatrix} \quad . \quad (1.21)$$

The request of large splittings among the light neutrino eigenvalues is equivalent to demanding that the determinant of m_ν^{eff} is much smaller than its trace. It is then possible to see at glance that two cases arise exhibiting large mixing, that is when the terms with respectively M_3 or M_2 at the denominator are dominant.

- **Case I: Large mixing from m_D**

One simple example of the first case is realized if $M_2 \sim M_3$ and $a, b \ll 1$. In order to have a large splitting, one must have $c \sim 1$, that is the right handed neutrino of the third generation couples with the same strength [43] to left handed ν_μ and ν_τ . The heaviest mass for light neutrinos is then $m_3 \sim v^2/M_3$. As already pointed out, in the hierarchical case the data from Super-Kamiokande give $m_3 \sim 0.05$ eV so that, if one assumes that v is a typical weak

scale, namely 200 GeV, then $M_3 \sim 10^{15}$ GeV, just a bit lower than the GUT scale.

It is worth to stress that this first mechanism makes use of a *strongly asymmetric Dirac matrix*, that is a matrix with a large left handed mixing already present. It has been observed [32, 44, 45, 46, 47] that in $SU(5)$ left handed charged lepton mixings correspond to right handed mixings for down quarks (more on this later on). Large right handed quark mixings are not in contrast with experiment and viable GUT models that correctly reproduce the data on fermion masses and mixings can be constructed following this mechanism.

- **Case II: Large mixing from nothing**

An alternative possibility [48] is to have dominance of the terms with M_2 at the denominator. This is achieved for any $c < 1$ if $a^2, b^2 > M_2/M_3$. The request for large splitting is then equivalent to the requirement that also $a \sim b$. Now it is the second generation right handed neutrino which is particularly light and which couples with the same strength [43] to left handed ν_μ and ν_τ . In order to be more specific, consider the following example with symmetric matrices. These matrices are interesting because, for instance, one could want to preserve left right symmetry at the GUT scale. Then, the observed smallness of left handed quark mixings would also demand small right handed mixings. Starting from

$$m_D = v \begin{bmatrix} \epsilon & x\epsilon \\ x\epsilon & 1 \end{bmatrix}, \quad M_{RR}^{-1} = \frac{1}{M_3} \begin{bmatrix} r_2 & 0 \\ 0 & 1 \end{bmatrix} \quad , \quad (1.22)$$

where ϵ is a small number, x is of $O(1)$ and $r_2 \equiv M_3/M_2$, then, via the see-saw, it is sufficient that $\epsilon^2 r_2 \gg 1$ in order to have approximately:

$$m_\nu^{eff} = \frac{v^2}{M_3} \epsilon^2 r_2 \begin{bmatrix} 1 & x \\ x & x^2 \end{bmatrix} \quad . \quad (1.23)$$

The determinant is naturally vanishing so that the mass eigenvalues are widely split and for $x \sim 1$ the mixing is nearly maximal. It is exactly maximal if $x = 1$. The see-saw mechanism has created *large mixing from almost nothing*: all relevant matrices entering the see-saw are nearly diagonal [48, 49], that is they are diagonalized by transformations that go into the identity in the limit of vanishing ϵ . Clearly, the crucial factorization of the small parameter ϵ^2 only arises if the light Majorana eigenvalue is coupled to ν_μ and ν_τ with comparable strength, that is $x \sim 1$. An interesting feature of this second case, in connection with a possible realization within a GUT scheme, is that it

requires $M_3 > v^2/m_3$, so that one can push M_3 , the scale of L violation, beyond the GUT scale. This is a desirable feature because, for instance, it is expected in $SU(5)$ if right handed neutrinos are present and also in the breaking of $SO(10)$ to $SU(5)$. Summarizing, the second case require a peculiar hierarchy in the Majorana eigenvalues in order to work, but it has however the good characteristic of being realized with nearly diagonal matrices.

In the next chapter we present concrete examples which realize Case I and Case II in a 3 by 3 context, the desired textures being motivated by the introduction of a suitable flavour symmetry.

Chapter 2

Models For Hierarchical Neutrinos with Large Atmospheric Mixing

2.1 A Model with Large Mixing from m_D

By left handed mixing we mean non diagonal matrix elements that can only be eliminated by a large rotation of the left handed fields. Thus the question is how to reconcile large left handed mixings in the leptonic sector with the observed near diagonal form of V_{CKM} , the quark mixing matrix. Strictly speaking, since $V_{CKM} = U_u^\dagger U_d$, the individual matrices U_u and U_d need not be near diagonal, but V_{CKM} does, while the analogue for leptons, U of eq. (1.12), cannot be near diagonal. However nothing forbids for quarks that, in the basis where m_u is diagonal, the d quark matrix has large non diagonal terms that can be rotated away by a pure right handed rotation.

Right handed mixings for quarks can correspond to left handed mixings for leptons. In fact, in the context of SUSY $SU(5)$ [50] there is a very attractive hint of how this can be realized. In the $\bar{5}$ of $SU(5)$ the d^c singlet appears together with the lepton doublet (ν, e) . The (u, d) doublet and e^c belong to the 10 and ν^c to the 1 and similarly for the other families. As a consequence, in the simplest model with mass terms arising from only Higgs pentaplets, the Dirac matrix of down quarks is the transpose of the charged lepton matrix: $m_D^d = (m_D^l)^T$. Thus, indeed, a large mixing for right handed down quarks corresponds to a large left handed mixing for

charged leptons because, at leading order we may have ¹:

$$m_D^d = (m_D^l)^T = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 1 \end{bmatrix} v_d \quad (2.1)$$

In the same simplest approximation with 5 or $\bar{5}$ Higgs, the up quark mass matrix is symmetric, so that left and right mixing matrices are equal in this case. Then small mixings for up quarks and small left handed mixings for down quarks are sufficient to guarantee small V_{CKM} mixing angles even for large d quark right handed mixings. If these small mixings are neglected, we expect:

$$m_D^u = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} v_u \quad . \quad (2.2)$$

The crucial fact is that *when the charged lepton matrix is diagonalized the large left handed mixing of the charged leptons is transferred to the neutrinos*. To see this, note that in $SU(5)$ we can always diagonalize the u mass matrix by a rotation of the fields in the 10, the Majorana matrix M by a rotation of the 1 and the effective neutrino matrix m_ν by a rotation of the $\bar{5}$. In this basis the d quark mass matrix fixes V_{CKM} and the charged lepton mass matrix fixes U , the neutrino mixing matrix.

2.1.1 The Choice of Horizontal Abelian Charges

We give here an explicit example [32] of the mechanism under discussion in the framework of a unified SUSY $SU(5)$ theory with an additional $U(1)_F$ flavour symmetry [51]. This model is to be taken as merely indicative, in that some important problems, like, for example, the cancellation of chiral anomalies, are not tackled here. Most importantly, in the following minimal version, it has the unpleasant feature of predicting the same spectrum for down quarks and charged leptons. The problem of the differentiation of these spectra will be addressed in the following chapters. Despite this, we find it impressive that the general pattern of all what we know on fermion masses and mixings is correctly reproduced at the level of orders of magnitude. We regard the present model as a low-energy effective theory valid at energies close to $M_{GUT} \ll M_{Pl}$. We can think to obtain it by integrating out the heavy modes from an unknown underlying fundamental theory defined at an energy scale close to M_{Pl} . From this point of view the gauge anomalies generated by the

¹Here and in the following we write Dirac mass matrices with the convention $\bar{L}m_DR$.

light supermultiplets listed below can be compensated by another set of supermultiplets with masses above M_{GUT} , already eliminated from the low-energy theory. In particular, we assume that these additional supermultiplets are vector-like with respect to $SU(5)$ and chiral with respect to $U(1)_F$. Their masses are then naturally expected to be of the order of the $U(1)_F$ breaking scale, which, in the following discussion, turns out to be near M_{Pl} . It is possible to check explicitly the possibility of canceling the gauge anomalies in this way but, due to our ignorance about the fundamental theory, it is not particularly instructive to illustrate the details here.

In this model the known generations of quarks and leptons are contained in triplets Ψ_{10}^a and $\Psi_{\bar{5}}^a$, ($a = 1, 2, 3$) transforming as 10 and $\bar{5}$ of $SU(5)$, respectively. Three more $SU(5)$ singlets Ψ_1^a describe the right handed neutrinos. We start by discussing the Yukawa coupling allowed by $U(1)_F$ -neutral Higgs multiplets φ_5 and $\varphi_{\bar{5}}$ in the 5 and $\bar{5}$ $SU(5)$ representations and by the flavon X , singlet of $SU(5)$ with $F = -1$.

We assign to these fields the following F -charges:

field	Ψ_{10}	$\Psi_{\bar{5}}$	Ψ_1	X
F	(3, 2, 0)	(3, 0, 0)	(1, -1, 0)	-1

Table 2.1: F charge assignments.

2.1.2 Charged Fermion Mass Matrices

In the quark sector we obtain ²:

$$m_D^u = (m_D^u)^T = \begin{bmatrix} \lambda^6 & \lambda^5 & \lambda^3 \\ \lambda^5 & \lambda^4 & \lambda^2 \\ \lambda^3 & \lambda^2 & 1 \end{bmatrix} v_u \quad , \quad m_D^d = \begin{bmatrix} \lambda^6 & \lambda^3 & \lambda^3 \\ \lambda^5 & \lambda^2 & \lambda^2 \\ \lambda^3 & 1 & 1 \end{bmatrix} v_d \quad , \quad (2.3)$$

from which we get the order-of-magnitude relations:

$$\begin{aligned} m_u : m_c : m_t &= \lambda^6 : \lambda^4 : 1 \\ m_d : m_s : m_b &= \lambda^6 : \lambda^2 : 1 \end{aligned} \quad (2.4)$$

²In eq. (2.3) the entries denoted by 1 in m_D^u and m_D^d are not necessarily equal. As usual, such a notation allows for $O(1)$ deviations. Also remember that we write Dirac mass matrices with the convention $\bar{L}m_DR$.

and

$$V_{us} \sim \lambda, \quad V_{ub} \sim \lambda^3, \quad V_{cb} \sim \lambda^2. \quad (2.5)$$

Here $v_u \equiv \langle \varphi_5 \rangle$, $v_d \equiv \langle \varphi_{\bar{5}} \rangle$ and λ denotes the ratio between the vev of X and an ultraviolet cut off identified with the Planck mass M_{Pl} : $\lambda \equiv \langle X \rangle / M_{Pl}$. To correctly reproduce the observed quark mixing angles, we take λ of the order of the Cabibbo angle. The fermion spectrum depends on $\langle X \rangle$ and M_{Pl} only through the ratio λ . In chapter 4 we will see how λ can be dynamically generated by minimizing the X -dependent effective potential of the theory. For non-negative F -charges, the elements of the quark mixing matrix V_{CKM} depend only on the charge differences of the left handed quark doublet [51]. Up to a constant shift, this defines the choice in Table 2.1. Equal F -charges for $\Psi_{\bar{5}}^{2,3}$ are then required to fit m_b and m_s . We will comment on the lightest quark masses later on.

At this level, the mass matrix for the charged leptons is the transpose of m_D^d :

$$m_D^l = (m_D^d)^T \quad (2.6)$$

and we find:

$$m_e : m_\mu : m_\tau = \lambda^6 : \lambda^2 : 1 \quad (2.7)$$

The $O(1)$ off-diagonal entry of m_D^l gives rise to a large left handed mixing in the 23 block which corresponds to a large right handed mixing in the d mass matrix. Thus, this situation corresponds to Case I of section 1.3.3.

2.1.3 Neutrino Sector

Two different situations arise allowing or not for the presence of the additional anti-flavon field $\bar{X}$, with $F = 1$. Let us first consider what happens allowing for the pair X and $\bar{X}$.

Two Flavons

In the neutrino sector, the Dirac and Majorana mass matrices are given by:

$$m_D = \begin{bmatrix} \lambda^4 & \lambda^2 & \lambda^3 \\ \lambda & \lambda' & 1 \\ \lambda & \lambda' & 1 \end{bmatrix} v_u, \quad M = \begin{bmatrix} \lambda^2 & 1 & \lambda \\ 1 & \lambda'^2 & \lambda' \\ \lambda & \lambda' & 1 \end{bmatrix} \bar{M}, \quad (2.8)$$

where $\lambda' \equiv \langle \bar{X} \rangle / M_{Pl}$ and $\bar{M}$ denotes the large mass scale associated to the right handed neutrinos: $\bar{M} \gg v_{u,d}$.

After diagonalization of the charged lepton sector and after integrating out the heavy right handed neutrinos we obtain the following neutrino mass matrix in the low-energy effective theory:

$$m_\nu = \begin{bmatrix} \lambda^6 & \lambda^3 & \lambda^3 \\ \lambda^3 & 1 & 1 \\ \lambda^3 & 1 & 1 \end{bmatrix} \frac{v_u^2}{\bar{M}} \quad (2.9)$$

where we have taken $\lambda \sim \lambda'$. The $O(1)$ elements in the 23 block are produced by combining the large left handed mixing induced by the charged lepton sector and the large left handed mixing in m_D . A crucial property of m_ν is that, as a result of the see-saw mechanism and of the specific $U(1)_F$ charge assignment, the determinant of the 23 block is *automatically* of $O(\lambda^2)$.

It is easy to verify that the eigenvalues of m_ν satisfy the relations:

$$m_1 : m_2 : m_3 = \lambda^4 : \lambda^2 : 1 \quad . \quad (2.10)$$

The atmospheric neutrino oscillations require $m_3^2 \sim 10^{-3} \text{ eV}^2$. From eq. (2.9), taking $v_u \sim 250 \text{ GeV}$, the mass scale $\bar{M}$ of the heavy Majorana neutrinos turns out to be close to the unification scale, $\bar{M} \sim 10^{15} \text{ GeV}$. The squared mass difference between the lightest states is of $O(\lambda^4) m_3^2$, appropriate to the MSW solution to the solar neutrino problem. Finally, beyond the large mixing in the 23 sector, m_ν provides a mixing angle $s_{12} \sim (\lambda/2)$ in the 12 sector, close to the range preferred by the SMA-MSW solution. In general U_{e3} is non-vanishing, of $O(\lambda^3)$.

In general, the charge assignment under $U(1)_F$ allows for non-canonical kinetic terms that represent an additional source of mixing. Such terms are allowed by the underlying flavour symmetry and it would be unnatural to tune them to the canonical form. The results quoted up to now remain unchanged after including the effects related to the most general kinetic terms, via appropriate rotations and rescaling in the flavour space (see also ref. [52]).

Only One Flavon Field

A general problem common to all models dealing with flavour is that of recovering the correct vacuum structure by minimizing the effective potential of the theory. It may be noticed that the presence of two multiplets X and $\bar{X}$ with opposite F charges could hardly be reconciled, without adding extra structure to the model, with a large common vev for these fields, due to possible analytic terms of the kind $(X\bar{X})^n$ in the superpotential [53]. It is then interesting to explore the consequences of allowing only the negatively charged X field in the theory.

It can be immediately recognized that, while the quark mass matrices of eq. (2.3) are unchanged, in the neutrino sector the Dirac and Majorana matrices get modified into:

$$m_D = \begin{bmatrix} \lambda^4 & \lambda^2 & \lambda^3 \\ \lambda & 0 & 1 \\ \lambda & 0 & 1 \end{bmatrix} v_u \quad , \quad M = \begin{bmatrix} \lambda^2 & 1 & \lambda \\ 1 & 0 & 0 \\ \lambda & 0 & 1 \end{bmatrix} \bar{M} \quad . \quad (2.11)$$

The zeros are due to the analytic property of the superpotential that makes impossible to form the corresponding F invariant by using X alone. These zeros should not be taken literally, as they will be eventually filled by small terms coming, for instance, from the diagonalization of the charged lepton mass matrix and from the transformation that put the kinetic terms into canonical form. It is however interesting to work out, in first approximation, the case of exactly zero entries in m_D and M , when forbidden by F .

The neutrino mass matrix obtained via see-saw from m_D and M has the same pattern as the one displayed in eq. (2.9). A closer inspection reveals that the determinant of the 23 block is identically zero, independently from λ . This leads to the following pattern of masses:

$$m_1 : m_2 : m_3 = \lambda^3 : \lambda^3 : 1 \quad , \quad m_1^2 - m_2^2 = O(\lambda^9) m_3^2 \quad . \quad (2.12)$$

Moreover the mixing in the 12 sector is almost maximal:

$$\frac{s}{c} = \frac{\pi}{4} + O(\lambda^3) \quad . \quad (2.13)$$

For $\lambda \sim 0.2$, both the squared mass difference $(m_1^2 - m_2^2)/m_3^2$ and $\sin^2 2\theta_{sol}$ are remarkably close to the values required by the VO solution to the solar neutrino problem. This property remains reasonably stable against the perturbations induced by small terms (of order λ^5) replacing the zeros, coming from the diagonalization of the charged lepton sector and by the transformations that render the kinetic terms canonical.

We find quite interesting that also the VO solution, requiring an intriguingly small mass difference and a bimaximal mixing, can be reproduced, at least at the level of order of magnitudes, in the context of a "minimal" model of flavour compatible with SUSY $SU(5)$. In this case the role played by supersymmetry is essential, a non-supersymmetric model with X alone not being distinguishable from the version with both $\bar{X}$ and X , as far as low-energy flavour properties are concerned.

In chapter 4 we will again exploit this mechanism based on the presence of only one flavon X , but with slightly different charge assignments for the matter field $\Psi_{\bar{5}}$, so that the LOW solution to the solar neutrino problem will naturally emerge.

2.1.4 Outlook

We have constructed two examples which realize Case I of section 1.3.3. Obviously, the order of magnitude description offered by this model is not intended to account for all the details of fermion masses. Even neglecting the parameters associated with the CP violating observables, some of the relevant observables are somewhat marginally reproduced. For instance we obtain $m_u/m_t \sim \lambda^6$ which is perhaps too large. However we find it remarkable that in such a simple scheme most of the 12 independent fermion masses and the 6 mixing angles turn out to have the correct order of magnitude. Notice also that our model prefers large values of $\tan \beta \equiv v_u/v_d$. This is a consequence of the equality $F(\Psi_{10}^3) = F(\Psi_5^3)$. In this case the Yukawa couplings of top and bottom quarks are expected to be of the same order of magnitude, while the large m_t/m_b ratio is attributed to $v_u \gg v_d$ (there may be factors $O(1)$ modifying these considerations, of course). To keep $\tan \beta$ small, one could suppress m_b/m_t by adopting different F -charges for the Ψ_5^3 and Ψ_{10}^3 . Alternatively, as will be explicitly shown in chapter 4, one can assign F -charges also to the Higgs pentaplets.

Additional contributions to flavour changing processes (both in the quark and in the lepton [54] sectors) and to CP violating observables are generally expected in a SUSY GUT. However, a reliable estimate of the corresponding effects would require a much more detailed definition of the theory than attempted here. Crucial ingredients such as the mechanism of supersymmetry breaking and its transmission to the observable sector have been ignored. We are implicitly assuming that the omission of this aspect of the flavour problem does not substantially alter our discussion. For instance, it has been pointed out [55] that in $SU(5) \otimes U(1)_F$ based models with large $\tan \beta$ and radiatively induced flavour mixing, one generally expects the branching ratio of the flavour changing process $\mu \rightarrow e\gamma$ to be comparable to the present experimental upper bound (the exact values depending on the particular model considered). On the other hand, for $\tan \beta = O(1)$, the branching ratio turns out to be smaller by about four orders of magnitude.

A common problem of all $SU(5)$ unified theories based on a minimal Higgs structure is represented by the relation $m_D^l = (m_D^d)^T$ that, while leading to the successful $m_b = m_\tau$ boundary condition at the GUT scale, provides the wrong prediction $m_d/m_s = m_e/m_\mu$ (which, however, is an acceptable order of magnitude equality). In section 3.5, we will describe in some detail this problematic feature of minimal GUT models and briefly review the proposed solutions. An interesting possibility to overcome this problem and improve the picture of fermion masses will be adopted in the model of chapter 4, where we will exploit the presence of an additional supermultiplet transforming in the 75 representation of $SU(5)$.

2.2 A Model with Large Mixing from Nothing

Without loss of generality we can go to a basis where both the charged lepton Dirac mass matrix m_D^l and the Majorana matrix M for the right handed neutrinos are diagonal. In fact, after diagonalization of the charged lepton Dirac mass matrix, we still have the freedom of a change of basis for the right handed neutrino fields, in that the right handed charged lepton and neutrino fields, as opposed to left handed fields, are uncorrelated by the $SU(2) \times U(1)$ gauge symmetry. We can use this freedom to make the Majorana matrix diagonal: $M^{-1} = V^T d_M V$ with $d_M = \text{Diag}[1/M_1, 1/M_2, 1/M_3]$.

In order to understand how to generalize to the 3 by 3 context the mechanism already described in Case II of section 1.3.3 (see eq. (1.22)), it is useful to start by considering symmetric matrices. These matrices are very interesting because, for example, one could want to preserve left-right symmetry at the GUT scale. Then, the observed smallness of left handed mixings for quarks would also demand small right handed mixings.

One simple class of examples is the following one [48]. We start from

$$m_D = v \begin{bmatrix} \epsilon'' & \epsilon' & y \epsilon' \\ \epsilon' & \epsilon & x \epsilon \\ y \epsilon' & x \epsilon & 1 \end{bmatrix}, \quad M^{-1} = \frac{1}{\Lambda} \begin{bmatrix} r_1 & 0 & 0 \\ 0 & r_2 & 0 \\ 0 & 0 & r_3 \end{bmatrix}, \quad (2.14)$$

where, unless otherwise stated, x and y are $O(1)$; ϵ , ϵ' and ϵ'' are independent small numbers and $r_i \equiv M_3/M_i$. Note that the 23 blocks of these matrices closely match eq. (1.22). We expect $\epsilon'' \ll \epsilon' \ll \epsilon \ll 1$ and, perhaps, also $r_1 \gg r_2 \gg r_3 = 1$, if the hierarchy for right handed neutrinos follows the same pattern as for known fermions. Depending on the relative size of the ratios r_i/r_j , ϵ/ϵ' and ϵ'/ϵ'' , we can have models with dominance of any of the $r_{1,2}$.

For example, we set $x = 1$ (keeping y of $O(1)$) and assume $r_2 \epsilon^2 \gg r_1 \epsilon'^2, r_3$, together with $r_2 \epsilon'^2 \gg r_1 \epsilon''^2$ and $r_2 \epsilon \gg r_1 \epsilon''$. In this limit, with good accuracy we obtain:

$$m_\nu = \frac{v^2}{\Lambda} r_2 \epsilon^2 \begin{bmatrix} \frac{\epsilon'^2}{\epsilon^2} & \frac{\epsilon'}{\epsilon} & \frac{\epsilon'}{\epsilon} \\ \frac{\epsilon'}{\epsilon} & 1 + \frac{r_1 \epsilon'^2}{r_2 \epsilon^2} & 1 \\ \frac{\epsilon'}{\epsilon} & 1 & 1 + \frac{r_3}{r_2 \epsilon^2} \end{bmatrix}. \quad (2.15)$$

Note that, for $\epsilon \epsilon'' \leq \epsilon'^2$:

$$\text{Det}[m_\nu] = \left(\frac{v^2}{\Lambda} \right)^3 r_1 r_2 r_3 \epsilon'^4. \quad (2.16)$$

Approximate eigenvalues, in units of v^2/Λ and with $m_3 > m_2 > m_1$, are given by

$$m_3 \sim 2r_2\epsilon^2, \quad m_2 \sim \frac{1}{2}r_1\epsilon'^2, \quad m_1 \sim \frac{r_3\epsilon''^2}{\epsilon^2} \quad (2.17)$$

for $r_1\epsilon'^2 > r_3$ or, alternatively,

$$m_3 \sim 2r_2\epsilon^2, \quad m_2 \sim \frac{1}{2}r_3, \quad m_1 \sim \frac{r_1\epsilon'^4}{\epsilon^2} \quad (2.18)$$

for $r_1\epsilon'^2 < r_3$. Having set $x = 1$ the atmospheric neutrino mixing is nearly maximal. The solar neutrino mixing is instead generically small in these models, being proportional to ϵ'/ϵ . Thus the SMA-MSW solution is obtained. It is easy to find set of parameter values that lead to an acceptable phenomenology within these solutions. As an illustrative example we take:

$$\epsilon \sim \lambda^4, \quad \epsilon' \sim \lambda^6, \quad \epsilon'' \sim \lambda^{12}, \quad (2.19)$$

$$r_1 \sim \lambda^{-12}, \quad r_2 \sim \lambda^{-9}, \quad (2.20)$$

where $\lambda \sim \sin \theta_C$, θ_C being the Cabibbo angle. The neutrino mass matrices become

$$m_D = v \begin{bmatrix} \lambda^{12} & \lambda^6 & \lambda^6 \\ \lambda^6 & \lambda^4 & \lambda^4 \\ \lambda^6 & \lambda^4 & 1 \end{bmatrix}, \quad M = \Lambda \begin{bmatrix} \lambda^{12} & 0 & 0 \\ 0 & \lambda^9 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad (2.21)$$

and, in units of v^2/Λ , we obtain

$$m_3 \sim 1/\lambda, \quad m_2 \sim 1, \quad m_1 \sim \lambda^4. \quad (2.22)$$

The solar mixing angle θ_{12} is of order λ^2 , suitable to the SMA-MSW solution. Also $\theta_{13} \sim \lambda^2$.

It is also possible to arrange the parameters in eq. (2.14) in such a way that r_1 is dominant. Phenomenologically viable models can also be constructed in this case, always of the SMA-MSW type. For instance, we can take $y = 1$ (keeping x of $O(1)$) in order to obtain a large atmospheric mixing angle, and further assume the dominance of the r_1 terms, namely: $\epsilon'^2 r_1 \gg \epsilon'^2 r_2, r_3$, with $\epsilon''^2 r_1 \gg \epsilon'^2 r_2$ and $\epsilon'' r_1 \gg \epsilon r_2, r_3$. In this case we can approximate the light neutrino mass matrix as follows:

$$m_\nu = \frac{v^2}{\Lambda} r_1 \epsilon'^2 \begin{bmatrix} \frac{\epsilon''^2}{\epsilon'^2} & \frac{\epsilon''}{\epsilon'} & \frac{\epsilon''}{\epsilon'} \\ \frac{\epsilon''}{\epsilon'} & 1 + \frac{r_2 \epsilon^2}{r_1 \epsilon'^2} & 1 \\ \frac{\epsilon''}{\epsilon'} & 1 & 1 + \frac{r_3}{r_1 \epsilon'^2} \end{bmatrix}. \quad (2.23)$$

As an example we consider the following choice of parameters:

$$\epsilon \sim \lambda^4, \quad \epsilon' \sim \lambda^6, \quad \epsilon'' \sim \lambda^8, \quad (2.24)$$

$$r_1 \sim \lambda^{-14}, \quad r_2 \sim \lambda^{-6}. \quad (2.25)$$

In this case we have

$$m_D = v \begin{bmatrix} \lambda^8 & \lambda^6 & \lambda^6 \\ \lambda^6 & \lambda^4 & \lambda^4 \\ \lambda^6 & \lambda^4 & 1 \end{bmatrix}, \quad M = \Lambda \begin{bmatrix} \lambda^{14} & 0 & 0 \\ 0 & \lambda^6 & 0 \\ 0 & 0 & 1 \end{bmatrix}. \quad (2.26)$$

In units of v^2/Λ , we find

$$m_3 \sim 1/\lambda^2, \quad m_2 \sim 1 \gg m_1 \quad (2.27)$$

and $\theta_{12} \sim \lambda^2$.

As anticipated in section 1.3.3, an interesting feature of these textures, in connection with a possible realization within a GUT scheme, is that m_3 is much larger than v^2/Λ by construction. This means that the lepton number breaking scale Λ can be pushed up to the order of the GUT scale or even beyond. A large value of Λ , $\Lambda \geq M_{GUT}$, is naturally expected in $SU(5)$ if right handed neutrinos are present and in $SO(10)$ if the grand unified group is broken down to $SU(5)$ close to or slightly below the Planck mass. We recall that if m_3 is approximately given by v^2/Λ and $v \sim 250$ GeV, then $\Lambda \sim 10^{15}$ GeV.

Models based on symmetric matrices are directly compatible with left-right symmetry and therefore are naturally linked with $SO(10)$. This is to be confronted with models - like those described in the previous section - that have large right handed mixings for quarks, which, in $SU(5)$, can be naturally translated into large left handed mixings for leptons. In this connection it is interesting to observe that the proposed textures for the neutrino Dirac matrix can also work for up and down quarks. For example, the matrices

$$m_D^u \propto \begin{bmatrix} 0 & \lambda^6 & \lambda^6 \\ \lambda^6 & \lambda^4 & \lambda^4 \\ \lambda^6 & \lambda^4 & 1 \end{bmatrix}, \quad m_D^d \propto \begin{bmatrix} 0 & \lambda^3 & \lambda^3 \\ \lambda^3 & \lambda^2 & \lambda^2 \\ \lambda^3 & \lambda^2 & 1 \end{bmatrix}, \quad (2.28)$$

where for each entry the order of magnitude is specified in terms of $\lambda \sim \sin \theta_C$, lead to acceptable mass matrices and mixings. In fact $m_u : m_c : m_t = \lambda^8 : \lambda^4 : 1$ and $m_d : m_s : m_b = \lambda^4 : \lambda^2 : 1$. The V_{CKM} matrix receives a dominant contribution from

the down sector in that the up sector angles are much smaller than the down sector ones. The same kind of texture can also be adopted in the charged lepton sector ³.

It is simple to realize that models of this sort cannot be derived from a broken $U(1)_F$ horizontal symmetry, with a single field that breaks spontaneously $U(1)_F$. For symmetric matrices, left and right $U(1)$ charges must be equal: then, for example, if $m_{33} \sim l^{2q_3} \sim 1$ or $q_3 = 0$, then $m_{22} \sim l^{2q_2}$ is different from $m_{23} \sim l^{q_2}$ etc.

However, the symmetry requirement is not really necessary to produce a large atmospheric mixing angle from neutrino mass matrices characterized by small mixings. Indeed, by considering for simplicity the 2 by 2 case, instead of eq. (1.22), we could equally well have started from a Dirac mass matrix of the kind:

$$m_D \propto \begin{bmatrix} \epsilon & x' \\ x\epsilon & 1 \end{bmatrix} , \quad M^{-1} \propto \begin{bmatrix} r_2 & 0 \\ 0 & 1 \end{bmatrix} , \quad (2.29)$$

with x' any number smaller than or equal to one. Once the condition $\epsilon^2 r_2 \gg 1$ is satisfied, the neutrino mass matrix of eq. (1.23) is obtained again, independently from x' . The generalization of this mechanism to the full 3 by 3 case is straightforward, and analogous to the one discussed above for symmetric matrices. This leaves much more freedom for model building and does not necessarily bound the realization of these schemes to left-right-symmetric scenarios.

The solar mixing angle is generically small in the class of models explicitly discussed above. However, small mixing angles in the Dirac and Majorana neutrino mass matrices do not exclude a large solar mixing angle. For instance, this is generated from the asymmetric, but nearly diagonal mass matrices:

$$m_D = v \begin{bmatrix} \lambda^6 & 0 & 0 \\ \lambda^6 & \lambda^4 & 0 \\ 0 & \lambda^4 & 1 \end{bmatrix} , \quad M = \Lambda \begin{bmatrix} \lambda^{12} & 0 & 0 \\ 0 & \lambda^{10} & 0 \\ 0 & 0 & 1 \end{bmatrix} . \quad (2.30)$$

They give rise to a light neutrino mass matrix of the kind:

$$m_\nu = \begin{bmatrix} \lambda^2 & \lambda^2 & 0 \\ \lambda^2 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix} \frac{v^2}{\lambda^2 \Lambda} , \quad (2.31)$$

which is diagonalized by large θ_{12} and θ_{23} and small θ_{13} . The mass hierarchy is suitable to the LMA-MSW solution.

³Notice that the possibility of treating on equal footing all fermion mass matrices is also present in the framework of an extended flavour democracy. See for instance Ref. [56] and references therein.

2.2.1 Renormalization Effects

These results are stable under renormalization from the high-energy scale Λ where the mass matrices are produced down to the electroweak scale. Indeed, suppose that we start with a supersymmetric theory whose superpotential w at the scale $\Lambda \geq M_{GUT}$ is given by:

$$w = Qy^u U^c H_u + Qy^d D^c H_d + Ly^\nu N^c H_u + Ly^e E^c H_d + \frac{1}{2} N^c M N^c \quad , \quad (2.32)$$

where

$$y^e(\Lambda) = \begin{bmatrix} 0 & 0 \\ 0 & y_\tau \end{bmatrix} \quad , \quad y^\nu(\Lambda) = \begin{bmatrix} \epsilon & x\epsilon \\ x\epsilon & 1 \end{bmatrix} \quad , \quad M(\Lambda) = \begin{bmatrix} M_2 & 0 \\ 0 & M_3 \end{bmatrix} \quad . \quad (2.33)$$

Here we focus on the two heaviest generations. The matrices $y^{u,d,e,\nu}$ describe the Yukawa couplings of the theory, related to the mass matrices through $m_{u,\nu} = y^{u,\nu} v_u$ and $m_{d,e} = y^{d,e} v_d$, where $v_{u,d}$ are the vacuum expectation values of the neutral scalar components in the superdoublets $H_{u,d}$. The mass scales are ordered according to $\Lambda \geq M_3 \gg M_2 \gg M_W$, M_W denoting the electroweak scale. At the scale M_2 , after integrating out the superfields N^c associated to the heavy right handed neutrinos, the mass matrix for the light neutrinos, including the renormalization effects, is given by:

$$m_\nu(M_2) = \frac{v_u^2}{M_2} \begin{bmatrix} (y_{22}^\nu)^2 & y_{22}^\nu y_{32}^\nu \\ y_{22}^\nu y_{32}^\nu & (y_{32}^\nu)^2 \end{bmatrix} + O\left(\frac{1}{M_3}\right) \quad . \quad (2.34)$$

The couplings y_{22}^ν and y_{32}^ν are evaluated at the scale M_2 . If the running of M is computed by taking $\epsilon = 0$, then M_2 stays approximately constant when going from Λ down to M_2 . Notice that, if we neglect the terms of $O(1/M_3)$, the determinant of $m_\nu(M_2)$ vanishes, even including the renormalization effects. These effects modify the mixing angle at the scale M_2 and they depend on how the Yukawa coupling y_{22}^ν and y_{32}^ν change from Λ to M_2 . The running of the neutrino Yukawa couplings is governed by the equation:

$$\frac{dy^\nu}{dt} = \frac{1}{16\pi^2} \left[\text{tr}(3y^u y^{u\dagger} + y^\nu y^{\nu\dagger}) - 4\pi(3\alpha_2 + \frac{3}{5}\alpha_1) + 3y^\nu y^{\nu\dagger} + y^e y^{e\dagger} \right] y^\nu \quad , \quad (2.35)$$

from $\mu = \Lambda$ to $\mu = M_3$, with $t = \log \mu$. At M_3 , the superfield N_3^c is integrated out. This produces terms of order $1/M_3$, which are neglected in the present discussion. From $\mu = M_3$ down to $\mu = M_2$ the running of y_{22}^ν and y_{32}^ν is described by an equation formally identical to the previous one, with

$$y^\nu \equiv \begin{bmatrix} y_{22}^\nu & y_{23}^\nu \\ y_{32}^\nu & y_{33}^\nu \end{bmatrix} \rightarrow y^\nu \equiv \begin{bmatrix} y_{22}^\nu \\ y_{32}^\nu \end{bmatrix} \quad . \quad (2.36)$$

To assess the size of the renormalization effects on y'_{22} and y'_{32} , we have integrated numerically the above equations in the approximation of constant $y^{u,e}$ and $\alpha_{1,2}$. This should provide a rough estimate of the effects. As an example we take $\Lambda = 10^{18}$ GeV, $M_3 = 10^{16}$ GeV and $M_2 = 10^8$ GeV, $\alpha_2 = (5/3)\alpha_1 = 1/24$; we choose $y_\tau = 0.6$, corresponding to the large $\tan \beta \equiv v_u/v_d$ regime, which enhances the renormalization effects of y' . Assuming as initial conditions $y'_{22}(\Lambda) = y'_{23}(\Lambda) = y'_{32}(\Lambda) = 0.05$ and $y'_{33}(\Lambda) = 1$, we find that y'_{22} and y'_{32} are modified by about 10% at the scale M_2 . The mixing angle, nearly maximal at the scale Λ , remains close to maximal: $\sin^2 2\theta_{23} \sim 0.98$ in the example at hand.

The running from M_2 down to the electroweak scale does not appreciably affect neither the vanishing determinant condition, nor a nearly maximal neutrino mixing angle. This can be seen by analyzing the renormalization group equation for m_ν , which, in the one-loop approximation, is given by:

$$\frac{dm_\nu}{dt} = \frac{1}{8\pi^2} \left\{ \left[\text{tr}(3y^{u\dagger}y^u) - 4\pi(3\alpha_2 + \frac{3}{5}\alpha_1) \right] m_\nu + \frac{1}{2} [m_\nu y^{e\dagger}y^e + (y^{e\dagger}y^e)^T m_\nu] \right\} . \quad (2.37)$$

The solution is:

$$m_\nu(M_W) = C \begin{bmatrix} m_{\nu 22}(M_2) & m_{\nu 32}(M_2) B \\ m_{\nu 32}(M_2) B & m_{\nu 33}(M_2) B^2 \end{bmatrix} , \quad (2.38)$$

where

$$C = \exp \left[\int_{t_0}^t c(t') dt' \right] , \quad B = \exp \left[\int_{t_0}^t b(t') dt' \right] , \quad t_0 = \log(M_W) , \quad t = \log(M_2) \quad (2.39)$$

and

$$c(t) = \frac{1}{8\pi^2} \left[\text{tr}(3y^{u\dagger}y^u) - 4\pi(3\alpha_2 + \frac{3}{5}\alpha_1) \right] (t) , \quad (2.40)$$

$$b(t) = \frac{y_\tau^2(t)}{16\pi^2} . \quad (2.41)$$

We see from eq. (2.38) that if the determinant of m_ν vanishes at the scale M_2 , then it vanishes also at the weak scale. The mixing angle at the weak scale is given by:

$$\sin^2(2\theta)(M_W) = \frac{4m_{\nu 32}^2(M_2)B^2}{4m_{\nu 32}^2(M_2)B^2 + (m_{\nu 33}(M_2)B^2 - m_{\nu 22}(M_2))^2} . \quad (2.42)$$

The condition of maximal mixing at the scale M_2 is $m_{\nu 33}(M_2) = m_{\nu 22}(M_2)$. It is easy to see that, if this condition is met, then the first correction to $\sin^2(2\theta)(M_W)$ is

of second order in the parameter $(B - 1)$. Numerically, we find that $\sin^2(2\theta)(M_W)$ is reduced only by a of few percent.

Summarizing, the renormalization effects do not appreciably modify the results obtained at the tree level for eigenvalues and mixing angle of the neutrino mass matrix. They manifest themselves as $O(1)$ coefficients that, for all practical purposes, can be absorbed in the definition of the classical parameters. An analogous analysis can be carried out in the 3 by 3 case, with similar results. This is in agreement with the analysis of ref. [37], as far as the case of hierarchical neutrino masses is concerned.

2.2.2 An Explicit GUT Example with Broken Flavour Symmetry

From our general discussion it is clear that some amount of correlation between the Dirac and the Majorana sectors is required in order to reproduce, at the order-of-magnitude level, the observed pattern of oscillations (assuming from now on the SMA-MSW solution for solar neutrinos). In order to show that this correlation could be induced by an underlying flavour symmetry and, at the same time, to present an explicit grand unified example of the class of textures considered here, we [48] sketch the features of an $SU(5)$ model with a spontaneously broken flavour symmetry based on the gauge group $U(1)_A \times U(1)_B$.

The three generations are described by Ψ_{10}^a and $\Psi_{\bar{5}}^a$, ($a = 1, 2, 3$) transforming as 10 and $\bar{5}$ of $SU(5)$, respectively. Three more $SU(5)$ singlets Ψ_1^a describe the right handed neutrinos. Here we focus on the Yukawa coupling allowed by Higgs multiplets φ_5 and $\varphi_{\bar{5}}$ in the 5 and $\bar{5}$ $SU(5)$ representations and by three flavons, φ_A^- , φ_B^- and φ_B^+ , taken, to begin with, as $SU(5)$ singlets. The charge assignment of the various fields is summarized in Table 2.2. We assume that the flavon fields develop the following set of vevs:

$$\langle \varphi_A^- \rangle \sim \lambda \Lambda \quad , \quad \langle \varphi_B^- \rangle \sim \lambda \Lambda \quad , \quad \langle \varphi_B^+ \rangle \sim \Lambda \quad , \quad (2.43)$$

where Λ denotes the cutoff of the theory and provides the mass scale suppressing the higher-dimensional operators invariant under the gauge and the flavour groups.

In the quark sector we obtain ⁴

$$m_D^u = (m_D^u)^T = \begin{bmatrix} \lambda^8 & \lambda^6 & \lambda^4 \\ \lambda^6 & \lambda^4 & \lambda^2 \\ \lambda^4 & \lambda^2 & 1 \end{bmatrix} v_u \quad , \quad m_D^d = \begin{bmatrix} \lambda^6 & \lambda^4 & \lambda^3 \\ \lambda^5 & \lambda^3 & \lambda^2 \\ \lambda^3 & \lambda & 1 \end{bmatrix} v_d \quad , \quad (2.44)$$

⁴In eq. (2.44) the entries denoted by 1 in m_D^u and m_D^d are not necessarily equal. As usual, such a notation allows for $O(1)$ deviations.

	Ψ_{10}	$\Psi_{\bar{5}}$	Ψ_1	φ_5	$\varphi_{\bar{5}}$	φ_A^-	φ_B^-	φ_B^+
$SU(5)$	10	$\bar{5}$	1	5	$\bar{5}$	1	1	1
$U(1)_A$	(3,2,0)	(3,1,0)	(6,3,0)	0	0	-1	0	0
$U(1)_B$	(1,0,0)	(-2,-2,-1)	(0,2,0)	0	0	0	-1	+1

Table 2.2: Quantum numbers of matter and flavon multiplets.

from which we get the order-of-magnitude relations:

$$\begin{aligned} m_u : m_c : m_t &= \lambda^8 : \lambda^4 : 1 \\ m_d : m_s : m_b &= \lambda^6 : \lambda^3 : 1 \end{aligned} \quad (2.45)$$

and

$$V_{us} \sim \lambda, \quad V_{ub} \sim \lambda^3, \quad V_{cb} \sim \lambda^2. \quad (2.46)$$

Here $v_u \equiv \langle \varphi_5 \rangle$, $v_d \equiv \langle \varphi_{\bar{5}} \rangle$. Notice that V_{ub} and V_{us} are dominated by the down-quark contribution. The mass matrix for the charged leptons is the transpose of m_D^d :

$$m_D^l = (m_D^d)^T \quad (2.47)$$

and we find

$$m_e : m_\mu : m_\tau = \lambda^6 : \lambda^3 : 1. \quad (2.48)$$

At this level we obtain the well-known prediction $m_b = m_\tau$, together with the unsatisfactory relation $m_d/m_s = m_e/m_\mu$ (which, however, is an acceptable order of magnitude equality).

In the neutrino sector, the Dirac and Majorana mass matrices are given by:

$$m_D = \begin{bmatrix} \lambda^9 & \lambda^6 & \lambda^3 \\ \lambda^7 & \lambda^4 & \lambda \\ \lambda^6 & \lambda^4 & 1 \end{bmatrix} v_u, \quad M = \begin{bmatrix} \lambda^{12} & \lambda^{11} & \lambda^6 \\ \lambda^{11} & \lambda^{10} & \lambda^5 \\ \lambda^6 & \lambda^5 & 1 \end{bmatrix} \Lambda. \quad (2.49)$$

Notice that while in the model under consideration m_D is asymmetric, it is diagonalized by small unitarity transformations, i.e. transformations that go to the identity

in the limit of vanishing λ , in both the left and the right sectors. After diagonalization of the charged-lepton sector and after integrating out the heavy right handed neutrinos, we obtain the following neutrino mass matrix in the low-energy effective theory:

$$m_\nu = \begin{bmatrix} \lambda^4 & \lambda^2 & \lambda^2 \\ \lambda^2 & 1 & 1 \\ \lambda^2 & 1 & 1 \end{bmatrix} \frac{v_u^2}{\lambda^2 \Lambda} \quad . \quad (2.50)$$

The $O(1)$ elements in the 23 block are produced only by the interplay of the left handed mixing in m_D and the hierarchy in the Majorana sector. The contribution from the charged lepton sector is subleading. An important property of m_ν is that, as a result of the see-saw mechanism and of the specific $U(1)_F$ charge assignment, the determinant of the 23 block is of $O(\lambda^2)$ (in units of $(v_u^2/\lambda^2\Lambda)^2$). This gives rise to the desired hierarchy between atmospheric and solar frequencies. It is easy to verify that the eigenvalues of m_ν satisfy the relations:

$$m_1 : m_2 : m_3 = \lambda^8 : \lambda^2 : 1 \quad . \quad (2.51)$$

The atmospheric neutrino oscillations require $m_3^2 \sim 3.5 \times 10^{-3} \text{ eV}^2$. From eq. (2.50), taking $v_u \sim 250 \text{ GeV}$, the mass scale Λ of the heavy Majorana neutrinos turns out to be close to $2 \cdot 10^{16} \text{ GeV}$. The squared mass difference between the lightest states is of $O(\lambda^4) m_3^2$, appropriate to the MSW solution of the solar neutrino problem. Finally, beyond the large mixing in the 23 sector, m_ν provides a mixing angle $\theta_{12} \sim \lambda^2$ in the 12 sector, close to the range preferred by the SMA-MSW solution. In general U_{e3} is non-vanishing, of $O(\lambda^2)$. We find it encouraging that the general pattern of all that we know on fermion masses and mixings is correctly reproduced at the level of orders of magnitude.

This model is to be taken merely as an illustration, among many other possibilities, of the scenario outlined: small mixing angles for the fundamental fermions giving rise to a large mixing angle for the light neutrinos. For this reason we do not attempt to address some important problems, such as the cancellation of chiral anomalies, which we implicitly postpone to an energy scale higher than the unification scale. We have not dealt here with the problem of recovering the correct vacuum structure by minimizing the effective potential of the theory. Also, the order of magnitude description offered by this model is not intended to account for all the details of fermion masses. Even neglecting the parameters associated with the CP violating observables, some of the relevant observables are somewhat marginally reproduced. For instance, as already emphasized, a common problem of all $SU(5)$ unified theories, based on a minimal Higgs structure, is represented by the relation $m_D^l = (m_D^d)^T$, which, while leading to the successful $m_b = m_\tau$ boundary condition at the GUT scale, provides the wrong prediction $m_d/m_s = m_e/m_\mu$. We might

overcome this problem and improve the picture [57] by assuming that the flavon fields φ_A^- , $\varphi_B^\pm$ or a suitable subset of them transform in the adjoint representation of $SU(5)$. The product $(24)^q \bar{5}$ (where q is a positive integer) contains both the $\bar{5}$ and the $\overline{45}$ $SU(5)$ representations and the couplings would allow for a differentiation between the down quarks and the charged leptons. Again, since the purpose of the present model was only illustrative, we do not insist here on recovering fully realistic mass matrices. Finally, additional contributions to flavour-changing processes and to CP-violating observables are generally expected in a supersymmetric GUT. However, a reliable estimate of the corresponding effects would require a much more detailed definition of the theory than attempted here. Crucial ingredients such as the mechanism of supersymmetry breaking and its transmission to the observable sector have been ignored in the present note.

In this example we have given up a possible symmetric structure of the textures in favour of a relatively simple and economic realization of the underlying flavour symmetry. It is possible, within the $SU(5)$ or $SO(10)$ gauge theories, to provide slightly more elaborate examples where this symmetric structure is manifest. Here we have insisted on requiring both the Dirac matrices and the Majorana matrix M to be quasi-diagonal (that is with all off-diagonal entries ij suppressed with respect to the largest of the diagonal ii and jj ones). In some alternative models, all Dirac matrices are indeed quasi-diagonal, but large mixings are present in the Majorana matrix M . For instance, the idea of a large atmospheric mixing angle from intrinsically small mixings in the Dirac mass matrices has recently been advocated in the context of a $U(2)$ flavour theory [58]. In this model the Dirac neutrino mass matrix is the one displayed in eq. (2.14), with y of $O(\epsilon)$ (i.e. $(m_D)_{13} \ll (m_D)_{12}$) and $\epsilon'' \sim \epsilon'^2$. The $U(2)$ symmetry favours an r_1 dominance, by adequately suppressing the M_{11} element. However, to have both ν_μ and ν_τ with approximately the same coupling to the light Majorana mass eigenstate, this should possess a component of order ϵ' along the ν_3^c direction. In turn, this property is only exhibited if also the M_{33} entry is sufficiently small⁵. As a consequence, there are two heavy Majorana states with similar masses and there is a large mixing in the 23 Majorana sector. In this case, large mixings, which are absent from the Dirac mass matrices, arise from the Majorana sector, a physically similar possibility but technically different from that examined here.

⁵Technically, this suppression can be realized in the framework of an $SO(10)$ gauge group, by assuming that there are no flavour singlets that couple to $\nu_3^c \nu_3^c$.

2.2.3 Outlook

Summarizing, in most models that describe neutrino oscillations with nearly maximal mixings, there appear large mixings in at least one of the matrices m_D , m_D^l , M (i.e. the neutrino and charged-lepton Dirac matrices and the right-handed Majorana matrix). Here we have discussed the peculiar possibility that large neutrino mixing is only produced by the see-saw mechanism starting from all nearly diagonal matrices. Although this possibility is certainly rather special, we have shown that models of this sort can be constructed without an unrealistic amount of fine tuning and are well compatible with grand unification ideas and the related phenomenology for quark and lepton masses.

Chapter 3

Problems of SUSY GUTs

The idea that all particle interactions merge into a unified theory at very high energies [59] is so attractive, for both aesthetical and phenomenological reasons, that this concept has become widely accepted by now. Indeed, the quantitative success of coupling unification in SUSY Grand Unified Theories (GUTs) [60] can be hardly interpreted as just a coincidence. In addition, support has also come from the developments on neutrino oscillations [1], which point to lepton number violation at a scale close to the coupling unification one.

Despite the fact that they constitute promising ideas beyond the standard model, *SUSY GUTs are in a not so safe ground as at first sight could appear*. This chapter is devoted to a critical reanalysis of those problematic aspects found in constructing realistic models.

Assuming gauge unification, minimal models of GUTs based on $SU(5)$, $SO(10)$, ... have been considered in detail. Also, many articles have addressed particular aspects of GUTs models like proton decay, fermion masses and, recently, neutrino masses and mixings. In the previous chapter, indeed, the attention has focused on the phenomenology of neutrino masses and mixings. But on the one hand, *minimal models are not plausible as they need a large amount of fine tuning* and are therefore highly unnatural, for example with respect to the doublet-triplet splitting problem or proton decay. On the other hand, *analyses of particular aspects of GUTs often leave aside the problem of embedding the sector under discussion into a consistent whole*.

So the problem arises of going beyond minimal toy models by formulating sufficiently realistic, not unnecessarily complicated, relatively complete models that can serve as benchmarks to be compared with experiment. More appropriately, instead of "realistic" one should say "not grossly unrealistic" because it is clear that many important details cannot be sufficiently controlled and assumptions must be made.

Finally, the success or failure of such program can decide whether a stage of gauge unification is a likely possibility or not.

In this chapter we will review SUSY GUT's basic weak-points and analyze critically the current status of solutions. As will be seen, most of these solutions are partial, have some problematic feature and are in general quite complicated. The crucial problem is to find a consistent picture of the Higgs sector which leads to light doublets but triplets heavy enough to escape current bounds on proton decay [61]. In addition, the field content must not destroy coupling unification. During this survey, we will devote special attention to those solutions that will be adopted in the following chapter, where an example of semi-realistic model will be proposed.

3.1 The Doublet-Triplet Splitting Problem

The phenomenological motivation for weak scale SUSY is the solution of the hierarchy problem. While the mass of the Higgs doublet should be around 100 GeV, both coupling unification and non observation of higgsino mediated proton decay require the Higgs triplet mass to lie around 10^{16} GeV, the precise value depending on the details of the model under discussion. Achieving this doublet-triplet splitting [62] is the most compelling problem one has to face in GUTs, because it requires to fine-tune a parameter at the level of 14 orders of magnitude. It is worth to remind that, with respect to this problem, SUSY GUTs stay in a much better situation than non SUSY GUTs. Thanks to non-renormalization theorems, in the former the fine-tuning has to be done only once, that is at tree level, while in the latter it has to be done at all orders of the perturbative expansion.

After a short presentation of the transparent way in which the doublet-triplet splitting problem appears in the case of minimal $SU(5)$, this section has the purpose of summarizing the status of the most interesting mechanisms which avoids this fine-tuning also insisting on their weak-points. In many models, in fact, even if one succeeds in achieving the desired splitting at tree level, this could be upset by a lifting of the doublet mass, due to either radiative corrections or non renormalizable operators. In addition, one has to be careful because some of the possible mechanisms can destroy coupling unification.

3.1.1 The Doublet-Triplet Splitting Problem in Minimal $SU(5)$

In minimal SUSY $SU(5)$ the Higgs sector consists of one chiral superfields in the adjoint representation, Σ , and one pair of fundamental and antifundamental, $H, \bar{H}$.

The breaking of $SU(5)$ is provided by the following superpotential terms:

$$w_{Higgs} = M_{24} \Sigma_B^A \Sigma_A^B + \lambda \Sigma_B^A \Sigma_C^B \Sigma_A^C + c \overline{H}_A \Sigma_B^A H^B + M_5 \overline{H}_A H^A \quad , \quad (3.1)$$

where $A, B, \dots = 1, \dots, 5$ and summation over repeated indices has to be understood. In the limit of unbroken SUSY, minimization of this part of superpotential gives the following $SU(3) \otimes SU(2) \otimes U(1)$ symmetric vevs ¹:

$$\langle \Sigma \rangle = \frac{2}{3} \frac{M_{24}}{\lambda} \text{diag}(2, 2, 2, -3, -3) \quad , \quad (3.2)$$

$$\langle H \rangle = 0 \quad , \quad (3.3)$$

$$\langle \overline{H} \rangle = 0 \quad .$$

This superpotential also ensure heavy masses to those Σ multiplets which are not eaten by the Higgs mechanism (see Table 3.7). Due to the structure of the Σ vacuum, doublets and triplets acquire different masses:

$$M_D = M_5 - \frac{2c}{\lambda} M_{24} \quad , \quad M_T = M_5 + \frac{4c}{3\lambda} M_{24} \quad . \quad (3.4)$$

To have light doublets one has to impose that the parameters M_5 and M_{24} , which should assume values around 10^{16} GeV, satisfy

$$M_5 \cong \frac{2c}{\lambda} M_{24} \quad (3.5)$$

with an accuracy of at least the 14th decimal, so that $M_D = O(100\text{GeV})$. This hierarchy problem is inherent to all GUTs. SUSY doesn't explain the origin of the two scales, but renders this constraint stable against radiative corrections, so that, at least, the fine-tuning has to be done only once. Non SUSY GUTs, in contrast, need the tuning to be repeated at all perturbative orders.

Similar considerations also apply to the case of minimal $SO(10)$, where $H, \overline{H}$ lie in the 10 representations of $SO(10)$.

3.2 Solutions to the Doublet-Triplet Splitting Problem

The actual problem is to achieve the doublet-triplet splitting in a natural way, without fine-tuning. Some solutions that overcome this problem have been proposed,

¹Actually there are three supersymmetric degenerate minima: in one of them $SU(5)$ is unbroken, while in the other two it is broken respectively to $SU(4) \otimes U(1)$ and $SU(3) \otimes SU(2) \otimes U(1)$. The degeneracy could eventually be lifted and the MSSM vacuum chosen due to supergravity corrections.

based on the introduction of Higgs representations with useful group-theoretical properties, which do the job of allowing a mass only for triplets but not for doublets.

Several attempts have been done and someone of the most important will be summarized in the following. Despite the fact that basic ideas are indeed very elegant, emphasis will be given to some problems which remain unsolved. In addition, it turns out that embedding these solutions in a realistic and not too cumbersome framework is not an easy task.

3.2.1 The Missing Doublet Mechanism

One of the most economical solution the doublet-triplet splitting problem is represented by the Missing Doublet Mechanism (MDM) [63]. It works in $SU(5)$, which is the lower rank group containing the standard model, and is realized just replacing the 24 of Higgses by a 75, Y , and by further adding a pair of 50, $\overline{50}$ Higgses. Unfortunately, this proposal suffers of some weakness which, however, can be overcome.

Description of the Mechanism

The part of the superpotential responsible for the breaking of $SU(5)$ reads

$$w_{Higgs} = c_1 Y_{CD}^{AB} Y_{EF}^{CD} Y_{AB}^{EF} + M_Y Y_{CD}^{AB} Y_{AB}^{CD} \quad , \quad (3.6)$$

where, as usual, $A, B, \dots = 1, 2, \dots, 5$. These terms provide Y with a vev $\langle Y \rangle \equiv M_Y/(2c_1) \sim M_{GUT}$ and give a mass to all physical components of Y , i.e. those that are not absorbed by the Higgs mechanism (see Table 4.2). The SUSY vacuum of the 75 breaks $SU(5)$ uniquely to $SU(3) \otimes SU(2) \otimes U(1)$:

$$\langle Y_{CD}^{AB} \rangle = -\langle Y \rangle \left[\left(\delta_\gamma^\alpha \delta_\delta^\beta + 2\delta_c^a \delta_d^b - \frac{1}{2} \delta_C^A \delta_D^B \right) - (A \leftrightarrow B) \right] \quad , \quad (3.7)$$

where Greek indices assume values from 1 to 3 and Latin indices from 4 to 5.

The doublet-triplet splitting is realized by the following superpotential terms,

$$w_{MDM} = c_2 H^A Y_{BC}^{DE} H_{50}^{BCFG} \epsilon_{ADEF G} + c_3 \overline{H}_A Y_{DE}^{BC} H_{\overline{50}BCFG} \epsilon^{ADEF G} \\ + M_{50} H_{50}^{ABCD} H_{\overline{50}ABCD} \quad . \quad (3.8)$$

Note that the direct term $M_5 \overline{H}_A H^A$ must not to appear. The splitting occurs because the 50 of chiral superfields contains a $(\overline{3}, 1, 1/3)$, i.e. a coloured antitriplet, $SU(2)$ singlet (of electric charge 1/3) but no colourless doublet $(1, 2, -1/2)$. Denoting the colour triplets contained in H , $\overline{H}$, H_{50} , $H_{\overline{50}}$ by H_{3u} , H_{3d} , H'_{3d} and H'_{3u} respectively, after $SU(5)$ breaking:

$$w_{MDM} \ni -4\sqrt{3}c_2 \langle Y \rangle H'_{3d} H_{3u} - 4\sqrt{3}c_3 \langle Y \rangle H_{3d} H'_{3u} + M_{50} H'_{3d} H'_{3u} + \dots \quad (3.9)$$

where dots stand for mass terms for the other multiplets contained in $50, \overline{50}$. Thus, w_{MDM} doesn't produce any mass term for the doublets but, on the other hand, the Higgs colour triplets mix with the analogous states in the 50. The resulting triplet mass matrix is of the see-saw form:

$$\hat{m}_T = \begin{bmatrix} 0 & -4\sqrt{3}c_2\langle Y \rangle \\ -4\sqrt{3}c_3\langle Y \rangle & M_{50} \end{bmatrix} . \quad (3.10)$$

The eigenvalues of the matrix $\hat{m}_T\hat{m}_T^\dagger$ are the squares of:

$$m_{T1,2} = \frac{1}{2} \left[\sqrt{M_{50}^2 + 48 (c_2 + c_3)^2 \langle Y \rangle^2} \pm \sqrt{M_{50}^2 + 48 (c_2 - c_3)^2 \langle Y \rangle^2} \right] . \quad (3.11)$$

Note that $m_{T1}m_{T2} = 48 c_2 c_3 \langle Y \rangle^2$, so that triplets are naturally heavy. The effective mass that enters in the dimension 5 operators with $|\Delta B = 1|$ is the following combination of physical masses:

$$m_T = \frac{m_{T1}m_{T2}}{M_{50}} = \frac{48 c_2 c_3 \langle Y \rangle^2}{M_{50}} . \quad (3.12)$$

In the literature there is confusion [64, 65, 66, 67] on the numerical coefficients of the previous relations and also on the values for the heavy masses of the 75. In particular, many authors seem to forget the coefficient $4\sqrt{3}$ in eqs. (3.9), (3.10) and following. But this factor is important, because it gives in eq. (3.12) the large coefficient 48 that cannot be neglected when evaluating the effective triplet mass which suppress proton decay amplitudes. The formulae above have been rederived, by carefully paying attention to the group-theoretical factors. Appendix C contains a brief summary of this procedure.

Possible Criticisms

Criticism against the MDM is often moved by the common fear that its large rank representations are dangerous, mainly for two arguments. The first one is that, from the point of view of string theory, it is really hard to find examples containing such huge representations. From a more conservative point of view, however, embedding GUTs into a string theory is for sure interesting but not at all compelling. The second argument is that the large content of matter fields in the MDM causes the gauge coupling to blow up before M_{Pl} . In principle, a theory which is non perturbative before M_{Pl} is acceptable.

Apart from the two previous considerations, the MDM in its simplest version has mainly a weak point. No matter what symmetry is used to forbid the direct

mass term for $\overline{H}H$ in the superpotential, the doublet Higgs could take mass from non renormalizable operators of the form [68]:

$$\frac{\overline{H}HYY}{M_{Pl}} \quad . \quad (3.13)$$

If one believes Planck suppressed operators consistent with the theory are present, the doublet has a too big mass. The problem is exacerbated in MDM because the coupling blows up at a low scale and it is probably the associated strong scale which would suppress such operators. A promising way to overcome this obstacle was introduced in ref. [67] and is based on the introduction of an additional anomalous $U(1)$ flavour symmetry.

As will be clarified, the solution of the doublet-triplet splitting problem in terms of the 50, $\overline{50}$ and 75 representations is very suggestive because it automatically leads to improve the prediction of $\alpha_3(m_Z)$ and at the same time relaxes the constraints from proton decay.

3.2.2 The Sliding Singlet Mechanism

New possibilities emerge when introducing an additional singlet field S . The simpler one [69] is the Sliding Singlet Mechanism. It is realized by just adding to eq. (3.1) the term

$$fS\overline{H}_AH^A \quad . \quad (3.14)$$

where f is an adimensional coefficient. At tree level and considering also soft SUSY breaking terms, the vev of S can be arranged to cancel the large contributions to the doublet mass coming from the other terms. However, it is well known that this mechanism is unstable against radiative corrections [70], because tadpole diagrams induce the doublet to take a too big mass $\sim \sqrt{m_{3/2}M_{GUT}}$.

3.2.3 The Missing VEV Mechanism

The Missing VEV Mechanism [71] (MVM), which apply to the case of $SO(10)$, is based on the presence of a 45 Higgs representation. With respect to $SU(5)$, $45 = 1 + 24 + 10 + \overline{10}$, so that, in practice, having a singlet and a 24, it is based on the same idea of the Sliding Singlet. This can occur because the 45 representation is not constrained to be traceless. In fact, the 45 can have a vev of the form $\langle 45 \rangle \sim \text{diag}(1, 1, 1, 0, 0) \otimes i\sigma$. The term $10_1 45 10_2$, gives mass to triplets but not to doublets. Note that two 10 are required due to the antisymmetry of 45, so that there are four doublets and four triplets.

This mechanism seems very nice at first glance, but worrisome at second. In fact the problem arises of giving mass to just two doublets (or two linear combinations of them). This could be done, for instance, by introducing a term $M_2 10_2 10_2$ and forbidding any direct coupling with $10_1 10_1$. Heavy doublets have then a mass $\sim M_2$ and are in no way connected with the light ones. But there is also the need for triplets heavy enough to agree with experimental bounds on proton decay, and this is not very easy. For instance, by adding $M_2 10_2 10_2$, the triplets mass matrix $\hat{m}_T$ turns out to be of a see-saw form (exactly as in the case of the MDM but with the 45 instead of the 75 of $SU(5)$, see eq. (3.12)). The effective mass entering proton decay amplitudes is then proportional to $\langle 45 \rangle^2 / M_2$. But in order to achieve the needed suppression of proton decay amplitudes, we need $M_2 \leq 10^{-(2 \div 3)} \langle 45 \rangle \sim M_{GUT}$. Apart from the fact that the not-so-heavy doublets would give dangerously large threshold corrections, this requirement is clearly a fine-tuning.

The simplest versions of the MVM are then not realistic, but at the price of introducing a good deal of new fields, various models have been realized in the past in which the parameter playing the role of M_2 is suppressed because of symmetry reasons. Also the new fields produce large threshold corrections, so that one cannot exclude the accidental possibility of a general compensation of all their contributions.

However, dealing with a large number of GUT fields is not very attractive, because the predictivity of the model is highly reduced and, most importantly, the success of coupling unification within the MSSM would be merely a lucky coincidence. In addition, due to the increased number of Higgs representations that generally appears in many models, there is the problem of actually implementing a potential to get the desired minimum, paying attention to possible flat directions and without destabilizing the minimum with higher order terms. In fact, in order to lower the rank of $SO(10)$, one needs also one (or more) 16. But the couplings between 16 and 45 can destabilize the the desired vev of the 45. The literature is rich of interesting models [72, 73, 74] where this sort of compromise between the internal elegance and the necessity of agreeing with phenomenology appears clearly.

3.2.4 Higgses as pseudo-Goldstone Bosons

An interesting approach to the doublet-triplet splitting problem is obtained if one consider the Higgs doublets to be pseudo-Goldstone bosons related to a spontaneously broken accidental global symmetry of the Higgs sector. This distinguishes the doublets from the triplets in a very non trivial way. In this scenario the Higgs sector of theory does possess a spontaneously broken accidental global symmetry and the Higgs doublets are the associated Goldstone bosons. The accidental global symmetry is explicitly broken in the Yukawa sector. The Higgs doublets become

pseudo-Goldstone bosons with a mass determined by the scale of soft SUSY breaking terms. Because of non-renormalization theorems, this mass is protected against radiative corrections.

Despite that the idea is very elegant, the difficulty one finds in its actual realization is to motivate the accidental symmetry in a natural way. For instance, models involving a gauged $SU(5)$ symmetry and a global $SU(6)$ [75] unfortunately require an unpleasant fine-tuning of some parameters of the superpotential. In order to justify the accidental symmetry, an alternative possibility [76] is to assume that it arises because two sectors of the chiral superfields responsible for gauge symmetry breaking do not mix. But such a separation has to be imposed ad hoc.

3.3 Proton Decay

Proton decay is maybe the most interesting consequence of GUTs. Since a long time it has been representing a very hard challenge from the experimental point of view. The Kamiokande detector was constructed for this purpose and, after Super-Kamiokande, maybe a third-generation detector will be constructed. From the theoretical perspective this subject is not less engaging: the actual value of proton lifetime (if any) is determined by such an enormous amount of physics, from the GUT scale down to 1 GeV, that it offers the possibility of exploring various aspects of particle physics at the same time. The theoretical prediction is thus not at all accurate and strongly model dependent, but the common opinion is that it should lie just in the range already at hand of experimental searches or not very far.

A lot of weighty informations have already come from these studies:

- non SUSY $SU(5)$ is excluded because, due to the lightness of heavy gauge bosons, it predicts a lifetime induced by dimension 6 operators which is much lower than present bounds. Same conclusions hold for greater groups which break in one step to $SU(5)$, like $SO(10)$.
- minimal SUSY $SU(5)$ is in very serious troubles: with SUSY particles lighter than 1 TeV, dimension 5 operators would induce proton to decay mainly into $K^+\bar{\nu}$ at an exceedingly fast rate, in conflict with experiment. The case of $SO(10)$ is less definite, because the prediction of proton lifetime is linked to the particular Higgs content chosen to realize the breaking of $SO(10)$. However, it turns out that also in those models where as few as possible Higgs representations are introduced [42], the prediction of proton lifetime should not be far from the range explored by present searches.

Channel	Lower Limits (90 % CL) $\tau/B(10^{33}yr)$
$K^+\bar{\nu}$	1.9
$\pi^+\bar{\nu}$	0.025
$e^+\pi^0$	4.4
$\mu^+\pi^0$	3.3

Table 3.1: Lower Limits on Proton Decay Channels [77].

Prompted by the recent Super-Kamiokande improvements on proton partial lifetimes bounds [77], a renewed interest on this subject has arisen. Since GUTs without SUSY turned out to be disappealing, the attention has focused mainly on the question whether it is possible to construct examples of realistic SUSY GUT models. In such models, achieving a sufficient suppression of proton decay amplitudes from dimension 5 operators is of crucial importance. In fact, since the actual prediction of proton lifetime depends significantly of the details of the model, one is interested in arranging things in such a way to escape conflict with bounds.

To understand how this can be done reasonably, that is without introducing fine-tuning, it is useful to have a close look at the calculation of proton decay lifetime in the simplest possible case, namely minimal SUSY $SU(5)$. In this framework we will summarize the principal steps of the calculation of proton lifetime induced by dimension 5 operators [79]. We will treat independently Yukawa couplings of doublets and triplets. Also down quarks and charged leptons mass matrices will not be correlated, so that the calculation can be easily adapted also to non minimal models. We will show in some detail that, generically, the amplitude is dominated by only two terms of the Lagrangian and we will discuss, also quantitatively, how large can be the effect of the two CP violating phases (in addition to the CKM one) which enter this amplitude. We will comment on the source of the various uncertainties and give numerical examples. These have to be compared with the bounds collected in Table 3.1 from ref. [77].

3.3.1 Dimension Five Operators

The starting point is the most general renormalizable superpotential which could come from $SU(5)$. Yukawa couplings of doublets and triplets will be considered uncorrelated as well as down quarks and charged lepton mass matrices. The discussion can thus be adapted for any $SU(5)$ based model. The possible presence of any GUT superfield with vanishing vev will not alter the following discussion.

It is useful to write down the superpotential in terms of the MSSM fields. The triplets $H_{3u} \sim (3, 1, -1/3)$, $H_{3d} \sim (\bar{3}, 1, 1/3)$, together with the two light Higgs doublets H_{2u}, H_{2d} stay in a pair of $SU(5)$ pentaplets $H, \bar{H}$. Omitting colour and weak isospin indices, in the interaction basis the terms of the superpotential which are relevant to proton decay read:

$$\begin{aligned} w \ni & Q^T y_u U^c H_{2u} + Q^T y_d D^c H_{2d} + L^T y_e E^c H_{2d} \\ & + Q^T \hat{A} Q H_{3u} + U^{cT} \hat{B} E^c H_{3u} + Q^T \hat{C} L H_{3d} + U^{cT} \hat{D} D^c H_{3d} \\ & + m_T H_{3u} H_{3d} \quad , \end{aligned} \quad (3.15)$$

where Q, L, U^c, D^c and E^c denote as usual the chiral multiplets associated to the three fermion generations, $y_{u,d,e}, \hat{A}, \hat{B}, \dots$ are the GUT scale Yukawa matrices of doublets and triplets and m_T is the triplets mass.

By integrating out the colour triplets [78, 80, 81] one obtains the following effective superpotential

$$w_{eff} = \frac{1}{m_T} \left[Q \hat{A} Q Q \hat{C} L + U^c \hat{B} E^c U^c \hat{D} D^c \right] + \dots \quad , \quad (3.16)$$

dots standing for terms that do not violate baryon or lepton number. The former (latter) term, which involves four fields with only left (right) chirality, is referred to as LLLL (RRRR) operator. w_{eff} generates dimension 5 operators with two fermions and two sfermions in the Lagrangian.

At lower scales these dimension five operators give rise to the four-fermion operators relevant to proton decay, via a "dressing" process, that is through 1-loop diagrams with exchange of charginos and neutralinos. It turns out that the relevant contributions arise from chargino exchange [78, 79, 82]. When considering the operator $QQQL$, the most important dressing comes from wino exchange. There are four different terms in the Lagrangian, here called $O_i^L, (i = 1, \dots, 4)$, coming from the exchange of a wino ². Charged higgsino exchange provides instead the most important dressing of $U^c E^c U^c D^c$ [85]. Also in this case four Lagrangian terms arise,

²Gluino dressing contributions cancel among each other in case of degeneracy between first two generations of squarks [83, 84].

here called O_i^R , ($i = 1, \dots, 4$). The conventions adopted, the definition of the loop function f and all these operators are explicitly given in appendix A and in Table A.1.

3.3.2 Hunting Dominant Operators

The situation would then appear quite complicated. However, in the case of minimal $SU(5)$, where

$$-2\hat{A} = \hat{B} = y_u = y_u^T \quad , \quad -\hat{C} = \hat{D} = y_d = y_e^T \quad , \quad (3.17)$$

it turns out that, in each category, only one term gives the relevant contribution to proton decay amplitudes and that the contributions from the other three terms are suppressed by at least two powers of the Cabibbo angle, λ^2 . This is expected to be approximately true also in those models where the matrices $\hat{A}, \hat{B}, \dots$ do not differ significantly from those of the minimal case.

From eqs. (3.17), also the following relations follow

$$R_u^\dagger = L_u K \quad , \quad L_e = R_d^\dagger \quad , \quad R_e = L_d^\dagger \quad , \quad (3.18)$$

where $R_{u,d,e}, L_{u,d,e}$ are defined in Appendix A and K is a diagonal matrix of phases [86, 81] with determinant one. The two degree of freedom it contains are CP violating phases, in addition to the CKM one. Since proton decay amplitudes are insensitive to the overall phase, it is not restrictive to choose K to be of the following form

$$K = \text{diag}(e^{i\phi_1}, e^{i\phi_2}, 1) \quad . \quad (3.19)$$

In the mass eigenstates basis and making use of the previous definitions, the four-fermion operators of Table A.1 get greatly simplified. In fact, it turns out that the combinations in square brackets can be expressed just in terms of the diagonal Yukawa couplings of doublets and in terms of the CKM matrix, as shown in Table 3.2. This means that no trace is left of right handed mixings of fermions.

The relative importance of the various operators can be approximately understood expressing them in terms of powers of the Cabibbo angle $\theta_C = \lambda$. If the masses of the sparticles in the loop are such that they allow f (defined in appendix A) to vary in a small range, so that it can be quite safely factored out, and remembering the approximate GUT scale relations

$$\begin{aligned} m_u : m_c : m_t &= \lambda^8 : \lambda^4 : 1 \\ m_d : m_s : m_b &= \lambda^4 : \lambda^2 : 1 \end{aligned} \quad (3.20)$$

O_1^L	$[L_d^T(\hat{A} + \hat{A}^T)L_d]_{ij}[L_u^T\hat{C}L_e]_{kl} \rightarrow [V^T y_u^{(d)}K^+V]_{ij}[V^*y_d^{(d)}]_{kl} \equiv c_1^L{}_{ijkl}$
O_2^L	$[L_u^T(\hat{A} + \hat{A}^T)L_d]_{ij}[L_d^T\hat{C}L_e]_{kl} \rightarrow [y_u^{(d)}K^+V]_{ij}[y_d^{(d)}]_{kl} \equiv c_2^L{}_{ijkl}$
O_3^L	$[L_u^T(\hat{A} + \hat{A}^T)L_d]_{ij}[L_u^T\hat{C}L_e]_{kl} \rightarrow [y_u^{(d)}K^+V]_{ij}[V^*y_d^{(d)}]_{kl} \equiv c_3^L{}_{ijkl}$
O_4^L	$[L_u^T(\hat{A} + \hat{A}^T)L_u]_{ij}[L_d^T\hat{C}L_e]_{kl} \rightarrow [y_u^{(d)}K^+]_{ij}[y_d^{(d)}]_{kl} \equiv c_4^L{}_{ijkl}$
O_1^R	$[L_d^+y_u^*\hat{B}y_e^+L_e^*]_{kl}[R_u^*\hat{D}R_d^+]_{ij} \rightarrow [V^+(y_u^{(d)})^2Vy_d^{(d)}]_{kl}[KV^*y_d^{(d)}]_{ij} \equiv c_1^R{}_{klij}$
O_2^R	$[R_u^*\hat{B}y_e^+L_e^*]_{kl}[L_d^+y_u^*\hat{D}R_d^+]_{ij} \rightarrow [y_u^{(d)}Vy_d^{(d)}]_{kl}[V^+y_u^{(d)}KV^*y_d^{(d)}]_{ij} \equiv c_2^R{}_{klij}$
O_3^R	$[R_u^*\hat{B}R_e^+]_{kl}[L_d^+y_u^*\hat{D}y_d^+L_u^*]_{ij} \rightarrow [y_u^{(d)}V]_{kl}[V^+y_u^{(d)}KV^*(y_d^{(d)})^2V^T]_{ij} \equiv c_3^R{}_{klij}$
O_4^R	$[L_d^+y_u^*\hat{B}R_e^+]_{kl}[R_u^*\hat{D}y_d^+L_u^*]_{ij} \rightarrow [V^+(y_u^{(d)})^2V]_{kl}[KV^*(y_d^{(d)})^2V^T]_{ij} \equiv c_4^R{}_{klij}$

Table 3.2: In the case of minimal $SU(5)$, the combinations in square brackets of Table A.1 can be expressed in terms of the diagonal Yukawa couplings of doublets and in terms of the CKM matrix $V \equiv V_{CKM}$. See appendix A for the conventions adopted here.

$$V_{CKM} \sim \begin{bmatrix} 1 & \lambda & \lambda^3 \\ \lambda & 1 & \lambda^2 \\ \lambda^3 & \lambda^2 & 1 \end{bmatrix} \quad (3.21)$$

one immediately obtains Table 3.3.

One can see at glance that O_1^L, O_1^R are, within their groups, the dominant operators. The contributions to proton decay amplitudes from other operators are suppressed by at least λ^2 with respect to them, and can be safely neglected in the present estimate. The calculation then proceeds as summarized in Appendix A.

Generally, the dominant channel is $p \rightarrow K^+\bar{\nu}$. However, also $p \rightarrow \pi^+\bar{\nu}$ is important. In fact, the difference of a factor $\lambda \sim 0.25$ doesn't produce a factor 1/16 in rates. The π^+ channel is favoured by phase space for a factor of 2, as appears from Table A.2. Another factor 4 comes from Wick theorem. It results that π^+ is

	c_1^L	c_2^L	c_1^R	c_2^R
$p \rightarrow K^+ \bar{\nu}$	$\lambda^{10} \tan \beta$ ($\bar{\nu}_{\mu, \tau}$)	$\lambda^{12} \tan \beta$ ($\bar{\nu}_\mu$)	$\lambda^{10} \tan^2 \beta$ ($\bar{\nu}_\tau$)	$\lambda^{22} \tan^2 \beta$ ($\bar{\nu}_{\mu, \tau}$)
$p \rightarrow \pi^+ \bar{\nu}$	$\lambda^{11} \tan \beta$ ($\bar{\nu}_{\mu, \tau}$)	$\lambda^{14} \tan \beta$ ($\bar{\nu}_\mu$)	$\lambda^{11} \tan^2 \beta$ ($\bar{\nu}_\tau$)	$\lambda^{25} \tan^2 \beta$ ($\bar{\nu}_{\mu, \tau}$)
	c_3^L	c_4^L	c_3^R	c_4^R
$p \rightarrow K^0 l^+$	$\lambda^{14} \tan \beta$ (μ^+)	$\lambda^{12} \tan \beta$ (μ^+)	$\lambda^{17} \tan^2 \beta$ (e^+)	$\lambda^{14} \tan^2 \beta$ (μ^+)
$p \rightarrow \pi^0 l^+$	$\lambda^{13} \tan \beta$ (μ^+)	$\lambda^{14} \tan \beta$ (e^+)	$\lambda^{18} \tan^2 \beta$ (e^+)	$\lambda^{15} \tan^2 \beta$ (e^+)

Table 3.3: Relative importance of the combinations in square brackets appearing in Table 3.2, expressed in terms of powers of the Cabibbo angle $\theta_C = \lambda$. Note that we summed over neutrino flavours, which are experimentally indistinguishable.

suppressed by only a factor of 1/2 in rate, with respect to K^+ . As will be explained later, under certain circumstances linked to the two CP violating phases, the channel with π^+ could even be the dominant one.

As pointed out in [85], O_1^L, O_1^R can be comparable. For large $\tan \beta$ and/or $m_{\tilde{h}} \gg m_{\tilde{w}}$, O_1^R dominates. But large values of $\tan \beta$ are discouraging because they lead to faster proton decay, exacerbating a possible contrast with experiment. For small $\tan \beta$ and $m_{\tilde{h}} \leq m_{\tilde{w}}$, O_1^L gives the dominant contribution.

3.3.3 Proton Decay from Non-Renormalizable Operators

It has been pointed out [87] that also non-renormalizable operators can lead to proton decay. If one views GUTs as effective rather than fundamental theories, then one has to consider also the role of non-renormalizable operators. The following operators, where it is understood that we are taking light generations, are completely allowed by gauge symmetry,

$$w_{eff} \ni c_L \frac{QQQL}{\Lambda} + c_R \frac{U^c E^c U^c D^c}{\Lambda} \quad (3.22)$$

and lead, after the dressing process already described, to proton decay. Consider for instance the first term. It will be dressed as the LLLL operator arising from triplet exchange, so that, evaluating the ratio between its proton decay amplitude $\mathcal{A}_{nr}$ and the amplitude from heavy triplet exchange $\mathcal{A}_3$, loop functions cancel each other. The difference between the first terms of (3.16) and (3.22) resides only in their multiplicative coefficients and in the mass at denominator. Approximately, for the dominant channel $p \rightarrow K^+ \bar{\nu}$, with $\Lambda \sim M_{Pl}$, one finds

$$\frac{\mathcal{A}_{nr}}{\mathcal{A}_3} \sim \frac{c_L}{M_{Pl}} \frac{m_T}{\lambda^{10} \tan \beta} \quad (3.23)$$

For small $\tan \beta$ and $m_T \sim 2 \cdot 10^{16}$ GeV we have something like $\mathcal{A}_{nr}/\mathcal{A}_3 \sim c_L \lambda^{-7}$, which means a reduction of a factor $10^{8 \div 9}$ in lifetime for $c_L = O(1)$. With the former values of $\tan \beta$ and m_T , as we will see, the expectation for proton lifetime is roughly 10^{31} yr, which is lower than actual bounds. For non renormalizable operators one should then expect a lifetime of around 10^{22} yr. To restore agreement with the bounds, one has to impose c_L to be at least smaller than $\lambda^8 \sim 10^{-(5 \div 6)}$. This is indeed a very large fine-tuning. The most natural solution is to impose flavour symmetries, which could give a valid reason to justify the required smallness of c_L thus adequately suppressing these disturbing non renormalizable operators. Flavour symmetries are usually largely employed in model building, because they provide a clue to understand the observed picture of fermion masses and mixings. So, it would be interesting to study if, with the tool represented by flavour symmetries, one can solve both problems.

3.4 How to Prolong Proton Lifetime

There are several parameters entering the formula for proton lifetime in minimal $SU(5)$, so that one can should be careful not to underestimate theoretical errors.

For instance, by applying the chiral Lagrangian technique, one obtains the formulas for the calculation of proton decay rates which are given in Table A.2. One can see that the hadronic matrix elements $\alpha = -\beta$ are very important ingredients, because proton rate scale as their second power. Unfortunately, these quantities are not very well known. Commonly, their quoted range [88] is $0.03 \div 0.003 \text{ GeV}^3$ and the smallest value used in the calculations. However, it is worth to stress that with the largest value, the rate would be increased by roughly two orders of magnitude.

However, one could also construct elaborated models where one or more of the important parameters change with respect to the minimal model. So, it is of crucial importance to analyze how much the prediction for proton lifetime is affected if one moves a certain parameter in its possible range. In what follows we will briefly give a survey of the most interesting circumstances in which proton lifetime can be raised. Finally some numerical example will be provided. These are to be compared with those given in the next chapter, where a concrete model will be examined.

3.4.1 Altering Triplets Yukawa Couplings

Clearly, the previous discussion about dominant operators, apply only to those models where the matrices $\hat{A}, \hat{B}, \dots$ don't differ significantly from those of the minimal case. If, for some reason, such matrices are much more suppressed than in the minimal case, then proton decay rates would be reduced. This possibility has been explored [89] in the literature and, in general, the suppression of certain matrix elements follows from the group-theoretical construction of the model. One can also investigate if the same strategy employed to suppress the Yukawa couplings of triplets could also be effective in differentiating charged leptons and squarks spectra. Even if this is certainly an attractive possibility, it seems not very effective in increasing proton lifetime. Indeed, when implemented in a realistic framework, it is difficult, by only this mean, to adequately suppress all channels at the same time.

3.4.2 Raising the Triplet Mass

A really crucial parameter is the mass m_T . The dependence is strong, $\tau_p \sim m_T^{-4}$, so that a large value of m_T would drastically decrease the rates for all channels. As will be shown in the following, when considering the effect of heavy thresholds on the running of the gauge couplings, it follows that in minimal $SU(5)$ and with SUSY particles lighter than 1 TeV, m_T should not exceed 10^{16} GeV . This value causes proton decaying at a too fast rate, in conflict with experiment. Minimal SUSY $SU(5)$ is then in serious troubles. Interestingly, a large value for m_T is instead preferred in the MDM. Notice that in this case m_T is not a physical quantity but

an effective one, given by eq. (3.12), and could in principle even exceed M_{Pl} . In the next chapter we will discuss a realistic version of the MDM which maintains this attractive feature. On the contrary, in those realizations of the MDM and MVM where a large number of heavy representations lead to huge threshold corrections (often of both signs), the correlation between the value of m_T and the prediction for $\alpha_3(m_Z)$ is lost.

3.4.3 Moving the Two CP Violating Phases

There are phases entering the proton decay amplitudes [86] which don't affect any of the low energy observables, like for instance fermion masses and mixings³, but which play a really interesting role, because they can produce sizeable effects on proton rates. In fact, proton lifetime can easily span over one order of magnitude, just allowing the phases to vary in their range.

As usual, it is instructive to see this firstly in minimal $SU(5)$, and then comment on its extensions.

The phases of minimal $SU(5)$

In minimal $SU(5)$ the two phases can be clearly identified. Following the definition of eqs. (3.18) and (3.19), after a bit of work, one can see how they enter in the decay amplitude. In particular, considering the dominant operators O_1^L, O_1^R , one realizes that the dominant amplitudes are proportional to the following quantities

$$\begin{aligned}\mathcal{A}(K^+\bar{\nu}_\mu) &\propto (e^{i\phi_2}a_{\tilde{c}}^L + a_{\tilde{t}}^L) \\ \mathcal{A}(K^+\bar{\nu}_\tau) &\propto (e^{i\phi_2}a_{\tilde{c}}^L + a_{\tilde{t}}^L + e^{i\phi_1}a_{\tilde{t}^c}^R) \quad ,\end{aligned}\tag{3.24}$$

where $a_{\tilde{c}}^L, a_{\tilde{t}}^L$ and $a_{\tilde{t}^c}^R$ are complex numbers which contain the piece of amplitude coming respectively from $\tilde{c}$ and $\tilde{t}$ for wino dressing and from $\tilde{t}^c$ for higgsino dressing. $a_{\tilde{c}}^L, a_{\tilde{t}}^L$ are comparable, so that, depending on ϕ_2 , they can interfere constructively or destructively in the amplitude. One can have three possibilities. Firstly, if the contribution from higgsino exchange which enters in $\mathcal{A}(K^+\bar{\nu}_\tau)$ is smaller than the contribution from wino, a global suppression of the channel $p \rightarrow K^+\bar{\nu}$ can be achieved [82] for certain values of ϕ_2 . Note that, in this situation, the role of ϕ_1 is irrelevant. Secondly, if also $a_{\tilde{t}^c}^R$ is of the same order of $a_{\tilde{c}}^L, a_{\tilde{t}}^L$, also ϕ_1 , one can never achieve at the same time a suppression of both $\mathcal{A}(K^+\bar{\nu}_\mu)$ and $\mathcal{A}(K^+\bar{\nu}_\tau)$. Summing their corresponding rates, there isn't a sizeable variation of the total rate for

³In the diagonalization of the mass matrices, these phases can always be absorbed through suitable fields redefinitions, so that only one phase is left, the V_{CKM} one.

$p \rightarrow K^+ \bar{\nu}$ over the whole region of the space spanned by (ϕ_1, ϕ_2) . The third and last possibility is that a_{tc}^R dominates over the other contributions, in which case any effect of interference is totally lost. However this situation would require very large values of $\tan \beta$, leading to unacceptably fast proton decay.

In the first case we obtained a suppression of the main decay channel $p \rightarrow K^+ \bar{\nu}$. What about the channel $p \rightarrow \pi^+ \bar{\nu}$, for which the rate is in general expected to be just a factor of 2 smaller? In the first case, for those ϕ_2 values which produce destructive interference in the $K^+ \bar{\nu}$ rate, it becomes the dominant channel. In the second case, instead, it is always subdominant [85].

The phases in generic models

When constructing realistic $SU(5)$ models which break the relation $y_d = y_e^T$, then, in general, $\hat{A}, \hat{B}, \dots$ are no more linked to fermion masses. However, it is reasonable to expect their entries to be of the same order of magnitude, that is:

$$-2\hat{A} \sim \hat{B} \sim y_u \quad , \quad -\hat{C} \sim \hat{D} \sim y_d \sim y_e^T \quad . \quad (3.25)$$

In this situation one again finds the same results of the minimal case. In fact, the dominant amplitudes can be still written as

$$\begin{aligned} \mathcal{A}(K^+ \bar{\nu}_\mu) &\propto (b_{\tilde{c}}^L + b_{\tilde{t}}^L) \\ \mathcal{A}(K^+ \bar{\nu}_\tau) &\propto (b_{\tilde{c}}^L + b_{\tilde{t}}^L + b_{tc}^R) \quad , \end{aligned} \quad (3.26)$$

where $b_{\tilde{c}}^L, b_{\tilde{t}}^L$ and b_{tc}^R are complex numbers which contain the piece of the amplitude coming respectively from $\tilde{c}$ and $\tilde{t}$ for wino dressing and from $\tilde{t}^c$ for higgsino dressing. Thus, all the comments made when discussing eq. (3.24) still apply. For instance, when b_{tc}^R is not important, interference occurs depending on the relative phase between $b_{\tilde{c}}^L, b_{\tilde{t}}^L$.

3.4.4 Playing with SUSY Particles

In the proton decay amplitude the loop function f , defined in eq. (A.3), plays an important role. It is obtained by dressing the dimension 5 operators in order to convert them into four-fermions operators. This function depends on the masses of SUSY particles which run in the loop. To have an idea of the range of values in which f can lie, it is not a too bad approximation to assume that all squarks and sleptons are degenerate, with common mass m . In this case, the loop function for the exchange of the chargino $\tilde{c}$ of mass $m_{\tilde{c}}$ greatly simplifies

$$f_{\tilde{c}} = \frac{1}{16\pi^2} \frac{m_{\tilde{c}}}{(m^2 - m_{\tilde{c}}^2)} \left[m^2 - m_{\tilde{c}}^2 - m_{\tilde{c}}^2 \log \frac{m^2}{m_{\tilde{c}}^2} \right] \quad . \quad (3.27)$$

The behaviour of f is the following: by fixing m , it grows until $2m_{\tilde{e}}$ and then decreases; by fixing $m_{\tilde{e}}$, it grows when m decreases. In the limit $m_{\tilde{e}} \ll m$, then

$$f_{\tilde{e}} \propto \frac{m_{\tilde{e}}}{m^2} \quad . \quad (3.28)$$

This means that if sfermions are much more heavy than charginos, proton decay rates get greatly suppressed. However, it is not possible to push m towards values greater than 1 TeV, if one wants SUSY to be the solution of the hierarchy problem.

The hypothesis of degeneracy (or nearly degeneracy) of sfermions is useful because the various f can be factorized, that is, summing over $\bar{\nu}$ flavours, one can symbolically write:

$$\Gamma(p \rightarrow K^+ \bar{\nu}) \propto |f_{\tilde{w}} a_L + f_{\tilde{h}} a_R|^2 \quad (3.29)$$

where $a_{L,R}$ represent the contribution to amplitude coming respectively from wino and higgsino dressing. If $m_{\tilde{w}} = m_{\tilde{h}}$ the a_L contribution is the dominant one. Dominance of a_R occurs only if $m_{\tilde{w}} \ll m_{\tilde{h}}$ and/or $\tan\beta$ is large. To quantify the uncertainty on τ_p induced by the SUSY spectrum, one has to study the behaviour of f for reasonable values of $m_{\tilde{w},\tilde{h}}$ and sfermion masses. Considering typical SUGRA derived spectra with values of the soft breaking parameters in the range from 100 GeV to 1 TeV, one finds a variation of about one order of magnitude in rates.

3.4.5 Numerical Examples

The previous observations are clarified by the following example, where we compare two different set of textures for fermion masses which lead to exactly equal phenomenology, that is they both give the same masses and the same V_{CKM} .

The first set ⁴ is the well known Georgi-Jarlskog proposal [90]

$$y_u^{GJ} = \begin{bmatrix} 0 & a\lambda^6 & 0 \\ a\lambda^6 & 0 & b\lambda^2 \\ 0 & b\lambda^2 & 1 \end{bmatrix} \quad , \quad y_d^{GJ} = \begin{bmatrix} 0 & g\lambda^3 & 0 \\ g'\lambda^3 & j\lambda^2 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad . \quad (3.30)$$

For $\tan\beta = 3$ and with $\lambda = 0.23$, the values $a = 0.799$, $b = 0.813$ and $g = 0.005 + 0.372i$, $g' = 0.004 - 0.285i$, $j = 0.387$ give: $(m_u, m_c, m_t) = (0.0012, 0.28, 154)$ GeV, $(m_d, m_s, m_b) = (0.0009, 0.24, 1.17)$ GeV, $|V_{ud}| = 0.22$, $|V_{ub}| = 0.027$, $|V_{cb}| = 0.043$ and the Jarlskog invariant $I_G = 2.2 \cdot 10^{-5}$. These GUT scale values, when run to low energies, are compatible with the measured fermion masses and mixings. The

⁴In our convention Dirac mass terms are given by $\bar{L}mR$.

second set is obtained by replacing y_d^{GJ} with the following matrix possessing a large right-handed mixing in the 23 plane:

$$y_d^{HO} = \begin{bmatrix} 0 & A\lambda^3 & 0 \\ A^*\lambda^3 & 0 & B\lambda^2 \\ 0 & 1 & 1 \end{bmatrix} . \quad (3.31)$$

This particular form for the down quark mass matrix has been studied in detail by Hagiwara and Okamura [91]. For $\tan\beta = 3$ and $A = 0.274 + 0.475i$, $B = -0.217 - 0.711i$, one obtains exactly the previously quoted values of masses and mixings. These two set, due to the different values of CP violating phases, give different predictions for proton lifetime. Keeping $m_T = 2 \cdot 10^{16}$ GeV, $\alpha = -\beta = 0.003$ GeV³, $m_{\tilde{w}} = 250$ GeV, $m_{\tilde{h}} = 125$ GeV and assuming sfermions to be degenerate with $m_{\tilde{s}f} = 837$ GeV, then one predicts the values shown in Table 3.4.

	$\tau_p/B(p \rightarrow K^+\bar{\nu})$ $10^{31}yr$	$\tau_p/B(p \rightarrow \pi^+\bar{\nu})$ $10^{31}yr$
1 st set (GJ)	5.6	2.8
2 nd set (HO)	0.7	1.2

Table 3.4: Comparison of the predictions for proton lifetime between the two set of mass matrices.

The values of the various parameters adopted here are suitable for the case of minimal $SU(5)$. As pointed out, they suffer from large theoretical uncertainties. However one can recognize that minimal $SU(5)$ is in serious troubles, predicting a mean value for proton lifetime which is smaller than the present bound by two or three orders of magnitude.

In addition, one can also conclude that, even in the best constrained case of minimal $SU(5)$, results depend on the textures utilized, even if they give completely equivalent low energy phenomenology. The predictions differ for about one order of magnitude and also the main channel is different. This discrepancy has to be attributed entirely to the two CP violating phases. In fact, it is possible to recognize that the first case is an example of destructive interference for the amplitude of $K^+\bar{\nu}$ (see eq. (3.24)). The dominant mode is then $\pi^+\bar{\nu}$. In the second case, where

no destructive interference occurs, proton lifetime is smaller by about one order of magnitude and the mode $\pi^+\bar{\nu}$ is slightly subdominant. The result is that an uncertainty of about one order of magnitude arises just from the ignorance of the input textures.

One can wonder about the effect on rates induced by taking appropriate textures for y_e , and not just y_d^T , as required by minimal $SU(5)$. As will be discussed in the next section, one simple and well known possibility is to assume that certain elements of these matrices arise from a $\bar{45}$ of Higgses and not from a $\bar{5}$. In the former case, the element of y_e will be multiplied by a factor -3 with respect to the corresponding one of y_d^T . The matrices $\hat{C}, \hat{D}$, arising from the interaction of the $\bar{5}$ with matter, will have zero in the corresponding entry. For the first set, Georgi and Jarlskog pointed out that realistic Yukawa couplings for charged leptons can be obtained if the (22) element comes entirely from a $\bar{45}$, so that $m_e = 1/3 m_d, m_\mu = 3m_s, m_\tau = m_b$. For the second set, these relations can be achieved if $y_{d2,3}$ comes from a $\bar{45}$:

$$y_e^{GJ} = \begin{bmatrix} 0 & g'\lambda^3 & 0 \\ g\lambda^3 & -3j\lambda^2 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \quad y_e^{HO} = \begin{bmatrix} 0 & A^*\lambda^3 & 0 \\ A\lambda^3 & 0 & 1 \\ 0 & -3B\lambda^2 & 1 \end{bmatrix}. \quad (3.32)$$

Correspondingly, $\hat{C}_{22}^{GJ} = \hat{D}_{22}^{GJ} = 0$ and $\hat{C}_{23}^{HO} = \hat{D}_{23}^{HO} = 0$. It is easy to adapt the calculation of proton decay lifetime to this particular case. Under the same assumptions made to obtain Table 3.4 one now obtains Table 3.5.

	$\tau_p/B(p \rightarrow K^+\bar{\nu})$ $10^{31}yr$	$\tau_p/B(p \rightarrow \pi^+\bar{\nu})$ $10^{31}yr$
1 st set (GJ)	5.8	2.9
2 nd set (HO)	0.8	1.5

Table 3.5: Comparison of the predictions for proton lifetime between the two sets of mass matrices in the realistic case.

The results are very similar to those of the minimal case. Indeed this is a simple and very particular example, however, it is true also in general that the use of simple realistic textures also for charged leptons doesn't have very sizeable effects on proton

lifetime. Achieving a certain degree of suppression through a differentiation between triplet and doublet Yukawa couplings is a technically realizable possibility only for much more complicated models, where a number of ad hoc assumptions must be made [89]. It is clear that one is interested in looking for such mechanisms that lead to construct non minimal models where proton lifetime can be pushed comfortably above experimental bounds. As pointed out, a resolute way to do this, would be to have triplets much more heavy than in the minimal case. In the next chapter we will discuss one of these models.

3.5 Fermion Masses and Mixings

The aesthetically attractive fact of putting in the same representation conceptually different objects like quarks and leptons has mainly two consequences. The most apparent one is the fact that quarks and leptons mass matrices turn out to be strictly correlated. Despite the difficulties found, this can give interesting starting points to face the flavour problem. The second consequence is the possibility of violating baryon and lepton numbers. As shown, this leads to the striking prediction of proton lifetime, with values in the region already partially excluded by present bounds. Minimal models fail in correctly predicting both mass relations and proton lifetime.

In the following we will review the usual methods which can be adopted to share some light on the resolution of the first problem. It is clear that finally one is interested in understanding if these mechanisms are compatible or even help in the solution of the other problems like proton decay and the other typical GUT problems.

3.5.1 Mass Relations

The evident problem of minimal models is the prediction of wrong mass relation for down quarks and charged leptons at M_{GUT} . Assuming an exact R -parity discrete symmetry under which matter fields are odd and $H, \overline{H}$ are even, if the superpotential contains the following Yukawa interaction terms

$$w_{Yuk} = \frac{1}{4} \Psi_{10}^{AB} G_u \Psi_{10}^{CD} H^E \epsilon_{ABCDE} + \sqrt{2} \Psi_{10}^{AB} G_d \Psi_{\overline{5}A} \overline{H}_B \quad (3.33)$$

where $A, B, \dots = 1, \dots, 5$, $H, \overline{H}$ are the pentaplets which contain the pair of light doublets and G_u, G_d are 3×3 matrices in flavour space. Writing w_{Yuk} in terms of MSSM multiplets, the interactions with doublets read:

$$w_{Yuk} \ni Q^T G_u U^c H_{2u} + Q^T G_d D^c H_{2d} + L^T G_e E^c H_{2d} \quad . \quad (3.34)$$

Then we get the following Dirac mass matrices at M_{GUT} :

$$\hat{m}_u = G_u \frac{v_u}{\sqrt{2}} \quad , \quad \hat{m}_e^T = \hat{m}_d = G_d \frac{v_d}{\sqrt{2}} \quad , \quad (3.35)$$

where v_u, v_d parameterize the vevs of $H, \overline{H}$. The sharp transposition relation $\hat{m}_e^T = \hat{m}_d$ is in conflict with what is expected on phenomenological grounds, by evolving the values of down quarks and charged lepton masses from their values measured at the electroweak scale to the GU scale, i.e. the Georgi-Jarlskog relations:

$$m_e \cong \frac{1}{3}m_d \quad , \quad m_\mu \cong 3m_s \quad , \quad m_\tau \cong m_b \quad . \quad (3.36)$$

This problem is even exacerbated in minimal $SO(10)$, where $\hat{m}_u = \hat{m}_u^T = \hat{m}_{\nu D} \propto \hat{m}_d = \hat{m}_d^T = \hat{m}_e$. However, at least in $SO(10)$ there is naturally the place for right-handed neutrinos.

On the contrary, in $SU(5)$ the inclusion of neutrino masses represents an additional problem. A very simple way out is the addition of a gauge singlet field Ψ_1 , which contains right-handed neutrinos. If one views $SU(5)$ as descending from $SO(10)$, this is of course not only natural but also necessary. Then, in the superpotential these terms are allowed

$$\Psi_1 G_\nu \Psi_{\bar{5}A} H^A - \frac{1}{2} M \Psi_1 G_M \Psi_1 \quad (3.37)$$

which realize the see-saw mechanism. If M is just a bit smaller than M_{GUT} , then the atmospheric neutrino mass scale is naturally obtained. However one should emphasize that, because the term which gives mass to right-handed neutrinos doesn't break $SU(5)$, then M should in principle be expected of order of the cut-off of the theory, that could even be as large as M_{Pl} . The advantage of groups like $SO(10), SU(4) \otimes SU(2) \otimes SU(2)$ is that they directly possess a particle with the same quantum numbers of right-handed neutrinos, whose mass scale thus results for dynamical reasons of order of the GUT scale.

3.5.2 How to Achieve Realistic Spectra

The flavour problem consists in explaining the origin of the difference between the three fermionic families, difference which causes their masses and mixings to be organized in strongly and bizarre hierarchical structures. Probably this problem is linked to the more fundamental one: why just three families? It is clear that the idea of GU cannot alone answer these deep questions. The only thing it can offer is a much more constrained framework with respect to the MSSM, where one can

look for the criterion according to which the different mass matrices are linked one to each other. One can thus learn a lot about what would be allowed to be present at scales higher than the electroweak one and what could not at all be there.

To understand the situation it is convenient to take $SU(5)$ as reference. This is not at all restrictive because $SO(10)$ can be always thought in terms of its $SU(5)$ content. The problems one has to face are two: firstly a mechanism to produce a progressive suppression in the masses of the three generation, secondly a way to reproduce the correct values of masses and mixings. The usual road taken to overcome the first problem is by the use of flavour symmetries [51], for instance a continuous $U(1)$. This strategy is very successful, even if one is left with the discomfort of having done ad hoc assumptions for the charges. Apart from this, it turns out that it is very difficult to realize the correct spectra without tuning some parameter, even in the more complicated extensions of minimal GUT models.

Possible ways for accounting for neutrino masses and mixings have been already discussed. Here we will review the main strategies one can use to break the bad transposition relation between down quarks and charged leptons. Since $10 \times \bar{5} = 5 + 45$, a mass term for quarks and leptons can be realized only with a $\bar{5}$ and a $\bar{45}$. The $\bar{5}$, which is of course nothing but the pentaplet containing one of the light Higgs doublets, predicts the sharp transposition relation. So, to have a realistic spectrum there is the absolute need for a $\bar{45}$, which succeeds in breaking the unwanted transposition. The $\bar{45}$ can be a "true" or an effective one. The first scenario, with a true $\bar{45}$, was proposed a long time ago. Now it seems not very attractive because the true $\bar{45}$ contains a true doublet, which needs to be massive. One can then introduce another 45 to construct a direct mass term (and also to cancel anomalies). But it is well known that the two additional doublets cannot be light. If this were the case then the prediction for $\alpha_3(m_Z)$ assuming coupling unification would be given by:

$$\alpha_3(m_Z) = \alpha_{em}(m_Z) \frac{56 - 2\delta n_H}{(120 + 6\delta n_H)\sin^2\theta(m_Z) - (24 + 3\delta n_H)} \quad (3.38)$$

where δn_H is the number of light extra doublets, in addition to the two of the MSSM. Taking $\alpha_{em}(m_Z) = 1/128$ and $\sin^2\theta(m_Z) = 0.23$, one has:

$$\alpha_3(m_Z) = 0.12 \quad \delta n_H = 0 \quad (\text{MSSM}) \quad (3.39)$$

$$\alpha_3(m_Z) = 1.13 \quad \delta n_H = 2 \quad . \quad (3.40)$$

It is then absolutely unavoidable: to give a GUT scale mass to the four triplets; to maintain at the electroweak scale two linear combinations of the doublets in $\bar{5}$, $\bar{45}$ and 5 , 45 but raising at the GUT scale their two orthogonal combinations. Here come the difficulties, because this asymmetric splitting between doublets and triplets is a much more complicated version of the usual doublet triplet splitting problem.

To escape from these difficulties one can make complete use of the possibility of working with non renormalizable operators and explore the field offered by the effective 45. The simplest possible mechanism is the well known proposal made by Ellis and Gaillard [57] of considering the 45 of the product $\bar{5} \times 24 = \bar{5} + 45 + 70$. This is indeed a very economical solution, because the $\bar{5}$ is already there and the 24, which must have non vanishing vev, can be naturally the one responsible for the breaking of $SU(5)$. Thus, this mechanism uses just the field content of minimal $SU(5)$. The following operator

$$\sqrt{2} \Psi_{10\ i}^{AB} G_{dij} \Psi_{\bar{5}Cj} \overline{H}_B \frac{\langle \Sigma_A^C \rangle}{\Lambda} \quad (3.41)$$

where Λ is the cut-off, gives $G_{dij} = y_{dij} = -2/3y_{eji}$. Note that both channels $\bar{5}$ and 45 of the product $\bar{5} \times 24$ contribute to the mass. Also operators with more insertions of Σ can be useful in realistic examples. The number of the invariants obtained by contracting the indices in all possible ways grows with the number of insertions made, so that one can use the freedom in choosing their relative coefficient to fit the masses. This is however a kind of proof that realistic mass spectra are realizable rather than a genuine prediction of the model.

There are other possibilities, in addition to the use of a 24. In the MDM, for instance, the 75 responsible for the breaking of $SU(5)$ can be successfully well employed, without the introduction of additional fields. In fact, since $\bar{5} \times 75 = 45 + \bar{50} + 280$, in this case only the pure 45 channel contributes. For instance

$$\sqrt{2} \Psi_{10\ i}^{AB} G_{dij} \Psi_{\bar{5}Cj} \overline{H}_D \frac{\langle Y_{AB}^{CD} \rangle}{\Lambda} \quad (3.42)$$

gives $G_{dij} = y_{dij} = -1/3y_{eji}$. In the next chapter the strategy of inserting the 75 to obtain a phenomenologically acceptable spectrum of fermions will be employed in the context of a realistic version of the MDM.

3.6 Coupling Unification

The most outstanding support to the idea of Grand Unification comes from the quantitative success of coupling unification in SUSY GUTs [60]. Assuming just the field content of the MSSM (with two light Higgses), in the leading order approximation (LO), that is at one loop, the RGE for the running coupling measured at m_Z read:

$$\frac{1}{\alpha_i(Q)} = \frac{1}{\alpha_i(m_Z)} + \frac{b_i}{2\pi} \ln \left(\frac{Q}{m_Z} \right) \quad (3.43)$$

with

$$b_1 = \frac{33}{5} \quad , \quad b_2 = 1 \quad , \quad b_3 = -3 \quad . \quad (3.44)$$

Nowadays it is possible, thanks to the quite precise measurements of the three running gauge coupling at m_Z , to single out a not too broad range for the scale of SUSY unification M_{GUT} . Fig. 3.1 shows the flow of the gauge coupling in the one loop approximation starting at m_Z with the PDG data of last summer [92] , that is, in the $\overline{MS}$ scheme: $\alpha_{em}(m_Z) = 127.934 \pm 0.027$, $\sin^2 \theta(m_Z) = 0.23117 \pm 0.00016$, $m_Z = 91.1872 \pm 0.0021$ GeV and the world average $\alpha_3(m_Z) = 0.1184 \pm 0.0012$. As can be clearly seen, unification is very successful and the associated scale M_{GUT} turns out to be in the small range $(1.9 \div 2.1) \cdot 10^{16}$ GeV, with the previously quoted errors.

Now it is also well established that GUT without SUSY is neatly disfavoured, as appears clearly from the comparison of figs. 3.1 and 3.2. For the SM equations analogous to (3.43) hold, but with $b_1 = 41/10$, $b_2 = -19/6$, $b_3 = -7$.

Going beyond LO turns out to be necessary because the corrections from two loop running on the prediction for $\alpha_3(m_Z)$ are of the order of 10%. It is reasonable however to disregard three loop running, which would give correction of only about 1% on $\alpha_3(m_Z)$. In the following we will describe the calculations relative to the two loop prediction of $\alpha_3(m_Z)$ starting from $\alpha_1(m_Z)$, $\alpha_2(m_Z)$ and assuming coupling unification. We will give numerical results for the case of minimal $SU(5)$. They will be useful for comparison with the analogous predictions obtained in the MDM case, whose analysis will be present in detail in the next chapter.

Different GUT models give different predictions for $\alpha_3(m_Z)$ because the contribution to its running coming from threshold effects is sizeable. If one is convinced that the success of coupling unification in the LO approximation is not just a coincidence, then, in model building, one is constrained to introduce as few as possible GUT fields, which in addition have to lie in not too large representations. It is clear that working with a lot of field is equivalent to have a lot of degrees of freedom at disposal, which can be arranged ad hoc to reproduce the data. The elegance of the model is completely lost, but at least one possess a sort of existence proof of the realizability of low energy data starting from a GUT scenario. Working with few fields, instead, it is much more hard to give a completely satisfactory picture of particle phenomenology. However, this latter point of view is the only one from which a true deeper understanding of fundamental constituents could come.

SUSY Standard Model




Figure 3.1: Evolution of the running gauge couplings within the MSSM. The curves are obtained in the 1 loop approximation by evolving the gauge couplings from their measured values at m_Z [92]: $\alpha_1(m_Z) = 0.016945 \pm 0.000007$, $\alpha_2(m_Z) = 0.033813 \pm 0.000003$ (from $\alpha_{em}(m_Z) = 127.934 \pm 0.027$, $\sin^2 \theta(m_Z) = 0.23117 \pm 0.00016$, $m_Z = 91.1872 \pm 0.0021$ GeV) and $\alpha_3(m_Z) = 0.1184 \pm 0.0012$.

Standard Model




Figure 3.2: The same as fig. 3.1 but within the SM.

3.6.1 Coupling Unification Beyond the Leading Order Approximation

It is well known that, going beyond the LO approximation, in the minimal version of SUSY $SU(5)$ the central value of $\alpha_3(m_Z)$ required by the constraint of coupling unification is somewhat large: $\alpha_3(m_Z) \approx 0.13$ [64, 96, 97, 100, 101].

From the one-loop renormalization group evolution of the couplings in the MSSM with only two light Higgs doublets, eqs. (3.43), by evolving two couplings from their values measured at m_Z and imposing the constraint of unification, one can run back to low energies with the third coupling thus giving a prediction for its value at m_Z . In this procedure it is preferable to take as input the two best measured couplings, that is $\alpha_1(m_Z), \alpha_2(m_Z)$. In fact, defining in the $\overline{MS}$ scheme $\alpha \equiv \alpha_{QED}(m_Z)$, $\sin^2 \theta \equiv \sin^2 \theta(m_Z)$,

$$\alpha_1(m_Z) = \frac{\alpha}{1 - \sin^2 \theta} \quad , \quad \alpha_2(m_Z) = \frac{\alpha}{\sin^2 \theta} \quad (3.45)$$

and the predictions are:

$$\begin{aligned} \ln(M_{GUT}^{(0)}/m_Z) &= \frac{\pi(3 - 8 \sin^2 \theta)}{14\alpha} \\ a_5^{(0)} &= \frac{28\alpha}{36 \sin^2 \theta - 3} \\ a_3^{(0)} &= \frac{28\alpha}{60 \sin^2 \theta - 12} \end{aligned} \quad (3.46)$$

where $a_3 \equiv \alpha_3(m_Z)$, $a_5 \equiv \alpha_5(M_{GUT})$ and the superscript (0) refers to a quantity evaluated in the LO approximation. It is worth to stress that the LO results are the same in minimal SUSY $SU(5)$ and in all its extensions where the MSSM β functions apply until M_{GUT} .

By using $\alpha^{-1} = 127.934$ and $\sin^2 \theta = 0.231$ one obtains $a_3^{(0)} = 0.118$ and $M_{GUT}^{(0)} = 2.1 \cdot 10^{16}$ GeV.

To go beyond the LO one must include two loop effects in the running of gauge couplings, threshold effects at the scale m_{SUSY} , close to the electroweak scale, and threshold effects at the large scale M_{GUT} . The unification scale M_{GUT} is not univocally defined beyond LO and here we choose to identify it with the mass of the $SU(5)$ superheavy gauge bosons m_V . Following ref. [64] it is possible to include these effects by writing:

$$\frac{1}{\alpha_i(m_Z)} = \frac{1}{a_5} + \frac{b_i}{2\pi} \ln \left(\frac{M_{GUT}}{m_Z} \right) + \frac{\delta_i}{\pi} \quad (3.47)$$

where $a_5 = \alpha_5(M_{GUT})$ and the next to leading corrections are contained in δ_i , $i = 1, 2, 3$ and can be separated according to their origin. The logarithmic contribution

coming from two loop running, $\delta_i(2)$, and the contributions of SUSY and GUT thresholds, respectively $\delta_i(l)$ and $\delta_i(h)$:

$$\delta_i = \delta_i(2) + \delta_i(l) + \delta_i(h) \quad (3.48)$$

The LO eqs. (3.46) are thus improved by:

$$\begin{aligned} \ln \left(\frac{M_{GUT}}{m_Z} \right) &= \ln \left(\frac{M_{GUT}^{(0)}}{m_Z} \right) + \delta_X \\ \frac{1}{a_5} &= \frac{1}{a_5^{(0)}} + \delta_5 \\ a_3 &= \frac{a_3^{(0)}}{(1 + a_3^{(0)} \delta_3)} \end{aligned} \quad (3.49)$$

where the various $\delta_{X,5,3}$ are linear combinations of δ_i :

$$\begin{aligned} \delta_X &= \frac{10}{28}(\delta_2 - \delta_1) \\ \delta_5 &= -\frac{1}{28}(33\delta_2 - 5\delta_1) \\ \delta_3 &= \frac{1}{7\pi}(5\delta_1 - 12\delta_2 + 7\delta_3) \end{aligned} \quad (3.50)$$

In order to obtain the prediction for a_3 it is necessary to specify the various contributions which enters $\delta_3 = \delta_3(2) + \delta_3(l) + \delta_3(h)$.

Two-loop running

The one from two-loop running is finite and given by:

$$\delta_i(2) = \sum_{j=1}^3 \frac{b_{ij}}{2b_j} \ln \left(1 + \frac{b_j}{2\pi} a_5^{(0)} \ln \left(\frac{M_{GUT}^{(0)}}{m_Z} \right) \right) \quad (3.51)$$

where

$$b_{ij} = \begin{bmatrix} 199/50 & 27/10 & 44/5 \\ 9/10 & 25/2 & 12 \\ 11/10 & 9/2 & 7 \end{bmatrix} \quad (3.52)$$

In particular

$$\delta_3(2) = -0.823 \quad . \quad (3.53)$$

SUSY Thresholds

The one loop thresholds from SUSY particles read:

$$\delta_i(l) = - \sum_j b_i^{(l)}(j) \ln \left(\frac{m_j}{m_Z} \right) \quad (3.54)$$

where j runs over all light particles of mass m_j and $b_i^{(l)}(j)$ can be read from Table 3.6.

sparticle	$b_1^{(l)}(j)$	$b_2^{(l)}(j)$	$b_3^{(l)}(j)$
gluinos	0	0	1
winos	0	2/3	0
higgsinos	1/5	1/3	0
extra Higgses	1/20	1/12	0
$\tilde{q}_L$	1/60	1/4	1/6
$\tilde{u}_R$	2/15	0	1/12
$\tilde{d}_R$	1/30	0	1/12
$\tilde{l}_L$	1/20	1/12	0
$\tilde{e}_R$	1/10	0	0

Table 3.6: SUSY thresholds.

To give a concrete example, one has to specify a certain SUSY spectrum. We will consider as representative the one given in Table B.1, where the additional freedom related to the parameters m_0 , $m_{1/2}$, μ , m_H can be fixed by choosing $0.8m_0 = 0.8m_{1/2} = 2\mu = m_H$ and taking as a definition $m_{SUSY} \equiv m_H$, so that all particle masses can be expressed in term of m_{SUSY} .

The result is then

$$\delta_3(l) = -0.503 + \frac{19}{28\pi} \ln \left(\frac{m_{SUSY}}{m_Z} \right) . \quad (3.55)$$

GUT Thresholds

Threshold contributions from heavy particles can be expressed as:

$$\delta_i(h) = c_i - \sum_j b_i^{(j)}(h) \ln \left(\frac{m_j}{m_{GUT}} \right) \quad (3.56)$$

with $c_1 = 0$, $c_2 = 1/6$, $c_3 = 1/4$ and the sum is over all heavy particles.

To specialize to the case of minimal $SU(5)$ one has to consider its heavy spectrum. After the Higgs mechanism, 12 complex fields of the 24 representation (the 24 would-be Goldstone bosons) are eaten by the 24_V of gauge vector bosons, which thus become massive. The other 12 complex fields in the 24 acquire instead a heavy mass. In addition to them, there are the colour triplets of the 5 and $\bar{5}$ representation. Thresholds can be evaluated according to:

state $(SU(3), SU(2), U(1))$	$b_1^{(h)}(j)$	$b_2^{(h)}(j)$	$b_3^{(h)}(j)$	mass
$(8, 1, 0)$	0	0	3/2	m_Σ
$(1, 3, 0)$	0	1	0	m_Σ
$(1, 1, 0)$	0	0	0	$0.2m_\Sigma$
$(3, 1, -1/3), (\bar{3}, 1, 1/3)$	1/5	0	1/2	m_T

Table 3.7: GUT thresholds. $m_\Sigma \equiv 5M_{24}$, with M_{24} defined as in section 3.1.1.

For this case one obtains

$$\delta_3^{min}(h) = -0.0114 - \frac{9}{14\pi} \ln \left(\frac{m_T}{M_{GUT}} \right) + \frac{3}{14\pi} \ln \left(\frac{m_\Sigma}{M_{GUT}} \right) . \quad (3.57)$$

Results

The first of eqs. (3.49) turns out to be very useful. The contributions to its corrective term δ_X coming from two-loop running and SUSY thresholds induced by the spectrum of Table B.1 are:

$$\delta_X(2) = 0.469 \quad (3.58)$$

$$\delta_X(l) = -0.036 - \frac{25}{84} \ln \left(\frac{m_{SUSY}}{m_Z} \right) . \quad (3.59)$$

Thresholds from GUT particles within minimal $SU(5)$ give:

$$\delta_X(h) = 0.060 - \frac{5}{14} \ln \left(\frac{m_\Sigma}{M_{GUT}} \right) + \frac{1}{14} \ln \left(\frac{m_T}{M_{GUT}} \right) . \quad (3.60)$$

Here in minimal $SU(5)$ it turns out clearly, but it is true in general that, keeping fixed m_{SUSY} , the first of eqs. (3.49) can be considered as a relation between the GUT masses. In this case it relates $m_V = M_{GUT}$ to m_Σ, m_T .

Substituting in $\delta_3(h)$ the expression thus obtained for M_{GUT} , one can always write δ_3 according to the following expression:

$$\delta_3 = k + \frac{1}{2\pi} \ln \left(\frac{m_{SUSY}}{m_Z} \right) - \frac{3}{5\pi} \ln \left(\frac{m_T}{M_{GUT}^{(0)}} \right) \quad (3.61)$$

This expression for δ_3 is general because we have subdivided the various contributions in a numerical coefficient plus logarithmic contributions. The logarithmic term with m_{SUSY} comes from particles with masses near m_{SUSY} . Similarly, the logarithmic term with m_T arises from particles with masses of order $m_T \approx M_{GUT}$. The definition of m_T is the mass of Higgs colour triplets in the minimal model or the effective mass that, in realistic models, plays the same role for proton decay [64, 96]. Notice that there isn't any dependence on m_Σ . The term indicated with $k = k(2) + k(SUSY) + k(M_{GUT})$ contains the contribution of two loop diagrams to the running couplings, $k(2)$, the threshold contribution from states near m_{SUSY} , $k(SUSY)$, and the threshold contribution from states near M_{GUT} , $k(M_{GUT})$. This way of subdividing the contribution is useful because the threshold contributions would vanish if all states had the mass m_{SUSY} or M_{GUT} so that only mass splittings contribute to $k(SUSY)$ and $k(M_{GUT})$.

The values of $k(2)$ and of $k(SUSY)$ are the same in minimal $SU(5)$ and in its realistic extensions, because only the heavy spectrum changes. Typical values are $k(2) = -0.733$ and $k(SUSY) = -0.510$. The value for $k(SUSY) = -0.510$ corresponds to the representative spectrum displayed in Table B.1. Note that the




Figure 3.3: Contours of $a_3 \equiv \alpha_s(m_Z)$ in the plane (m_{SUSY}, m_T) , for minimal $SU(5)$. The SUSY spectrum is parametrized as in Table B.1.

value of $k(M_{GUT})$ is practically zero for the 24 of the minimal model. The reason is that in minimal $SU(5)$ all coloured GUT multiplets are degenerate, as appears clearly giving a look to Table 3.7. The result is thus:

$$k = -0.733 - 0.510 = -1.243 \quad \text{minimal model} \quad (3.62)$$

In fig. 3.3 the prediction for a_3 in the plane (m_{SUSY}, m_T) is shown for the case of minimal $SU(5)$. It can be seen that, in this case, we need to take m_{SUSY} as large as possible and m_T as small as possible. But the smaller is m_T , the faster is proton decay. The best compromise is something like $m_{SUSY} \approx 1$ TeV and $m_T \approx M_{GUT}^{(0)}$, which leads to $\alpha_3(m_Z) \approx 0.13$ which is still rather large and proton decay is dangerously fast, in conflict with experimental limits. In fact, with this value for m_T , we already know from the discussion of section 3.4.5 that proton lifetime for the channel $p \rightarrow K^+ \bar{\nu}$ is expected to be around 10^{31} yr and that, in the minimal model, there is no way to push it above the experimental limit of $\sim 2 \cdot 10^{33}$ yr. Thus, we are lead to conclude that minimal SUSY $SU(5)$ is ruled out.

As already pointed out, models based on $SO(10)$ are highly model dependent, because the choice of Higgs fields which are introduced to realize the breaking of $SO(10)$ affects many predictions, among which proton decay and coupling unification. For instance, in the quite economic $SO(10)$ model of ref. [42], the corrections to $\alpha_3(m_Z)$ coming from representations like 10, $\overline{10}$, 16, $\overline{16}$ and 45 are expected to be positive and of order 10% with respect to the experimental value of $\alpha_3(m_Z)$, so that one needs to invoke negative corrections of the same order arising from Planck scale effects or other thresholds.

Chapter 4

A realistic SUSY $SU(5)$ GUT Model

It has emerged from the survey of the previous chapter that, despite the idea of GUT is really very attractive, its actual realization is not at all precisely defined. Consider also that at the Planck scale, M_{Pl} , the unification of gravity with gauge interactions is a general property of superstring theories. But in simple GUT models gauge interactions are unified at a distinctly lower mass scale M_{GUT} within the context of a renormalizable gauge theory. However, this neat separation between gauge unification and merging with gravity is not at all granted. The gap between M_{GUT} and M_{Pl} could be filled up by a number of threshold effects and several layers of additional states. Also, coupling unification could be realized without gauge unification, as suggested in some versions of superstring theory or in flipped $SU(5)$.

The model we aim at should not rely on large fine tunings and must lead to an acceptable phenomenology. This includes coupling unification with an acceptable value of $\alpha_s(m_Z)$, given α and $\sin^2\theta_W$ at m_Z , compatibility with the more and more stringent bounds on proton decay [61, 77], agreement with the observed fermion mass spectrum, also considering neutrino masses and mixings and so on.

Prompted by recent neutrino oscillation data some new studies on realistic GUT models have appeared in the context of $SO(10)$ or larger groups [42, 93]. Here we address the question whether the smallest SUSY $SU(5)$ symmetry group can still be considered as a basis for a realistic GUT model. We indeed present an explicit example of a realistic $SU(5)$ model [94], which uses a $U(1)$ flavour symmetry as a crucial ingredient. In principle the flavour symmetry could be either global or local.

We tentatively assume here that the flavour symmetry is global. This is more in the spirit of GUT's in the sense that all gauge symmetries are unified. The associated Goldstone boson receives a mass from the anomaly. Such a pseudo-Goldstone boson

can be phenomenologically acceptable in view of the existing limits on axion-like particles [95].

In this model the doublet-triplet splitting problem is solved by the MDM [63] stabilized by the flavour symmetry against the occurrence of doublet mass lifting due to non renormalizable operators. Relatively large representations (50, $\overline{50}$, 75) have to be introduced for this purpose. A good effect of this proliferation of states is that the value of $\alpha_s(m_Z)$ obtained from coupling unification in the next to the leading order perturbative approximation receives important negative corrections from threshold effects near the GUT scale. As a result, the central value changes from $\alpha_s(m_Z) \approx 0.130$ in minimal SUSY $SU(5)$ down to $\alpha_s(m_Z) \approx 0.116$, in better agreement with observation [64, 96, 97]. The same $U(1)$ flavour symmetry that stabilizes the missing partner mechanism is used to explain the hierarchical structure of fermion masses. In the neutrino sector, mass matrices very similar to the ones already presented in the first model of section 2.1.3 (and proposed in ref. [32]) are obtained. The large atmospheric neutrino mixing is due to a large left handed mixing in the lepton sector that corresponds to a large right handed mixing in the down quark sector. In the present particular version maximal mixing also for solar neutrinos is preferred. A possibly problematic feature of the model is that, beyond the unification point, when all the states participate in the running, the asymptotic freedom of $SU(5)$ is destroyed by the large number of matter fields. As a consequence, the coupling increases very fast and the theory becomes non perturbative below M_{Pl} . In the past models similar to ours have been considered, but were discarded just because they contain many additional states and tend to become non perturbative between M_{GUT} and M_{Pl} . We instead argue that these features are not necessarily bad. While the predictivity of the theory is reduced because of non renormalizable operators that are only suppressed by powers of M_{GUT}/Λ with $\Lambda < M_{Pl}$, still these corrections could explain the distortions of the mass spectrum with respect to the minimal model, the suppression of proton decay and so on. However, it is certainly true that also in this case, as for any other known realistic model, the resulting construction is considerably more complicated than in the corresponding minimal model.

4.1 The Model

The symmetry of the model is SUSY $SU(5) \otimes U(1)$. We define Q the charge associated with $U(1)$ and assume that the global $U(1)$ flavour symmetry is a good symmetry of the superpotential, that is it is not violated by non-perturbative effects through operators appearing in the superpotential w . The superpotential of the

model can be thought as composed of three parts:

$$w = w_1 + w_2 + w_3 \quad , \quad (4.1)$$

where w_1 accounts for the breaking of $SU(5)$, w_2 realizes the doublet-triplet splitting by means of the MDM and w_3 contains the Yukawa couplings.

4.1.1 $SU(5)$ Breaking

The w_1 term contains only the field Y of the representation 75 of $SU(5)$ with $Q = 0$:

$$w_1 = c_1 Y_{CD}^{AB} Y_{EF}^{CD} Y_{AB}^{EF} + M_Y Y_{CD}^{AB} Y_{AB}^{CD} \quad , \quad (4.2)$$

where $A, B, \dots = 1, 2, \dots, 5$. w_1 has the effect of providing Y with a vev $\langle Y \rangle \equiv M_Y/(2c_1) \sim M_{GUT}$ and of giving a mass to all physical components of Y , i.e. those that are not absorbed by the Higgs mechanism. The 75 uniquely breaks $SU(5)$ down to $SU(3) \otimes SU(2) \otimes U(1)$ through the following vev:

$$\langle Y_{CD}^{AB} \rangle = -\langle Y \rangle \left[\left(\delta_\gamma^\alpha \delta_\delta^\beta + 2\delta_c^a \delta_d^b - \frac{1}{2} \delta_C^A \delta_D^B \right) - (A \leftrightarrow B) \right] \quad , \quad (4.3)$$

where Greek indices assume values from 1 to 3 and Latin indices from 4 to 5.

4.2 Doublet-Triplet Splitting

The w_2 term induces the doublet-triplet splitting:

$$\begin{aligned} w_2 = & c_2 H^A Y_{BC}^{DE} H_{50}^{BCFG} \epsilon_{ADEF G} + c_3 \overline{H}_A Y_{DE}^{BC} H_{\overline{50}BCFG} \epsilon^{ADEF G} \\ & + c_4 H_{50}^{ABCD} H_{\overline{50}ABCD} X \quad . \end{aligned} \quad (4.4)$$

Here H and $\overline{H}$ are the usual pentaplets of Higgs fields, except that now they carry non opposite Q charges. It is not restrictive to take c_2 , c_3 and c_4 real and positive. The renormalizable couplings that appear in w_2 are the most general allowed by the $SU(5)$ and Q assignments of the fields in eq. (4.4) which are given in Table 4.1. The value of q will be specified later.

4.2.1 Limit of Unbroken SUSY

At the minimum of the potential, in the limit of unbroken SUSY, the vevs of the fields H , $\overline{H}$, H_{50} and $H_{\overline{50}}$ all vanish, while the X vev remains undetermined. These

field	Y	H	$\overline{H}$	H_{50}	$H_{\overline{50}}$	X
$SU(5)$	75	5	$\overline{5}$	50	$\overline{50}$	1
Q	0	$-q$	$q - 1$	q	$1 - q$	-1

Table 4.1: Q charge assignments.

can be easily understood by writing symbolically the equations for the minimum of the superpotential $w_i = 0$, the subscript i standing for derivation:

$$\begin{aligned}
w_Y &= 2M_Y Y + 3c_1 Y^2 + c_2 H H_{50} + c_3 \overline{H} H_{\overline{50}} \\
w_{50} &= c_2 H Y + c_4 X H_{\overline{50}} \\
w_{\overline{50}} &= c_3 \overline{H} Y + c_4 X H_{50} \\
w_H &= c_2 Y H_{50} \\
w_{\overline{H}} &= c_3 Y H_{\overline{50}} \\
w_X &= c_4 H_{50} H_{\overline{50}}
\end{aligned} \tag{4.5}$$

For any $\langle X \rangle$, a minimum is achieved by taking $\langle H \rangle = \langle \overline{H} \rangle = \langle H_{50} \rangle = \langle H_{\overline{50}} \rangle = 0$ and $\langle Y \rangle$ as shown in eq. (4.3).

As we shall see, when SUSY is softly broken the light doublets in H and $\overline{H}$ acquire a small vev while the X vev will be fixed near the cut-off Λ , of the order of the scale between M_{GUT} and M_{Pl} where the theory becomes strongly interacting (we shall see that we estimate this scale at around $10 - 20 M_{GUT}$, large enough that the approximation of neglecting terms of order M_{GUT}/Λ is not unreasonable). The MDM occurs as discussed in eqs. (3.9) and (3.10) with the only difference that the mass term for H_{50} and $H_{\overline{50}}$ originates from the vev of X . The $U(1)$ flavour symmetry protects the doublet Higgs to take mass from radiative corrections because no $H\overline{H}$ mass term is allowed. Also no non renormalizable terms of the form $H\overline{H}Y^m X^n$ ($m, n \geq 0$) are possible, because X has a negative Q charge. As anticipated in section 3.2.1, this version of the MDM was introduced in ref. [67] and overcomes the observation in ref. [68] that, in general, non renormalizable interactions spoil the mechanism. The Higgs colour triplets mix with the analogous states in the 50

and the resulting mass matrix is of the see-saw form:

$$\hat{m}_T = \begin{bmatrix} 0 & -4\sqrt{3}c_2\langle Y \rangle \\ -4\sqrt{3}c_3\langle Y \rangle & c_4\langle X \rangle \end{bmatrix} . \quad (4.6)$$

Defining $m_\phi = c_4\langle X \rangle$ the eigenvalues of the matrix $\hat{m}_T\hat{m}_T^\dagger$ are the squares of:

$$m_{T1,2} = \frac{1}{2} \left[\sqrt{m_\phi^2 + 48 (c_2 + c_3)^2 \langle Y \rangle^2} \pm \sqrt{m_\phi^2 + 48 (c_2 - c_3)^2 \langle Y \rangle^2} \right] . \quad (4.7)$$

and $m_{T1}m_{T2} = 48 c_2 c_3 \langle Y \rangle^2$. The effective mass that enters in the dimension 5 operators with $|\Delta B = 1|$ is $1/(\hat{m}_T^{-1})_{11}$:

$$m_T = \frac{m_{T1}m_{T2}}{m_\phi} = \frac{48 c_2 c_3 \langle Y \rangle^2}{c_4 \langle X \rangle} . \quad (4.8)$$

4.2.2 Broken Supersymmetry

The breaking of SUSY fixes a large vev for the field X by removing the corresponding flat direction, gives masses to s-partners, provides a small mass to the Higgs doublet and introduces a μ term. In the $(50, \bar{50}, X)$ sector, the terms that break SUSY softly are:

$$-\mathcal{L}_{soft} = m_x^2|x|^2 + m_{50}^2|c|^2 + m_{\bar{50}}^2|\bar{c}|^2 + m_{3/2}c_4A_4(c\bar{c}x + h.c.) + \dots \quad (4.9)$$

where c , $\bar{c}$ and x denote the scalar components of H_{50} , $H_{\bar{50}}$ and X respectively, m_x , m_{50} , $m_{\bar{50}}$ their soft breaking masses, $m_{3/2}$ the gravitino mass and A_4 a trilinear coupling. Note that c_4 has here the same role played in SUSY radiative electroweak breaking by the top Yukawa coupling. Dots stand for the remaining soft breaking terms, including mass terms for the scalar components of the Higgs doublet fields ¹.

Consider first the limit $m_x = m_{50} = m_{\bar{50}} = m_{3/2} \equiv m$ and $c_4A_4 = 1$. In the SUSY limit and neglecting the mixing between the $(50, \bar{50})$ and the $(5, \bar{5})$ sectors, fermions and scalars in the 50 and $\bar{50}$ have a common squared mass $c_4^2x^2$. When SUSY is broken by the soft terms for each fermion of mass c_4x there are two bosons of squared masses $c_4^2x^2 + m^2 \pm mx$. The x terms in the scalar potential at one loop accuracy are given by:

$$V = m^2x^2 + \frac{50}{64\pi^2} \left\{ 2(c_4^2x^2 + m^2 + mx)^2 \left[\log \left(\frac{c_4^2x^2 + m^2 + mx}{\Lambda^2} \right) - 3/2 \right] \right.$$

¹To find the minima of the scalar potential it is not restrictive to set to zero the imaginary part of x , which will be understood in the remaining part of this section.

$$\begin{aligned}
& + 2(c_4^2 x^2 + m^2 - mx)^2 \left[\log \left(\frac{c_4^2 x^2 + m^2 - mx}{\Lambda^2} \right) - 3/2 \right] \\
& - 4c_4^4 x^4 \left[\log \left(\frac{c_4^2 x^2}{\Lambda^2} \right) - 3/2 \right] \} \quad (4.10)
\end{aligned}$$

A numerical study of this potential in the limit $m \ll \Lambda$ and for c_4 of order one, shows that the minimum is at x of order Λ (somewhat smaller than Λ but close to it). The expression for V in eq. (4.10) provides a good approximation of the scalar potential only in a small region around Λ . Outside this region the perturbative approximation will break down. We take our numerical analysis as an indication that the minimum occurs near Λ .

To analyze in more detail the situation we consider the differential equation describing the evolution of the x mass and we identify $\langle X \rangle$ with the scale at which m_x^2 vanishes. With the soft terms of (4.9), we find that the following set of differential equations is appropriate to describe the running in range between M_{GUT} and the cut-off:

$$\begin{aligned}
\frac{dm_x^2}{d \ln Q} &= \frac{25}{4\pi^2} c_4^2 (m_x^2 + m_{50}^2 + m_{\overline{50}}^2 + m_{3/2}^2 A_4^2) \\
\frac{dm_{50}^2}{d \ln Q} &= \frac{dm_{\overline{50}}^2}{d \ln Q} = -\frac{84}{5\pi} \alpha \tilde{m}^2 + \frac{1}{8\pi^2} c_4^2 (m_x^2 + m_{50}^2 + m_{\overline{50}}^2 + m_{3/2}^2 A_4^2) \\
\frac{dc_4}{d \ln Q} &= -\frac{42}{5\pi} \alpha c_4 + \frac{13}{4\pi^2} c_4^3 \\
\frac{d(m_{3/2} A_4)}{d \ln Q} &= -\frac{84}{5\pi} \alpha \tilde{m} + \frac{13}{2\pi^2} m_{3/2} c_4^2 A_4 \\
\frac{d\tilde{m}}{d \ln Q} &= \frac{52}{2\pi} \alpha \tilde{m} \\
\frac{d\alpha}{d \ln Q} &= \frac{52}{2\pi} \alpha^2
\end{aligned} \quad (4.11)$$

where α is the running gauge coupling and $\tilde{m}$ are gaugino masses, which are supposed to be degenerate. The value of $\langle X \rangle$ depends on the boundary conditions assigned to the relevant parameters at the cut-off scale, in particular c_4 and the trilinear $c\bar{c}x$ coupling in $\mathcal{L}_{soft}$. To give a numerical example, if $\Lambda = 1.1 \cdot 10^{17}$ GeV and $\alpha(\Lambda) = 0.88$, then $\langle X \rangle \sim \Lambda/4$ is obtained by choosing the following reasonable values of the parameters at the cut-off scale: $m_x = m_{50} = m_{\overline{50}} = m_{3/2} = 800$ GeV, $\tilde{m} = 2$ TeV, $c_4 = 0.2$ and $A_4 = 0.7$. This particular situation is shown in fig. 4.1, where the running of $m_x^2, m_{50}^2, m_{\overline{50}}^2$ and $m_m^2 \equiv (m_{3/2} c_4 A_4)^2$ is displayed. In the next section we assume that the true minimum occurs at $x = \langle X \rangle = 0.25\Lambda$, that is, by defining $\lambda \equiv \langle X \rangle / \Lambda$, at $\lambda = 0.25$.




Figure 4.1: Running of $m_x^2, m_{50}^2, m_{\overline{50}}^2$ and $m_m^2 \equiv (m_{3/2} c_4 A_4)^2$. The solid (dashed) lines correspond to: $\Lambda = 1.1 \cdot 10^{17}$ GeV, $\alpha(\Lambda) = 0.88$, $m_x = m_{50} = m_{\overline{50}} = m_{3/2} = 800$ GeV, $\tilde{m} = 2$ TeV, $c_4 = 0.2$ ($c_4 = 0.4$) and $A_4 = 0.7$. The solid red line gives a $\lambda \equiv \langle X \rangle / \Lambda \sim 0.25$.

We have studied the dependence of $\langle X \rangle$ upon both the boundary values of c_4 and A_4 . We find that the dependence on $c_4(\Lambda)$ is logarithmic and that there is a mild linear dependence on the trilinear $c\bar{c}x$ term evaluated at Λ . The above dependence results mainly from the interplay of the differential equations describing the evolution of the x mass and the c_4 coupling, the quantities with the faster running in the region close to the cut-off Λ . We obtain: $\delta\langle X \rangle/\Lambda \approx 0.4 \delta c_4/c_4$. This means that, with the same boundaries of the previous numerical example but with c_4 doubled, the value of $\langle X \rangle$ becomes greater by a factor 2.2, as can be seen from fig. 4.1. The vev of X seems thus to be not dangerously unstable with respect to the boundaries. This situation differs from the analogous mechanism of radiative electroweak symmetry breaking, where a doubling of y_t can cause a variation of several orders of magnitude in the vev of the Higgs doublet. We have checked that, in our case, the mild dependence on c_4 is due to the running of c_4 itself which, in the appropriate energy window, is not negligible.

The complex field x describes two physical scalar particles. The one associated to the real part of x has a mass of order $m_{3/2}$ and couplings to the ordinary fermions suppressed by $1/\Lambda$. The particle associated to the imaginary part of x is massless only in the tree approximation. Due to the anomaly of the related $U(1)_Q$ current the particle acquires a mass of order $f_\pi m_\pi/\langle X \rangle$. Cosmological bounds on $\langle X \rangle$ have been recently reconsidered in ref. [95] where it has been observed that $\langle X \rangle$ of the order of the grand unification scale is not in conflict with observational data.

An alternative possibility is to assume that the $U(1)$ symmetry is local [67]. In this case the supersymmetric action contains a Fayet-Iliopoulos term and the associated D-term in the scalar potential provides a large vev for x , of the order of the cut-off scale Λ .

A μ term for the fields H and $\overline{H}$ of the appropriate order of magnitude can be generated according to the Giudice-Masiero mechanism [99]. Assume that the breaking of SUSY is induced by the θ^2 component of a chiral (effective) superfield S , singlet under $SU(5)$ with $Q = 0$. In the Kahler potential a term of the form

$$K = \frac{S^\dagger X^\dagger H \overline{H}}{\Lambda^2} + \text{h.c.} \quad (4.12)$$

is allowed. The vevs of S and X , $\langle S \rangle \sim \theta^2 m_{3/2} M_{Pl}$, $\langle X \rangle = \lambda \Lambda$ lead to an equivalent term in the superpotential of the desired form $\mu H \overline{H}$ with $\mu \sim \lambda m_{3/2} M_{Pl}/\Lambda$ which can be considered of the right order.

4.3 Coupling Unification

As was pointed out previously, it is well known that in the minimal version of SUSY $SU(5)$ the central value of $\alpha_3(m_Z)$ required by the constraint of coupling unification is somewhat large: $\alpha_3(m_Z) \approx 0.13$ [64, 96, 97, 100, 101]. In the model discussed here, where the doublet-triplet splitting problem is solved by introducing the $SU(5)$ representations 50, $\bar{50}$ and 75, the central value of $\alpha_3(m_Z)$ is modified by threshold corrections near M_{GUT} which bring the central value down by a substantial amount so that the final result can be in much better agreement with the observed value. As discussed in refs. [64, 96], this remarkable result arises because the 24 of the minimal model is replaced by the 75. The mass splittings inside these representations are dictated by the group embedding of the 75 singlet under $SU(3) \otimes SU(2) \otimes U(1)$ that breaks $SU(5)$. The difference in the threshold contributions from the 24 and the 75 has the right sign and amount to bring $\alpha_3(m_Z)$ down even below the observed value. The right value can then be obtained by moving m_{SUSY} and m_T in a reasonable range. The difference in the favored value of m_T with respect to the minimal model in order to reproduce the observed value of $\alpha_3(m_Z)$ goes in the right direction to also considerably alleviate the potential problems from the bounds on proton decay. Note that there are no additional threshold corrections from the 50 and $\bar{50}$ representations because the mass of these states arise from the coupling to the field X which is an $SU(5)$ singlet. Thus there are no mass splittings inside the 50 and the $\bar{50}$ and no threshold contributions. While there is no effect on the value of $\alpha_3(m_Z)$ the presence of the states in the 50 and $\bar{50}$ representations affects the value of the unified coupling at M_{GUT} and also spoils the asymptotic freedom of $SU(5)$ beyond M_{GUT} . We find it suggestive that the solution of the doublet-triplet splitting problem in terms of the 50, $\bar{50}$ and 75 representations automatically leads to improve the prediction of $\alpha_3(m_Z)$ and at the same time relaxes the constraints from proton decay. We now discuss this issue in more detail.

4.3.1 Estimate of $\alpha_3(m_Z)|_{\overline{MS}}$

The calculation proceeds exactly as in the previous chapter. With the same SUSY spectrum, all the differences on the values of a_5, M_{GUT}, a_3 , with respect to minimal SUSY $SU(5)$, are caused by the term $\delta_i(h)$ of (3.48).

Following the steps already encountered, $\delta_3(2), \delta_3(l)$ are those of eqs. (3.53) and (3.55), but $\delta_3(h)$ is now obtained substituting in eq. (3.56) the $b_i^{(h)}$ reported in Table 4.2.

state ($SU(3), SU(2), U(1)$)	$b_1^{(h)}(j)$	$b_2^{(h)}(j)$	$b_3^{(h)}(j)$	mass
$(8, 3, 0)$	0	8	$9/2$	m_Y
$(3, 1, 5/3)(\bar{3}, 1, -5/3)$	5	0	$1/2$	$0.8m_Y$
$(6, 2, 5/6)(\bar{6}, 2, -5/6)$	5	3	5	$0.4m_Y$
$(1, 1, 0)$	0	0	0	$0.2m_Y$
$(8, 1, 0)$	0	0	$3/2$	$0.2m_Y$
$(3, 1, -1/3), (\bar{3}, 1, 1/3)$	$1/5$	0	$1/2$	m_{T1}
$(3, 1, -1/3), (\bar{3}, 1, 1/3)$	$1/5$	0	$1/2$	m_{T2}
$50, \bar{50}$	$173/10$	$35/2$	17	m_ϕ

Table 4.2: GUT thresholds for the MDM. $m_Y \equiv 5M_Y$.

For this case one obtains

$$\delta_3(h) = 2.047 - \frac{9}{14\pi} \ln \left(\frac{m_T}{M_{GUT}} \right) + \frac{3}{14\pi} \ln \left(\frac{m_Y}{M_{GUT}} \right) \quad (4.13)$$

where m_T is the effective triplet mass. Note that this equation is very similar to eq. (3.57), apart from the non-logarithmic term. We can still use the first of eqs. (3.49), with the difference that thresholds from GUT particles within the MDM now give:

$$\delta_X(h) = -0.993 - \frac{5}{14} \ln \left(\frac{m_Y}{M_{GUT}} \right) + \frac{1}{14} \ln \left(\frac{m_T}{M_{GUT}} \right) . \quad (4.14)$$

Thus, keeping fixed m_{SUSY} , the first of eqs. (3.49) can be considered as a relation

between the GUT masses $m_V = M_{GUT}, m_Y, m_T$.

$$\begin{aligned}
\ln\left(\frac{M_{GUT}}{m_Z}\right) &= \ln\left(\frac{M_{GUT}^{(0)}}{m_Z}\right) \\
&+ [0.469]_{(2)} + \left[-0.036 - \frac{25}{84} \ln\left(\frac{m_{SUSY}}{m_Z}\right)\right]_{(l)} \\
&+ \left[-0.993 - \frac{5}{14} \ln\left(\frac{m_Y}{M_{GUT}}\right) + \frac{1}{14} \ln\left(\frac{m_T}{M_{GUT}}\right)\right]_{(h)} \quad (4.15)
\end{aligned}$$

Substituting it in $\delta_3(h)$ for M_{GUT} , as pointed out in 3.6.1, we can write the prediction for δ_3 in the following form,

$$\delta_3 = k + \frac{1}{2\pi} \ln\left(\frac{m_{SUSY}}{m_Z}\right) - \frac{3}{5\pi} \ln\left(\frac{m_T}{M_{GUT}^{(0)}}\right) \quad (4.16)$$

where again $k = k(2) + k(SUSY) + k(M_{GUT})$ and it contains the contribution of two-loop diagrams to the running couplings, $k(2)$, the threshold contribution of states near m_{SUSY} , $k(SUSY)$, and the threshold contribution from states near M_{GUT} , $k(M_{GUT})$. Remember that the threshold contributions would vanish if all states had the mass m_{SUSY} or M_{GUT} so that only mass splittings contribute to $k(SUSY)$ and $k(M_{GUT})$. The values of $k(2)$ and of $k(SUSY)$ are essentially the same in the minimal and the realistic model. As in the previous chapter we take: $k(2) = -0.733$ and $k(SUSY) = -0.510$, corresponding to the representative spectrum displayed in Table B.1. The value of $k(M_{GUT})$ was zero for the 24 of the minimal model, but now we have $k(M_{GUT}) = 1.857$ for the 75 of the realistic model. The 75 contains in fact non degenerate coloured multiplets. Thus we obtain:

$$\begin{aligned}
k &= -0.733 - 0.510 &= -1.243 &\quad \text{minimal model} \\
k &= -0.733 - 0.510 + 1.857 &= 0.614 &\quad \text{realistic model} \quad (4.17)
\end{aligned}$$

This difference is very important and makes the comparison with experiment of the predicted value of $\alpha_3(m_Z)$ much more favorable in the case of the realistic model. In fact for k large and negative as in the minimal model we need to take m_{SUSY} as large as possible and m_T as small as possible. But the smaller is m_T , the faster is proton decay. The best compromise is something like $m_{SUSY} \approx 1$ TeV and $m_T \approx M_{GUT}^{(0)}$, which leads to $\alpha_s(m_Z) \approx 0.13$ which is still rather large and proton decay is too fast. This is to be confronted with the case of the realistic model where k is instead positive and large enough to drag $\alpha_s(m_Z)$ below the observed value. We now prefer m_T to be larger than $M_{GUT}^{(0)}$ by typically a factor of 20-30, which




Figure 4.2: Contours of $a_3 \equiv \alpha_s(m_Z)$ in the plane (m_{SUSY}, m_T) , for the missing doublet model. The SUSY spectrum is parametrized as in Table B.1.

means a factor of 400-1000 of suppression for the proton decay rate with respect to the minimal model. For example, for $m_{SUSY} \approx 0.25$ TeV and $m_T \approx 6 \cdot 10^{17}$ GeV we obtain $\alpha_3(m_Z) \approx 0.116$ which is acceptable. The predictions for a_3 versus (m_{SUSY}, m_T) in the minimal and the missing doublet model are shown in figs. 3.3 and 4.2.

Due to the large matter content the model is not asymptotically free and the gauge coupling constant α_5 blows up at the scale

$$M_{GUT} \exp\left(\frac{\pi}{26a_5}\right) . \quad (4.18)$$

Taking into account that threshold effects modify a_5 with respect the LO value $a_5^{(0)} \approx 1/24$, we find that the pole occurs near 10^{17} GeV, the precise value depending on the details of the heavy spectrum.

4.3.2 Theoretical Uncertainties

Clearly, there is an uncertainty on $k(SUSY)$, related to the possibility of varying both the parameterization used and the relations assumed between the parameters $m_0, m_{1/2}, \mu, m_H$. An overall uncertainty of ± 0.005 on a_3 has been estimated in ref. [100] by scanning many models consistent with electroweak symmetry breaking, a neutral lightest supersymmetric particle and sparticle masses above the experimental bound and below ~ 2 TeV.

We stress that in the present analysis we are assuming universal soft breaking parameters at the cut-off scale. If we relax this assumption, then a larger range for a_3 could be obtained. Indeed a_3 is particularly sensitive to wino and gluino masses, that, at the electroweak scale, are approximately in the ratio 1:3 when a universal boundary condition on gaugino masses is imposed. It has been observed that, by inverting this ratio, a negative contribution of about -0.01 to a_3 is obtained [102].

Another source of uncertainty on a_3 is related to the unknown physics above the cut-off scale of the grand unified theory. There can be threshold effects due to new heavy particles or even non perturbative effects that arise in the underlying fundamental theory. These effects can be estimated from the non-renormalizable operators, suppressed by $\langle Y \rangle / \Lambda$, that split the gauge couplings at the scale $M_{GUT}^{(0)}$ [103]. For generic, order one coefficients and barring cancellations among different terms, the presence of these operators may affect a_3 by additional contributions of about $\pm 0.005 M_{Pl} / \Lambda$ [100]. As we will see the model under discussion requires $\Lambda \approx 0.1 M_{Pl}$. Therefore, to maintain the good agreement between the experimental and predicted values of a_3 , we need a suppression of this contribution by about a factor of 10, which we do not consider too unnatural.

In view of the large theoretical uncertainties on a_3 we cannot firmly conclude that the gauge coupling unification fails in the minimal model while it is completely successful in the realistic one. However we find very encouraging that, by solving the doublet-triplet splitting problem within the missing partner model, acceptable values of a_3 can be easily obtained and that they are directly related to a proton lifetime potentially larger than in the minimal model.

4.4 Yukawa Couplings

The w_3 term contains the Yukawa interactions of the quark and lepton fields Ψ_{10} , $\Psi_{\bar{5}}$ and Ψ_1 , transforming as the representations 10, $\bar{5}$ and 1 of $SU(5)$ respectively. We assume an exact R -parity discrete symmetry under which Ψ_{10} , $\Psi_{\bar{5}}$ and Ψ_1 are odd whereas H , $\bar{H}$, H_{50} and $H_{\bar{50}}$ are even. The w_3 term is symbolically given by

$$\begin{aligned} w_3 = & \Psi_{10} G_u(X, Y) \Psi_{10} H + \Psi_{10} G_d(X, Y) \Psi_{\bar{5}} \bar{H} + \Psi_{\bar{5}} G_\nu(X, Y) \Psi_1 H \\ & + M \Psi_1 G_M(X, Y) \Psi_1 + \Psi_{10} G_{\bar{50}}(X, Y) \Psi_{10} H_{\bar{50}} \quad . \end{aligned} \quad (4.19)$$

The Yukawa matrices G_u , G_d , G_ν , G_M and $G_{\bar{50}}$ depend on X and Y and the associated mass matrices on their vevs. The last term does not contribute to the mass matrices because of the vanishing vev of $H_{\bar{50}}$, but is important for proton decay. The pattern of fermion masses is determined by the $U(1)$ flavour symmetry that fixes the powers of $\lambda \equiv \langle X \rangle / \Lambda$ for each entry of the mass matrices. In fact X is the only field with non vanishing Q that takes a vev. The powers of λ in the mass terms are fixed by the Q charges of the matter field Ψ and of the Higgs fields H and $\bar{H}$. We can then specify the charge q that appears in Table 4.1 and the Q charges of the matter fields Ψ in order to obtain realistic textures for the fermion masses. We choose $q = 2$, so that we have:

$$Q(H) = -2 \quad \text{and} \quad Q(\bar{H}) = 1 \quad , \quad (4.20)$$

and, for matter fields

$$Q(\Psi_{10}) = (4, 3, 1) \quad , \quad Q(\Psi_{\bar{5}}) = (4, 2, 2) \quad , \quad Q(\Psi_1) = (1, -1, 0) \quad . \quad (4.21)$$

The Yukawa mass matrices are of the form:

$$G_r(\langle X \rangle, \langle Y \rangle)_{ij} = \lambda^{n_{ij}} G_r(\langle Y \rangle)_{ij}, \quad r = u, d, \nu, M \quad . \quad (4.22)$$

We expand $G_r(\langle Y \rangle)_{ij}$ in powers of $\langle Y \rangle$ and consider the lowest order term at first. Taking $G_r(0)_{ij}$ of order 1 and n_{ij} as dictated by the above charge assignments we

obtain ²:

$$\begin{aligned}
m_u &= \frac{1}{\sqrt{2}} \begin{bmatrix} \lambda^6 & \lambda^5 & \lambda^3 \\ \lambda^5 & \lambda^4 & \lambda^2 \\ \lambda^3 & \lambda^2 & 1 \end{bmatrix} v_u , & m_d &= m_e^T = \frac{1}{\sqrt{2}} \begin{bmatrix} \lambda^5 & \lambda^3 & \lambda^3 \\ \lambda^4 & \lambda^2 & \lambda^2 \\ \lambda^2 & 1 & 1 \end{bmatrix} v_d \lambda^4 , \\
m_\nu &= \frac{1}{\sqrt{2}} \begin{bmatrix} \lambda^3 & \lambda & \lambda^2 \\ \lambda & 0 & 1 \\ \lambda & 0 & 1 \end{bmatrix} v_u , & m_{maj} &= \begin{bmatrix} \lambda^2 & 1 & \lambda \\ 1 & 0 & 0 \\ \lambda & 0 & 1 \end{bmatrix} M .
\end{aligned} \tag{4.23}$$

For a correct first approximation of the observed spectrum we need $\lambda \approx \lambda_C \approx 0.22$, λ_C being the Cabibbo angle. These mass matrices are very similar to those already introduced in section 2.1.3, with two important differences.

We have here that $\tan \beta = v_u/v_d \approx m_t/m_b \lambda^4$, which is small. The factor λ^4 is obtained as a consequence of the Higgs and matter fields charges Q , while in chapter 2 the H and $\overline{H}$ charges were taken as zero. We recall that a value of $\tan \beta$ near 1 is an advantage for suppressing proton decay. A small range of $\tan \beta$ around one is currently disfavored by the negative results of the SUSY Higgs search at LEP [98]. Of course we could easily avoid this range, if necessary.

Remember that the zero entries in the mass matrices of the neutrino sector occur because the negatively Q -charged X field has no counterpart with positive Q -charge. Neglected small effects could partially fill up the zeroes. As explained in ref. [32] these zeroes lead to near maximal mixing also for solar neutrinos. At variance of the example of section 2.1.3, due to the different charge assignments for Ψ_{51} , the LOW solution to the solar neutrino problem emerges.

As pointed out, a problematic aspect of this zeroth order approximation to the mass matrices is the relation $m_d = m_e^T$, which is good as an order of magnitude relation because relates large left handed mixings for leptons to large right handed mixings for down quarks [32, 44, 45, 46, 47]. However the implied equalities $m_b/m_\tau = m_s/m_\mu = m_d/m_e = 1$ are good only for the third generation while need to be corrected by factors of 3 for the first two generations. As already mentioned in the discussion about the mass matrices presented in section 3.5, the necessary corrective terms can arise from the neglected terms in the expansion in $\langle Y \rangle$ of $G_r(\langle Y \rangle)_{ij}$ [57]. The higher order terms correspond to non renormalizable operators with the insertion of n factors of the 75, which break the transposition relation between m_d and m_e . For this purpose we would like the expansion parameter $\langle Y \rangle/\Lambda$ to be not too small in order to naturally provide the required factors of 3. We will present in

²In our convention Dirac mass terms are given by $\bar{L}mR$ and the light neutrinos effective mass matrix is $m_\nu m_{maj}^{-1} m_\nu^T$.

the following explicit examples of parameter choices that lead to a realistic spectrum without unacceptable fine tuning.

4.4.1 A Numerical Example

To reproduce the fermion mass spectrum we must further specialize the Yukawa couplings introduced in eq. (4.19):

$$\begin{aligned}
w_3 = & \frac{1}{4} \Psi_{10}^{AB} G_u \Psi_{10}^{CD} H^E \epsilon_{ABCDE} + \frac{\sqrt{3}}{2} \Psi_{10}^{AB} G_{\overline{50}} \Psi_{10}^{CD} H_{\overline{50}ABCD} \\
& + \sqrt{2} \Psi_{10}^{AB} G_d \Psi_{\overline{5}A} \overline{H}_B + \frac{1}{\Lambda} \sqrt{2} \Psi_{10}^{AB} F_d \Psi_{\overline{5}C} \overline{H}_D Y_{AB}^{CD} \\
& + \Psi_{\overline{5}A} G_\nu \Psi_1 H^A - \frac{1}{2} M \Psi_1 G_M \Psi_1 + \dots
\end{aligned} \tag{4.24}$$

where G_r ($r = u, d, \nu, M, \overline{50}$) is proportional to $G_r(\langle X \rangle, 0)$ of eq. (4.19). We have explicitly introduced a term linear in Y , whose couplings are described by the $\langle X \rangle$ -dependent matrix F_d . There are other terms linear in Y/Λ , not explicitly given above. In particular we may insert Y/Λ in the renormalizable term providing masses to the up type quarks. We neglect such a term since, on the one hand, the matrix G_u is already sufficient to correctly reproduce the up quark masses and, on the other hand, this operator would not significantly modify the results for proton decay. As we will show, Y/Λ is close to 0.1 in our model and higher order terms in the Y/Λ expansion can be safely neglected. The interaction term involving $H_{\overline{50}ABCD}$ does not contribute to the fermion spectrum, but it will be relevant for the proton decay amplitudes.

The term linear in Y in the previous equation is sufficient to differentiate the spectra in the charged lepton and down quark sectors. We get the following Dirac mass matrices:

$$m_{u,\nu} = y_{u,\nu} \frac{v_u}{\sqrt{2}} \quad , \quad m_{d,e} = y_{d,e} \frac{v_d}{\sqrt{2}} \quad , \tag{4.25}$$

$$\begin{aligned}
y_u &= G_u & y_\nu &= G_\nu \\
y_d &= G_d + \frac{\langle Y \rangle}{\Lambda} F_d & y_e^T &= G_d - 3 \frac{\langle Y \rangle}{\Lambda} F_d \quad ,
\end{aligned} \tag{4.26}$$

where v_u, v_d and $\langle Y \rangle$ parameterize the vevs of $H, \overline{H}$ and Y respectively. The fermion spectrum can be easily fitted by appropriately choosing the numerical values of the matrices G_u, G_d, F_d, G_ν and G_M . The most general fitting procedure would leave a large number of free parameters. Here we limit ourselves to the discussion of one particular example. In agreement with eq. (4.23), we take:

$$G_u = \begin{bmatrix} (-0.51 + 0.61i)\lambda^6 & (0.42 - 0.70i)\lambda^5 & (0.27 + 0.86i)\lambda^3 \\ (0.42 - 0.70i)\lambda^5 & (-0.39 + 0.52i)\lambda^4 & (-0.30 - 1.14i)\lambda^2 \\ (0.27 + 0.86i)\lambda^3 & (-0.30 - 1.14i)\lambda^2 & 1.39 \end{bmatrix}, \quad (4.27)$$

$$G_d = \lambda^4 \begin{bmatrix} (2.39 - 1.11i)\lambda^5 & (0.33 - 0.59i)\lambda^3 & (0.13 + 0.45i)\lambda^3 \\ (0.87 + 0.55i)\lambda^4 & (2.76 + 0.89i)\lambda^2 & (0.69 - 0.51i)\lambda^2 \\ (-1.50 + 0.94i)\lambda^2 & (0.45 + 1.78i) & 1.94 \end{bmatrix}, \quad (4.28)$$

$$\frac{\langle Y \rangle}{\Lambda} F_d = \lambda^4 \begin{bmatrix} (0.38 - 0.18i)\lambda^5 & (-0.16 - 0.06i)\lambda^3 & (0.07 + 0.04i)\lambda^3 \\ (-0.08 - 0.05i)\lambda^4 & (-0.20 - 0.15i)\lambda^2 & (0.15 - 0.11i)\lambda^2 \\ (-0.09 + 0.12i)\lambda^2 & (0.07 + 0.14i) & -0.19 \end{bmatrix}, \quad (4.29)$$

$$G_\nu = \begin{bmatrix} (-0.78 - 0.19i)\lambda^3 & (0.52 - 0.34i)\lambda & (1.38 + 0.39i)\lambda^2 \\ (-1.23 - 0.34i)\lambda & 0 & (1.04 + 1.31i) \\ (0.45 + 1.18i)\lambda & 0 & 0.8 + 1.2i \end{bmatrix}, \quad (4.30)$$

$$G_M = \begin{bmatrix} (1.50 + 0.55i)\lambda^2 & (1.41 + 1.19i) & (0.35 - 1.53i)\lambda \\ (1.41 + 1.19i) & 0 & 0 \\ (0.35 - 1.53i)\lambda & 0 & 1.26 + 1.48i \end{bmatrix}. \quad (4.31)$$

While the generic pattern of G_u , G_d , F_d , G_ν and G_M is dictated by the $U(1)$ flavour symmetry, the precise values of the coefficients multiplying the powers of λ are chosen to reproduce the data. We take $\lambda \equiv \langle X \rangle / \Lambda = 0.25$, $\tan \beta = 1.5$ and $M = 0.9 \cdot 10^{15}$ GeV. In $SU(5)$ the matrix G_u contains two additional phases [86], ϕ_1 and ϕ_2 that have been set to zero in eq. (4.27). These phases do not affect the fermion spectrum but enter the proton decay amplitude. When discussing the proton decay we will analyze also the dependence on ϕ_1 and ϕ_2 .

From the above matrices we obtain, at the unification scale:

$$\begin{aligned} m_t &= 200 \text{ GeV} & m_c &= 0.27 \text{ GeV} & m_u &= 0.9 \text{ MeV} \\ m_b &= 1.0 \text{ GeV} & m_s &= 26 \text{ MeV} & m_d &= 1.1 \text{ MeV} \\ m_\tau &= 1.1 \text{ GeV} & m_\mu &= 71 \text{ MeV} & m_e &= 0.34 \text{ MeV} \end{aligned} \quad (4.32)$$

$$|V_{us}| = 0.22 \quad |V_{ub}| = 0.0022 \quad |V_{cb}| = 0.052 \quad J = 1.9 \cdot 10^{-5}, \quad (4.33)$$

where J is the CP-violating Jarlskog invariant. In the neutrino sector, we find:

$$m_1 = 0.81 \cdot 10^{-3} \text{ eV}, \quad m_2 = 0.88 \cdot 10^{-3} \text{ eV}, \quad m_3 = 0.061 \text{ eV}. \quad (4.34)$$

More precisely:

$$\Delta m_{sol}^2 \equiv m_2^2 - m_1^2 = 1.1 \cdot 10^{-7} \text{ eV}^2, \quad \Delta m_{atm}^2 \equiv m_3^2 - m_2^2 = 3.7 \cdot 10^{-3} \text{ eV}^2. \quad (4.35)$$

The neutrino mixing angles are

$$\theta_{12} \sim \frac{\pi}{4}, \quad \theta_{23} \sim \frac{\pi}{4}, \quad \theta_{13} = 0.06. \quad (4.36)$$

Until now we have not specified the value of $\langle Y \rangle / \Lambda$. We know that $\langle Y \rangle$ should be around $M_{GUT}^{(0)}$. The cut-off Λ cannot be too close to $\langle Y \rangle$, otherwise most of the spectrum of the model would lie beyond the cut-off. At the same time Λ cannot be too large: it is bounded from above by the scale at which the $SU(5)$ gauge coupling blows up, which, as we see from eq. (4.18), occurs more or less one order of magnitude below the Planck mass. This is welcome. If, for instance, we take $\langle Y \rangle / \Lambda = 0.05 - 0.1$, it is reasonable to neglect multiple insertions of Y in the Yukawa operators. At the same time, in the example given in eq. (4.29), the coefficients of the powers of λ in F_d remain of order one, even if $\langle Y \rangle / \Lambda$ is as small as 0.05. A too large cut-off would have been ineffective in separating down quarks from charged leptons unless we had chosen unnaturally large coefficients in the allowed operators. What is usually considered a bad feature of the missing partner mechanism - the lack of perturbativity before the Planck mass - turns out here to be an advantage to provide a correct description of the fermion spectrum.

In the neutrino sector, we obtain a bimaximal neutrino mixing with the so-called LOW solution to the solar neutrino problem³. Within the same $U(1)$ flavour symmetry considered here we could as well reproduce the quasi-vacuum oscillation solution, by appropriately tuning the order one coefficients in G_ν and G_M . We recall that the value of M required to fit the observed atmospheric oscillations is probably somewhat small in the context of $SU(5)$, where a larger scale, closer to the cut-off Λ , is expected. This feature might be improved by embedding the model in $SO(10)$ where M is directly related to the $B - L$ breaking scale.

In conclusion, the known fermion spectrum can be reproduced starting from a superpotential with order one dimensionless coefficients. Mass matrix elements for charged leptons and down quarks match only within 10 – 20%, due to $\langle Y \rangle / \Lambda \approx 0.1$, and this produces the required difference between the two sectors. The neutrino mixing is necessarily bimaximal in our model, with either the LOW or the vacuum oscillation solution to the solar neutrino problem.

³As pointed out, in chapter 1, latest preliminary results from Super-Kamiokande [77], including constraints from day-night spectra, seem in fact to prefer bimaximal neutrino mixing and to revamp interest in the LOW solution.

4.5 Proton Decay

Similarly to the case of minimal $SU(5)$, we expect that the main contribution to proton decay comes from the dimension five operators [79] originating at the grand unified scale when the colour triplet superfields are integrated out [78, 80, 81]. We denote the colour triplets contained in $H, \bar{H}, H_{50}, H_{\bar{50}}$ by H_{3u}, H_{3d}, H'_{3d} and H'_{3u} , respectively. The part of the superpotential depending on these superfields reads:

$$\begin{aligned}
w = & H_{3u} \left[-\frac{1}{2} Q G_u Q + U^c G_u E^c \right] \\
& + H_{3d} \left[Q \hat{C} L + U^c \hat{D} D^c \right] \\
& + H'_{3u} \left[\frac{1}{4} Q G_{\bar{50}} Q + U^c G_{\bar{50}} E^c \right] \\
& - 4\sqrt{3}c_2 \langle Y \rangle H'_{3d} H_{3u} - 4\sqrt{3}c_3 \langle Y \rangle H_{3d} H'_{3u} + c_4 \langle X \rangle H'_{3d} H'_{3u} + \dots \quad (4.37)
\end{aligned}$$

where

$$\hat{C} = -G_d - \frac{\langle Y \rangle}{\Lambda} F_d, \quad \hat{D} = G_d - \frac{\langle Y \rangle}{\Lambda} F_d \quad (4.38)$$

and Q, L, U^c, D^c and E^c denote as usual the chiral multiplets associated to the three fermion generations. Notice that, at variance with the minimal $SU(5)$ model, an additional interaction term depending on H'_{3u} is present, which originates from the interaction of $\bar{50}$ with the 10 matter fields. In Appendix C we give a short discussion on its properties and justify the choice of the numerical coefficient $\sqrt{3}/2$ in eq (4.24). By integrating out the colour triplets we obtain the following effective superpotential:

$$w_{eff} = \frac{1}{m_T} \left[Q \hat{A} Q Q \hat{C} L + U^c \hat{B} E^c U^c \hat{D} D^c \right] + \dots \quad (4.39)$$

where m_T has been defined in eq. (4.8),

$$\hat{B} = \left(G_u + \frac{4\sqrt{3}c_2 \langle Y \rangle}{c_4 \langle X \rangle} G_{\bar{50}} \right), \quad -2\hat{A} = \left(G_u - \frac{2\sqrt{3}c_2 \langle Y \rangle}{c_4 \langle X \rangle} G_{\bar{50}} \right) \quad (4.40)$$

and dots stand for terms that do not violate baryon or lepton number. Minimal $SU(5)$ is recovered by setting $G_{\bar{50}} = F_d = 0$. As discussed, in that case the matrices $\hat{A}, \hat{B}, \hat{C}$ and $\hat{D}$ are determined by G_u (in which now also ϕ_1 and ϕ_2 play a role) and G_d and therefore strictly related to the fermionic spectrum [78]. In our case we have a distortion due to the terms proportional to $G_{\bar{50}}$ and F_d . These distortions have

different physical origins. On the one hand the terms containing F_d are required to avoid the rigid mass relation of minimal $SU(5)$. They are suppressed by $\langle Y \rangle / \Lambda$ and we expect a mild effect from them [89]. On the other hand the terms containing G_{50} are a consequence of the missing doublet mechanism and they might be as important as those of the minimal model.

To estimate the proton decay rates we should specify some important parameters. First of all the mass m_T . From the discussion of the threshold corrections we know that a large value for m_T is preferred in our model. We should however check that this value can be obtained with reasonable choices of the parameters at our disposal. The unification conditions constrain $\langle Y \rangle$ in a small range around 10^{16} GeV. More precisely, after the inclusion of threshold corrections and two loop effects, eq. (4.15) becomes a relation among the masses of the super-heavy particles and the leading order quantity $M_{GUT}^{(0)}$. This relation limits the allowed range for the heavy gauge vector bosons and indirectly, through eq. (C.41), pushes the $SU(5)$ breaking vev $\langle Y \rangle$ close to 10^{16} GeV. Then the cut-off scale and the X vev are fixed by the phenomenological requirements $\langle Y \rangle \approx 0.05 \Lambda$ and $\lambda \equiv \langle X \rangle / \Lambda = 0.25$. Large values of m_T could be obtained either by taking $m_{T_1} m_{T_2}$ large or by choosing a small m_ϕ . This last possibility is however not practicable, since the gauge coupling α_5 blows up at approximately $20 m_\phi$: it is not reasonable to push m_ϕ below $5 \cdot 10^{15}$ GeV. The only remaining freedom to obtain the desired large value for m_T is represented by the coefficients c_2 and c_3 that, however, cannot be taken arbitrarily large. We should also check that all the heavy spectrum remains below Λ and this requirement imposes a further constraint on our parameters. A choice that respects all these requirements is provided by ⁴:

$$c_1 = 0.035 \quad , \quad c_2 = c_3 = 2.8, \quad c_4 = 0.7 \quad , \quad (4.41)$$

$$\langle X \rangle = 3.0 \cdot 10^{16} \text{ GeV}, \quad \langle Y \rangle = 5.7 \cdot 10^{15} \text{ GeV} \quad (\Lambda = 1.2 \cdot 10^{17} \text{ GeV}) \quad , \quad (4.42)$$

which leads to:

$$M_{GUT} = 2.9 \cdot 10^{16} \text{ GeV} \quad , \quad m_\phi = 2.0 \cdot 10^{16} \text{ GeV} \quad , \quad (4.43)$$

$$m_{T_1} = 1.2 \cdot 10^{17} \text{ GeV} \quad , \quad m_{T_2} = 1.0 \cdot 10^{17} \text{ GeV} \quad , \quad m_T = 6 \cdot 10^{17} \text{ GeV} \quad . \quad (4.44)$$

The heavy sector of the particle spectrum is displayed in fig. 4.3.

To evaluate the loop function f we also need the spectrum of the supersymmetric particles. As an example we take here the same spectrum considered in Table B.1,

⁴The values of the coefficients c_2 and c_3 here adopted raise doubts on the validity of the perturbative approach that has been exploited in several aspects of the present analysis. We adhere to this choice also to show the difficulty met to obtain acceptable phenomenological results within a not too complicated scheme.




Figure 4.3: Heavy sector of the spectrum.

with $m_{SUSY} = 250$ GeV. This leads to a squark mass of about 800 GeV, a slepton mass of approximately 350 GeV a wino mass of 250 GeV and a charged higgsino mass of 125 GeV.

The matrix $G_{\overline{50}}$ is not directly related to any accessible observable quantity and the only constraint we have on it comes from the $U(1)$ flavour symmetry that requires the following general pattern:

$$G_{\overline{50}} = \begin{bmatrix} \lambda^7 & \lambda^6 & \lambda^4 \\ \lambda^6 & \lambda^5 & \lambda^3 \\ \lambda^4 & \lambda^3 & \lambda \end{bmatrix} . \quad (4.45)$$

With the above numerical values we also obtain:

$$\left| \frac{2 \sqrt{3} c_2 \langle Y \rangle}{c_4 \langle X \rangle} \right| \simeq 2.7 . \quad (4.46)$$

The texture for $G_{\overline{50}}$ has an overall suppression factor λ compared to G_u . Therefore the contribution of $G_{\overline{50}}$ to the matrices $\hat{A}$ and $\hat{B}$ is comparable or even slightly larger than the minimal contribution provided by G_u . The dominant operators are then the ones already described and the calculation can be carried out following Table A.2.

At variance of the minimal case, here the amplitude with G_u and the one with $G_{\overline{50}}$ interfere. This interference can be either constructive or destructive, depending on the relative phases between the two terms. We have scanned several examples for $G_{\overline{50}}$, obtained by generating random coefficients for the order one variables in eq. (4.45). By keeping fixed all the remaining parameters we obtain a proton decay rate in the range $2 \cdot 10^{32} \div 4 \cdot 10^{34}$ yr for the channel $p \rightarrow K^+ \bar{\nu}$ and a rate between $3 \cdot 10^{32}$ yr and $5 \cdot 10^{34}$ yr for the channel $p \rightarrow \pi^+ \bar{\nu}$. For comparison, considering the same choice of parameters but setting $G_{\overline{50}} = F_d = 0$, we obtain $9 \cdot 10^{32}$ yr and $2 \cdot 10^{33}$ yr respectively, for the above channels. The present 90% CL bound on $\tau/BR(p \rightarrow K^+ \bar{\nu})$ is $1.9 \cdot 10^{33}$ yr [77]. These estimates have been obtained by setting to zero the two physical phases ϕ_1, ϕ_2 contained in the matrix G_u . These additional parameters may increase the uncertainty on the proton lifetime. For instance, in

minimal $SU(5)$, the proton decay rates for the channels considered above, change by about one order of magnitude when ϕ_1 and ϕ_2 are freely varied between 0 and 2π . Even when the inverse decay rates for the channels $K^+\bar{\nu}$ and $\pi^+\bar{\nu}$ are as large as 10^{34} yr, they remain the dominant contribution to the proton lifetime. Indeed, since the heavy vector boson mass, M_{GUT} , is equal to $2.9 \cdot 10^{16}$ GeV in our model, the dimension 6 operators provide an inverse decay rate for the channel $e^+\pi^0$ larger than 10^{36} yr.

The effective theory considered here breaks down at the cut-off scale Λ . We expect additional non-renormalizable operators contributing to proton decay amplitudes from the physics above the cut-off. We have already stressed that, by assuming dimensionless coupling constants of order one, in unified models without flavour symmetries the proton lifetime induced by these operators is unacceptably short, even when $\Lambda = M_{Pl}$ [83, 87]. In our case these contributions are adequately suppressed by the $U(1)$ symmetry. If we compare the amplitude A_{nr} induced by the new non-renormalizable operators with the amplitude A_3 coming from the triplet exchange, we obtain, for the generic decay channel,

$$\frac{A_{nr}}{A_3} \approx \frac{48 c_2 c_3}{c_4} \left(\frac{\langle Y \rangle}{\Lambda} \right)^2 \approx 1 \quad . \quad (4.47)$$

This supports the conclusion that a proton lifetime range considerably larger than the one estimated in minimal models is expected in our case.

Conclusions

We have considered the problem of obtaining the large atmospheric neutrino mixing in the framework of GUT models.

First, we have analyzed the most interesting mechanisms which provide this large mixing.

Usually the large atmospheric mixing arises because of the presence of large mixings in the Dirac and/or Majorana right mass matrices. In GUT models this situation is naturally realized by exploiting the possible presence of large right mixings for down quarks, which correspond to large left mixing for charged leptons.

We have pointed out an alternative mechanism, which does not require the presence of any large mixings. In fact, even when both the Dirac and Majorana right matrices possess very small mixings, the large atmospheric mixing entirely arises from the see-saw mechanism if two conditions apply, that is if the spectrum of right Majorana masses is sufficiently splitted and if the left handed ν_μ and ν_τ couple with comparable strength to the right handed ν_μ^c . As opposite to the first mechanism, this situation can be realized even if all left and right mixings of quarks and leptons are very small. This feature is particularly attractive from the theoretical point of view, because it implies that this mechanism is well compatible with those GUT models where right and left mixings are similar.

Despite the fact that GUT has been studied since a long time, still a lot of its aspects require a better understanding. It would be really improbable for this idea to survive if inside it there were not something true. But it is clear that minimal versions of SUSY GUTs are already ruled out because unrealistic. We have addressed the question whether it is possible to construct a not too complicated, relatively complete and realistic model which can serve as benchmark to be compared with experiment. This model must also correctly reproduce neutrino masses and mixings.

We have then constructed an example of SUSY $SU(5)$ GUT model, with an additional $U(1)$ flavour symmetry, which is not plagued by the need for large amounts of fine tunings, like those associated with doublet-triplet splitting in the minimal model, and leads to an acceptable phenomenology. This includes coupling unification with a value of $\alpha_3(m_Z)$ in much better agreement with the data than in the

minimal version, an acceptable pattern for fermion masses and mixing angles, also including neutrino masses and mixings, and the possibility of a slower proton decay than in the minimal version, compatible with the present limits (in particular the limit from Super-Kamiokande of about $2 \cdot 10^{33}$ yr for the channel $p \rightarrow K^+ \bar{\nu}$). In the neutrino sector the preferred solution is one with nearly maximal mixing both for atmospheric and solar neutrinos.

The $U(1)$ flavour symmetry plays a crucial role by protecting the light doublet Higgs mass from receiving large mass contributions from higher dimension operators, by determining the observed hierarchy of fermion masses and mixings and by providing an adequate suppression for non renormalizable operators inducing proton decay. Of course, the $U(1)$ symmetry can only reproduce the order of magnitude of masses and mixings, while more quantitative relations among masses and mixings can only arise from a non abelian flavour symmetry.

A remarkable feature of the model is that the presence of the representations 50, $\overline{50}$ and 75, demanded by the Missing Doublet Mechanism for the solution of the doublet-triplet splitting problem, directly produces, through threshold corrections at M_{GUT} from the 75, a decrease of the value of $\alpha_3(m_Z)$ that corresponds to coupling unification and an increase of the effective mass that mediates proton decay by a factor of typically 20-30. As a consequence the value of the strong coupling is in better agreement with the experimental value and the proton decay rate is smaller by a factor 400-1000 than in the minimal model. The presence of these large representations also has the consequence that the asymptotic freedom of $SU(5)$ is spoiled and the associated gauge coupling becomes non perturbative below M_{Pl} . We argue that this property far from being unacceptable can actually be useful to obtain better results for fermion masses and proton decay.

Clearly such a model is not unique: our version is the simplest realistic model that we could construct. We think it is interesting because it proves that SUSY $SU(5)$ GU is not excluded and offers a benchmark for comparison with experiment. For example, even including all possible uncertainties, it is difficult in this class of models to avoid the conclusion that proton decay must occur with a rate which is only a factor 10-50 from the present bounds. Failure to observe such a signal would require some additional specific mechanism in order to further suppress the decay rate.

Finally, it is a generic feature of realistic models that the region between M_{GUT} and M_{Pl} becomes populated by many states with different thresholds and also non perturbative phenomena occur. This suggests that the reality can be more complicated than the neat separation between the Grand Unification and the string regime which is postulated in the simplest toy models of GUTs.

There is then the need for further ideas on the theoretical side. It would be

of crucial importance to actually understand the origin of the flavour symmetry. The simple $U(1)$ used in the present model is satisfactory at the level of orders of magnitude, but is not able to really predict the observed values of masses and mixing. Maybe the flavour has a deeper origin, in a fundamental theory of particles that includes also gravity beyond the strong and electroweak interactions.

Acknowledgements

I warmly thank Guido Altarelli and Ferruccio Feruglio for their invaluable and pleasant collaboration. I would like to thank also Giovanni Costa, Gianfranco Sartori, Gianpaolo Valente and Fabio Zwirner for useful discussions.

Bibliography

- [1] Super-Kamiokande Collaboration, Y. Fukuda et al., Phys. Rev. Lett. 81 (1998) 1562, hep-ex/9807003; Phys. Lett. B433 (1998) 9; Phys. Lett. B436 (1998) 33; Phys. Rev. Lett. 82 (1999) 2644.
- [2] B. Pontecorvo, Sov. Phys. JETP 26 (1968) 984.
- [3] Z. Maki, M. Nakagawa and S. Sakata, Prog. Theor. Phys. 28 (1962) 870.
- [4] Soudan-2 Collaboration, W. W. M. Allison et al., Phys. Lett. B449 (1999) 137.
- [5] MACRO Collaboration, M. Ambrosio et al., Phys. Lett. B434 (1998) 451.
- [6] Kamiokande Collaboration, Phys. Rev. Lett. 81 (1998) 2016.
- [7] CHOOZ Collaboration, M. Apollonio et al., Phys. Lett. B 420 (1998) 397; Phys. Lett. B466 (1999) 415.
- [8] F. Boehm et al., hep-ex/0003022.
- [9] G. L. Fogli, E. Lisi, A. Marrone and G. Scioscia, Phys. Rev. D59 (1999) 033001.
- [10] V. Barger, B. Kayser, J. Learned, T. Weiler and K. Whisnant, hep-ph/0008019.
- [11] G. L. Fogli, E. Lisi and A. Marrone, hep-ph/0009299; G. L. Fogli, E. Lisi, D. Montanino and A. Palazzo, Phys. Rev. D61 (2000) 073009, hep-ph/9910387.
- [12] V. Barger, J. G. Learned, S. Pakvasa and T. J. Weiler, Phys. Rev. Lett. 82 (1999) 2640.
- [13] E. Lisi, A. Marrone and D. Montanino, Phys. Rev. Lett. 85 (2000) 1166, hep-ph/0002053.

- [14] R. Barbieri, P. Creminelli and A. Strumia, Nucl. Phys. B585 (2000) 28, hep-ph/0002199.
- [15] G. L. Fogli, E. Lisi, A. Marrone and G. Scioscia, Nucl. Instrum. Meth. A451 (2000) 10.
- [16] K2K Collaboration, T. Ishida, hep-ex/0008047.
- [17] MINOS Collaboration, E. Ables et al., FERMILAB-PROPOSAL-P-875.
- [18] A. Rubbia, hep-ex/0008071.
- [19] For a short review see A. Cervera et al., hep-ph/0007281, and references therein.
- [20] The data from Homestake, Gallex, Sage, Kamiokande and Superkamiokande on solar neutrinos are summarised and the list of references given in J. N. Bahcall, P. I. Krastev and A. Yu Smirnov, hep-ph/9807216; Super-Kamiokande Collaboration, Y. Fukuda et al., Phys. Rev. Lett. 82 (1999) 1810; Phys. Rev. Lett. 82 (1999) 2430.
- [21] J. N. Bahcall, S. Basu and M. H. Pinsonneault, Phys. Lett. B433 (1998) 1.
- [22] SNO Collaboration, J. Boger et al., Nucl. Instrum. Meth. A449 (2000) 172.
- [23] BOREXINO Collaboration, F. P. Calaprice, Nucl. Phys. Proc. Suppl. 87 (2000) 180.
- [24] KamLAND Collaboration, L. de Braekeleer, Nucl. Phys. Proc. Suppl. 87 (2000) 312.
- [25] C. Athanassopoulos et al., Phys. Rev. Lett. 77 (1996) 3082; nucl-ex/9706006, nucl-ex/9709006.
- [26] B. Armbruster et al., Phys. Rev. C57 (1998) 3414; hep-ex/0008002.
- [27] G. L. Fogli, E. Lisi, A. Marrone and D. Montanino, hep-ph/0009269.
- [28] M. Gell-Mann, P. Ramond and R. Slansky in Supergravity, ed. P. van Nieuwenhuizen and D. Z. Freedman, North-Holland, Amsterdam, 1979, p.315; T. Yanagida, in Proceedings of the Workshop on the unified theory and the baryon number in the universe, ed. O. Sawada and A. Sugamoto, KEK report No. 79-18, Tsukuba, Japan, 1979. See also R. Mohapatra and G. Senjanovic, Phys. Rev. Lett. 44, 912 (1980).

- [29] N. Cabibbo, Phys. Lett. B72 (1978) 333.
- [30] A. De Rujula, M. B. Gavela and P. Hernandez, Nucl. Phys. B547 (1999) 21, hep-ph/9811390.
- [31] S.M. Bilenky and C. Giunti, hep-ph/9802201; V. Barger, S. Pakvasa, T.J. Weiler and K. Whisnant, hep-ph/9806387; A.J. Baltz, A.S. Goldhaber and M. Goldhaber, hep-ph/9806540; V. Barger, T.J. Weiler and K. Whisnant, hep-ph/9807319.
- [32] G. Altarelli and F. Feruglio, Phys. Lett. B439 (1998) 112, hep-ph/9807353; JHEP 11 (1998) 21, hep-ph/9809596; Phys. Lett. B451 (1999) 388, hep-ph/9812475.
- [33] G. Altarelli and F. Feruglio, Phys. Rep. 320 (1999) 295, hep-ph/9905536 and references therein.
- [34] L. Baudis et al., Phys. Rev. Lett. 83 (1999) 41, hep-ex/9902014.
- [35] F. Vissani, hep-ph/9708483; H. Georgi and S.L. Glashow, Phys. Rev. D61 (2000) 097301, hep-ph/9808293.
- [36] A. Ioannisian, J. W. F. Valle, Phys. Lett. B332 (1994) 93, hep-ph/9402333; M. Fukugita, M. Tanimoto, T. Yanagida, Phys. Rev. D57 (1998) 4429, hep-ph/9709388; M. Tanimoto, hep-ph/9807283 and hep-ph/9807517; H. Fritzsch, Z. Xing, hep-ph/9808272; R. N. Mohapatra, S. Nussinov, hep-ph/9808301 and hep-ph/9809415; M. Fukugita, M. Tanimoto, T. Yanagida, hep-ph/9809554; Yue-Liang Wu, hep-ph/9810491; J. I. Silva-Marcos, hep-ph/9811381; H. Fritzsch, Z. Xing, hep-ph/9903499; S. K. Kang and C. S. Kim, hep-ph/9811379; C. Wetterich, hep-ph/9812426; R. Barbieri, L. J. Hall, G. L. Kane and G. G. Ross, hep-ph/9901228.
- [37] J. Ellis and S. Lola, hep-ph/9904279; J. A. Casas, J. R. Espinosa, A. Ibarra and I. Navarro, Nucl. Phys. B556 (1999) 3, hep-ph/9904395; Nucl. Phys. B569 (2000) 82, hep-ph/9905381; hep-ph/9910420; hep-ph/0004166; E. Ma, hep-ph/9907400; P. H. Chankowski, W. Krolikowski, S. Pokorski, hep-ph/9910231; S. Lola, hep-ph/0005093.
- [38] R. Barbieri, G. G. Ross and A. Strumia, JHEP 9910 (1999) 020, hep-ph/9906470.
- [39] R. Barbieri, L. J. Hall, D. Smith, A. Strumia and N. Weiner, JHEP 9812 (1998) 017, hep-ph/9807235.

- [40] A. Ghosal, hep-ph/9905470.
- [41] J. A. Casas, J. R. Espinosa, A. Ibarra and I. Navarro, JHEP 9909 (1999) 015, hep-ph/9906281; Nucl. Instrum. Meth. A451 (2000) 69.
- [42] K. S. Babu, J. C. Pati and F. Wilczek, Nucl. Phys. B566 (2000) 33, hep-ph/9812538.
- [43] S. F. King, Phys. Lett. B439 (1998) 350, hep-ph/9806440; Nucl. Phys. B562 (1999) 57, hep-ph/9904210; hep-ph/9912492; S. Davidson and S. F. King, Phys. Lett. B445 (1998) 191, hep-ph/9808333; S. F. King and M. Oliveira, hep-ph/0009287.
- [44] G. C. Branco, L. Lavoura and F. Mota, Phys. Rev. D39 (1989) 3443; G. C. Branco and J. I. Silva-Marcos, Phys. Lett. B331 (1994) 390.
- [45] Z. Berezhiani and A. Rossi, Phys. Lett. B367 (1996) 219; JHEP 9903 (1999) 002, hep-ph/9811447; S. Lola and G. G. Ross, Nucl. Phys. B553 (1999) 81, hep-ph/9902283.
- [46] C. H. Albright and S. M. Barr, Phys. Rev. D58 (1998) 013002, hep-ph/9712448; Phys. Lett. B452 (1999) 287, hep-ph/9901318; C. H. Albright, K. S. Babu and S. M. Barr, Phys. Rev. Lett. 81 (1998) 1167, hep-ph/9802314.
- [47] Z. Berezhiani and Z. Tavartkiladze, Phys. Lett. B409 (1997) 220, hep-ph/9612232; R. Barbieri, L. Giusti, L. Hall and A. Romanino, Nucl. Phys. B550 (1999) 32-40, hep-ph/9812239; N. Haba and N. Okamura, Eur. Phys. J. C14 (2000) 347-365, hep-ph/9906481.
- [48] G. Altarelli, F. Feruglio and I. Masina, Phys. Lett. B472 (2000) 382, hep-ph/9907532.
- [49] A. Y. Smirnov, Phys. Rev. D48 (1993) 3264; M. Jezabek and Y. Sumino, Phys. Lett. B440 (1998) 327, hep-ph/9807310; Phys. Lett. B457 (1999) 139, hep-ph/9904382.
- [50] G. G. Ross, "Grand Unified Theories", Benjamin, 1985; R. N. Mohapatra, "Unification and Supersymmetry" Springer-Verlag, 1986; C. Kounnas, A. Masiero, D. V. Nanopoulos and K. A. Olive, "Grand Unification with and without Supersymmetry and Cosmolgical Implications", World Sci. 1984.
- [51] C. Froggatt and H. B. Nielsen, Nucl. Phys. B147 (1979) 277.

- [52] M. Leurer, Y. Nir and N. Seiberg, Nucl. Phys. B420 (1994) 468, hep-ph/9310320.
- [53] P. Binetruy, N. Irges, S. Lavignac and P. Ramond, Phys. Lett. B403 (1997) 38, hep-ph/9612442; N. Irges and S. Lavignac, Phys. Lett. B424 (1998) 293, hep-ph/9712239.
- [54] W. Buchmuller, D. Delepine and F. Vissani, Phys. Lett. B459 (1999) 171, hep-ph/9904219.
- [55] W. Buchmuller, D. Delepine and L. T. Handoko, Nucl. Phys. B576 (2000) 445, hep-ph/9912317.
- [56] E. Kh. Akhmedov, G. C. Branco, F. R. Joaquim and J. I. Silva-Marcos, Phys. Lett. B498 (2001) 237.
- [57] J. Ellis and M. K. Gaillard, Phys. Lett. B88 (1979) 315.
- [58] R. Barbieri, P. Creminelli and A. Romanino, Nucl. Phys. B559 (1999) 17, hep-ph/9903460; T. Blazek, S. Raby and K. Tobe, hep-ph/9903340; B. Stech, hep-ph/9905440.
- [59] J. C. Pati and A. Salam, Phys. Rev. D10 (1974) 275; H. Georgi and S. Glashow, Phys. Rev. Lett. 32 (1974) 438; A. Buras, J. Ellis and D. Nanopoulos, Nucl. Phys. B135 (1978) 66; H. Georgi and D. Nanopoulos, Nucl. Phys. B155 (1979) 52.
- [60] S. Dimopoulos, S. Raby and F. Wilczek, Phys. Rev. D24 (1981) 1681; W. J. Marciano and G. Senjanovic, Phys. Rev. D25 (1982) 3092.
- [61] The Soudan 2 collaboration, Phys. Lett. B427 (1998) 217, hep-ex/980303; The Super-Kamiokande Collaboration: M. Shiozawa et al., Phys. Rev. Lett. 81 (1998) 3319, hep-ex/9806014; Super-Kamiokande Collaboration: Y. Hayato, M. Earl et al., Phys. Rev. Lett. 83 (1999) 1529, hep-ex/9904020.
- [62] S. Dimopoulos and H. Georgi, Nucl. Phys. B150 (1981) 193; M. Sakai, Z. Phys. C11 (1981) 153; E. Witten, Nucl. Phys. B188 (1981) 573.
- [63] H. Georgi, Phys. Lett. B108 (1982) 283; A. Masiero, D. V. Nanopoulos, K. Tamvakis and T. Yanagida, Phys. Lett. B115 (1982) 380; B. Grinstein, Nucl. Phys. B206 (1982) 387.

- [64] Y. Yamada, Z. Phys. C60 (1993) 83; K. Hagiwara and Y. Yamada, Phys. Rev. Lett. 70 (1993) 709.
- [65] J. Hisano, T. Moroi, K. Tobe and T. Yanagida, Phys. Lett. B342 (1995) 138; J. Hisano, Y. Nomura and T. Yanagida, Prog. Theor. Phys. 98 No. 6 (1997) 1385.
- [66] A. Masiero, D. V. Nanopoulos, K. Tamvakis and T. Yanagida, Z. Phys. C17 (1983) 33; J. L. Lopez and D. V. Nanopoulos, Phys. Rev. D35 (1996) 2670, hep-ph/9508253.
- [67] Z. Berezhiani and Z. Tavartkiladze, Phys. Lett. B396 (1997) 150, hep-ph/9611277.
- [68] L. Randall and C. Csaki, hep-ph/9508208.
- [69] E. Witten, Phys. Lett. B105 (1981) 267; D. V. Nanopoulos and K. Tamvakis, Phys. Lett. B113 (1982) 151.
- [70] J. Polchinski and L. Susskind, Phys. Rev. D26 (1982) 3661; H. P. Nilles, M. Srednicki and D. Wyler, Phys. Lett. B124 (1982) 337; A. Lahanas, Phys. Lett. B124 (1982) 341.
- [71] S. Dimopoulos and F. Wilczek, NSF-ITP-82-07; M. Srednicki, Nucl. Phys. B202 (1982) 327.
- [72] K. S. Babu and S. M. Barr, Phys. Rev. D48 (1993) 5354; Phys. Rev. D50 (1994) 3529; Phys. Rev. D51 (1995) 2463; K. S. Babu and R. N. Mohapatra, Phys. Rev. Lett. 74 (1995) 2418.
- [73] G. Dvali and S. Pokorski, Phys. Lett. B379 (1996) 126, hep-ph/9601358; Z. Berezhiani and Z. Tavartkiladze, Phys. Lett. B409 (1997) 220, hep-ph/9612232.
- [74] C. H. Albright and S. M. Barr, hep-ph/0002155.
- [75] K. Inoue, A. Kakuto and H. Takano, Prog. Theor. Phys. 75 (1986) 664; A. Anselm and A. Johansen, Phys. Lett. B200 (1988) 331; R. Barbieri, G. Dvali and A. Strumia, Nucl. Phys. B391 (1993) 487.
- [76] R. Barbieri, G. Dvali and M. Moretti, Phys. Lett. B312 (1993) 137; R. Barbieri, G. Dvali, A. Strumia, Z. Berezhiani and L. Hall, Nucl. Phys. B432 (1994) 49; Z. Berezhiani, C. Csaki and L. Randall, Nucl. Phys. B444 (1995) 61,

- hep-ph/9501336; C. Csaki and L. Randall, Nucl. Phys. B466 (1996) 41, hep-ph/9512278.
- [77] Latest results from Super-Kamiokande have been presented in: Y. Totsuka, Talk at the “Susy 2K” Conference, CERN, June 2000.
- [78] J. Ellis, D.V. Nanopoulos and S. Rudaz, Nucl. Phys. B202 (1982) 43.
- [79] N. Sakai and T. Yanagida, Nucl. Phys. B197 (1982) 533; S. Weinberg, Phys. Rev. D26 (1982) 287.
- [80] S. Dimopoulos, S. Raby and F. Wilczek, Phys. Lett. B112 (1982) 133.
- [81] J. Hisano, H. Murayama and T. Yanagida, Nucl. Phys. B402 (1993) 46-84, hep-ph/9207279.
- [82] R. Arnowitt, A. H. Chamseddine and P. Nath, Phys. Rev. D32 (1985) 2348-2358.
- [83] J. Ellis, J. S. Hagelin, D. V. Nanopoulos and K. Tamvakis, Phys. Lett. B124 (1983) 484.
- [84] V. M. Belyaev and M. I. Vysotsky, Phys. Lett. B127 (1983) 215.
- [85] T. Goto and T. Nihei, Phys. Rev. D59 (1999) 115009, hep-ph/9808255; hep-ph/9909251.
- [86] J. Ellis, M.K. Gaillard and D. V. Nanopoulos, Phys. Lett. B88 (1979) 320; R. Arnowitt, A. H. Chamseddine and P. Nath, Phys. Lett. B156 (1985) 215-219.
- [87] J. Ellis, D. V. Nanopoulos and K. Tamvakis, Phys. Lett. B121 (1983) 123; H. Murayama and D. B. Kaplan, Phys. Lett. B336 (1994) 221-228; H. Murayama, hep-ph/9610419.
- [88] See JLQCD Collaboration: S. Aoki et al., Phys. Rev. D62 (2000) 014506, hep-lat/9911026 and references therein.
- [89] K. S. Babu and S. M. Barr, Phys. Lett. B381 (1996) 137, hep-ph/9506261; P. Nath, Phys. Rev. Lett. 76 (1996) 2218-2221, hep-ph/9512415; Phys. Lett. B381 (1996) 147, hep-ph/9602337; P. Nath and R. Arnowitt, Phys. Atom. Nucl. 61 (1998) 975-982, hep-ph/9708469; Z. Berezhiani, Z. Tavartkiladze and M. Vysotsky, hep-ph/9809301.
- [90] H. Georgi and C. Jarlskog, Phys. Lett. B86 (1979) 297.

- [91] K. Hagiwara and N. Okamura, Nucl. Phys. B548 (1999) 60, hep-ph/9811495.
- [92] The European Physical Journal C 15 Number 1-4 (2000) 1.
- [93] V. Lucas and S. Raby, Phys. Rev. D55 (1997) 6986; R. Barbieri, L. J. Hall, S. Raby and A. Romanino, Nucl. Phys. B493 (1997) 3-26, hep-ph/9610449; K. S. Babu, J. C. Pati and F. Wilczek, Phys. Lett. B423 (1998) 337-347, hep-ph/9712307; Q. Shafi and Z. Tavartkiladze, Nucl. Phys. B573 (2000) 40-56, hep-ph/9905202; hep-ph/9910314; Phys. Lett. B473 (2000) 272-280, hep-ph/9911264; C. H. Albright and S. M. Barr, hep-ph/0003251; hep-ph/0007145; Z. Berezhiani and A. Rossi, hep-ph/0003084; J. C. Pati, hep-ph/0005095; R. Dermisek, A. Mafi and S. Raby, hep-ph/0007213.
- [94] G. Altarelli, F. Feruglio and I. Masina, JHEP11(2000) 40, hep-ph/0007254.
- [95] G. F. Giudice, E. W. Kolb and A. Riotto, hep-ph/0005123.
- [96] J. Bagger, K. Matchev and D. Pierce, Phys. Lett. B348 (1995) 443-450, hep-ph/9501277.
- [97] L. Clavelli and P. W. Coulter, Phys. Rev. D51 (1995) 3913-3922; hep-ph/9507261.
- [98] T. Junk, Talk at the “Susy 2K” Conference, CERN, June 2000.
- [99] G. F. Giudice and A. Masiero, Phys. Lett. B206 (1988) 480-484.
- [100] P. Langacker and N. Polonsky, Phys. Rev. D47 (1993) 4028-4045, hep-ph/9210235; Phys. Rev. D52 (1995) 3081-3086, hep-ph/9503214.
- [101] P. H. Chankowski, Z. Pluciennik and S. Pokorski, Nucl. Phys. B439 (1995) 23, hep-ph/9411233.
- [102] L. Roszowski and M. Shifman, Phys. Rev. D53 (1996) 404, hep-ph/9503358.
- [103] C. T. Hill, Phys. Lett. B135 (1984) 47; Q. Shafi and C. Wetterich, Phys. Rev. Lett. 52 (1984) 47; J. McDonald and C. E. Vayonakis, Phys. Lett. B144 (1984) 199; J. Ellis et al., Phys. Lett. B155 (1985) 381; M. Drees, Phys. Lett. B158 (1985) 409; L. J. Hall and U. Sarid, Phys. Rev. Lett. 70 (1993) 2673; A. Vayonakis, Phys. Lett. B307 (1993) 318.
- [104] T. Nihei and J. Arafune, Prog. Theor. Phys. 93 (1995) 665-669, hep-ph/9412325.

- [105] M. Claudson, M. B. Wise and L. J. Hall, Nucl. Phys. B195 (1982) 297; S. Chadha and M. Daniels, Nucl. Phys. B229 (1983) 105.
- [106] R. Shrock and L. Wang, Phys. Rev. Lett 41 (1978) 692.

Appendix A

Formulae for the Calculation of Proton Lifetime

A.1 d=5 B Violating Operators

The starting point is the most general renormalizable superpotential which could come from $SU(5)$. Yukawa couplings of doublets and triplets will be considered uncorrelated as well as down quarks and charged lepton mass matrices. The discussion can thus be adapted for any $SU(5)$ based model. Omitting colour and weak isospin indices, in the interaction basis the relevant terms read:

$$\begin{aligned}
 w \ni & Q^T y_u U^c H_{2u} + Q^T y_d D^c H_{2d} + L^T y_e E^c H_{2d} \\
 & + Q^T \hat{A} Q H_{3u} + U^{cT} \hat{B} E^c H_{3u} + Q^T \hat{C} L H_{3d} + U^{cT} \hat{D} D^c H_{3d} \\
 & + m_T H_{3u} H_{3d} \quad ,
 \end{aligned} \tag{A.1}$$

where Q , L , U^c , D^c and E^c denote as usual the chiral multiplets associated to the three fermion generations, $y_{u,d,e}$, $\hat{A}$, $\hat{B}$, ... are matrices in generation space and m_T is the triplets mass. By integrating out the colour triplets one obtains the following effective superpotential

$$w_{eff} = \frac{1}{m_T} \left[Q \hat{A} Q \hat{C} \hat{L} + U^c \hat{B} E^c U^c \hat{D} D^c \right] + \dots \quad , \tag{A.2}$$

dots standing for terms that do not violate baryon or lepton number.

At lower scales these dimension five operators give rise to the four-fermion operators relevant to proton decay, via a "dressing" process. When considering the operator $QQQL$, the most important dressing comes from wino exchange ¹. Charged

¹Gluino dressing contributions cancel among each other in case of degeneracy between first two generations of squarks [83, 84].

$O_1^L = \frac{g^2}{m_T} (\nu_l d_i) (d_j u_k) [L_d^T (\hat{A} + \hat{A}^T) L_d]_{ij} [L_u^T \hat{C} L_e]_{kl} (f_{\tilde{w}}(\tilde{u}, \tilde{e}) + f_{\tilde{w}}(\tilde{u}, \tilde{d}))$
$O_2^L = \frac{g^2}{m_T} (u_i d_j) (d_k \nu_l) [L_u^T (\hat{A} + \hat{A}^T) L_d]_{ij} [L_d^T \hat{C} L_e]_{kl} (f_{\tilde{w}}(\tilde{u}, \tilde{e}) + f_{\tilde{w}}(\tilde{u}, \tilde{d}))$
$O_3^L = -\frac{g^2}{m_T} (u_i d_j) (u_k e_l) [L_u^T (\hat{A} + \hat{A}^T) L_d]_{ij} [L_u^T \hat{C} L_e]_{kl} (f_{\tilde{w}}(\tilde{d}, \tilde{\nu}) + f_{\tilde{w}}(\tilde{u}, \tilde{d}))$
$O_4^L = \frac{g^2}{m_T} (u_j d_k) (u_i e_l) [L_u^T (\hat{A} + \hat{A}^T) L_u]_{ij} [L_d^T \hat{C} L_e]_{kl} (f_{\tilde{w}}(\tilde{d}, \tilde{u}) + f_{\tilde{w}}(\tilde{d}, \tilde{\nu}))$
$O_1^R = -\frac{1}{m_T} (\bar{d}_k \bar{\nu}_l) (u_i^c d_j^c) [L_d^+ y_u^* \hat{B} y_e^+ L_e^*]_{kl} [R_u^* \hat{D} R_d^+]_{ij} f_{\tilde{h}}(\tilde{u}^c, \tilde{e}^c)$
$O_2^R = -\frac{1}{m_T} (\bar{\nu}_l \bar{d}_i) (d_j^c u_k^c) [R_u^* \hat{B} y_e^+ L_e^*]_{kl} [L_d^+ y_u^* \hat{D} R_d^+]_{ij} f_{\tilde{h}}(\tilde{u}^c, \tilde{e}^c)$
$O_3^R = -\frac{1}{m_T} (\bar{d}_i \bar{u}_j) (u_k^c e_l^c) [R_u^* \hat{B} R_e^+]_{kl} [L_d^+ y_u^* \hat{D} y_d^+ L_u^*]_{ij} f_{\tilde{h}}(\tilde{u}^c, \tilde{d}^c)$
$O_4^R = -\frac{1}{m_T} (\bar{u}_j \bar{d}_k) (e_l^c u_i^c) [L_d^+ y_u^* \hat{B} R_e^+]_{kl} [R_u^* \hat{D} y_d^+ L_u^*]_{ij} f_{\tilde{h}}(\tilde{d}^c, \tilde{u}^c)$

Table A.1: Four-fermion operators arising from dimension 5 operators. For each operator it is understood to also add the h.c.

higgsino exchange provides instead the most important dressing of $U^c E^c U^c D^c$ [85].

The integration over the loop has the effect of producing the multiplicative factor $g^2 f_{\tilde{w}}(\tilde{a}, \tilde{b})$ for wino dressing (g being the $SU(2)$ gauge coupling) and $f_{\tilde{h}}(\tilde{a}, \tilde{b})$ for higgsino dressing. f is a function coming from the loop integration:

$$\begin{aligned}
f_{\tilde{e}}(\tilde{a}, \tilde{b}) &= \int \frac{d^4 k}{(2\pi)^4} \frac{-i}{k^2 - m_a^2} \frac{-i}{k^2 - m_b^2} \frac{-i}{\not{k} - m_{\tilde{e}}} \\
&= \frac{1}{16\pi^2} \frac{m_{\tilde{e}}}{(m_a^2 - m_b^2)} \left[\frac{m_a^2}{m_a^2 - m_{\tilde{e}}^2} \log \frac{m_a^2}{m_{\tilde{e}}^2} - \frac{m_b^2}{m_b^2 - m_{\tilde{e}}^2} \log \frac{m_b^2}{m_{\tilde{e}}^2} \right] . \quad (\text{A.3})
\end{aligned}$$

The operators are explicitly given in Table A.1, where we made the not too bad assumption of degeneracy between the generations of each sparticle (the dependence on sparticle masses is logarithmic), so that the final expression is more compact. We also turned to the mass eigenstate basis by indicating with $L_{u,d,e}$ and $R_{u,d,e}$ the unitary matrices that diagonalize the fermion mass matrices:

$$L_u^T y_u R_u^\dagger = y_u^{(d)} \quad - \quad L_d^T y_d R_d^\dagger = y_d^{(d)} \quad - \quad L_e^T y_e R_e^\dagger = y_e^{(d)} \quad , \quad (\text{A.4})$$

with $(y_f)_{(d)}$ ($f = u, d, e$) diagonal and positive. The quark mixing matrix is then $V_{CKM} = L_u^\dagger L_d$.

A.2 The Relevant Operators

As discussed in chapter 3, the most important operators are O_1^L and O_1^R , which are explicitly given by:

$$\begin{aligned} O_1^L &= C_{ijkl}^L (\nu_l d_i) (d_j u_k) + \text{h.c.} \\ O_1^R &= C_{ijkl}^R (d_k \nu_l) (\bar{u}_i^c \bar{d}_j^c) + \text{h.c.} \quad , \end{aligned} \quad (\text{A.5})$$

where i, j, k, l are generation indices and

$$\begin{aligned} C_{ijkl}^L &= \frac{K_L}{m_T} g^2 [f_{\tilde{w}}(\tilde{u}, \tilde{d}) + f_{\tilde{w}}(\tilde{u}, \tilde{e})] P_{ij} Q_{kl} \\ C_{ijkl}^R &= -\frac{K_R}{m_T} f_{\tilde{h}}(\tilde{u}^c, \tilde{e}^c) S_{kl}^* T_{ij}^* \end{aligned} \quad (\text{A.6})$$

$$P = 2 L_d^T \hat{A} L_d \quad Q = L_u^T \hat{C} L_e \quad (\text{A.7})$$

$$S = L_d^\dagger y_u^* \hat{B} y_e^\dagger L_e^* \quad T = R_u^* \hat{D} R_d^\dagger \quad .$$

Considering O_1^L , for the channel $p \rightarrow K^+ \bar{\nu}$ we have to analyze the contributions from both $(\nu_l d)$ (su), which corresponds to C_{121l}^L , and $(\nu_l s)$ (du), which corresponds to C_{211l}^L . Since C_{ijkl}^L is proportional to $P_{ij} Q_{kl}$, it is possible to identify immediately the important P, Q elements. For instance C_{121l}^L is linked to P_{12} and Q_{1l} . Note that P is symmetric, hence $C_{121l}^L = C_{211l}^L$. For minimal $SU(5)$, P, Q greatly simplify, as can be seen from Table 3.2. In this case

$$P_{12} = [V^T y_u^{(d)} K^+ V]_{12} \sim y_{top} \lambda^5 \quad , \quad (\text{A.8})$$

and $Q_{1l} = [V^* y_d^{(d)}]_{1l}$, so that

$$Q_{12} \sim Q_{13} \sim y_{bot} \lambda^3 \gg Q_{11} \quad . \quad (\text{A.9})$$

Thus, since

$$y_{top}y_{bot} \sim \lambda^2(\tan \beta + 1/\tan \beta) \quad , \quad (\text{A.10})$$

one obtains

$$P_{12}Q_{12(3)} \sim \lambda^{10}(\tan \beta + 1/\tan \beta) \quad , \quad (\text{A.11})$$

as shown in Table 3.3. For the channel $p \rightarrow \pi^+ \bar{\nu}$, we have only to consider $(\nu_l d)(du)$, that is $P_{11}Q_{1l}$. Since $P_{11} \sim \lambda P_{12}$, then

$$P_{11}Q_{12(3)} \sim \lambda^{11}(\tan \beta + 1/\tan \beta) \quad . \quad (\text{A.12})$$

In the case of O_1^R , the channel with K^+ is obtained for $(d\nu_l) (\overline{u^c s^c})$, corresponding to $S_{1l}T_{12}$, and $(s\nu_j) (\overline{u^c d^c})$, corresponding to $S_{2l}T_{11}$. In the minimal case, from

$$S = V^+ y_u^{(d)2} V y_d^{(d)} \quad , \quad (\text{A.13})$$

it turns out that the dominant contribution corresponds to $l = 3$, that is emission of $\bar{\nu}_\tau$, and $S_{13} \sim y_{top}^2 y_{bot} \lambda^3$, $S_{23} \sim y_{top}^2 y_{bot} \lambda^2$. From

$$T = K V^* y_d^{(d)} \quad (\text{A.14})$$

it follows $T_{12} \sim y_{bot} \lambda^3$, $T_{11} \sim y_{bot} \lambda^4$. Thus,

$$S_{13}T_{12} \sim y_{top}^2 y_{bot}^2 \lambda^6 \sim \lambda^{10}(\tan \beta + 1/\tan \beta)^2 \sim S_{23}T_{11} \quad . \quad (\text{A.15})$$

For the channel with π^+ , we have to consider $(d\nu_l) (\overline{u^c d^c})$, that is

$$S_{13}T_{11} \sim \lambda^{11}(\tan \beta + 1/\tan \beta)^2 \quad . \quad (\text{A.16})$$

These results are given in Table 3.3.

Note that we have also introduced $K_{L,R}$, which are constants accounting for the renormalization of the operators from the grand unification scale down to 1 GeV [78, 81, 104]. In our estimates we take $K_1 = K_2 = 10$.

The next step is to parameterize the partial rates according to the results of a chiral Lagrangian computation [105]. The rates for the dominant channels are given in Table A.2.

channel	rate
$K^+ \bar{\nu}_\mu$	$X_K \left \beta \left[\frac{2}{3} \frac{m_p}{m_B} D C_{1212}^L + \left(1 + \frac{m_p}{3m_B} (D + 3F) \right) C_{2112}^L \right] \right ^2$
$K^+ \bar{\nu}_\tau$	$X_K \left \beta \left[\frac{2}{3} \frac{m_p}{m_B} D C_{1213}^L + \left(1 + \frac{m_p}{3m_B} (D + 3F) \right) C_{2113}^L \right] \right.$ $\left. + \alpha \left[\frac{2}{3} \frac{m_p}{m_B} D C_{1312}^R + \left(1 + \frac{m_p}{3m_B} (D + 3F) \right) C_{2311}^R \right] \right ^2$
$\pi^+ \bar{\nu}_\mu$	$X_\pi \left \beta (1 + D + F) 2C_{1112}^L \right ^2$
$\pi^+ \bar{\nu}_\tau$	$X_\pi \left \beta (1 + D + F) 2C_{1113}^L + \alpha (1 + D + F) 2C_{1311}^R \right ^2$

Table A.2: Proton decay rates. We define $X_{K,\pi} = (m_p^2 - m_{K,\pi}^2)^2 / (32\pi m_p^3 f_\pi^2)$; m_p , m_K , m_π are the proton, K^+ and π^+ masses; m_B is an average baryon mass, f_π is the pion decay constant; D and F are coupling constants between baryons and mesons in the relevant chiral Lagrangian; β and α parameterize the hadronic matrix element. In our estimates we take $m_p = 0.938$ GeV, $m_B = 1.150$ GeV, $m_K = 0.494$ GeV, $m_\pi = 0.140$ GeV, $f_\pi = 0.139$ GeV, $D = 0.8$, $F = 0.45$ [106], $\beta = -\alpha = 0.003$ GeV³ [88].

Appendix B

Representative SUSY Spectrum

$SU(2) \otimes U(1)$ breaking effects are neglected. Numerical coefficients appearing in the previous Table are due to renormalization.

The additional freedom related to the parameters m_0 , $m_{1/2}$, μ , m_H is here fixed by choosing $0.8m_0 = 0.8m_{1/2} = 2\mu = m_H$ and taking as a definition $m_{SUSY} \equiv m_H$, so that all particle masses can be expressed in term of m_{SUSY} . This parameterization leads to $k(SUSY) = -0.510$.

sparticle	mass ²
gluinos	$(2.7m_{1/2})^2$
winos	$(0.8m_{1/2})^2$
higgsinos	μ^2
extra Higgses	m_H^2
squarks	$m_0^2 + 6m_{1/2}^2$
(sleptons) _L	$m_0^2 + 0.5m_{1/2}^2$
(sleptons) _R	$m_0^2 + 0.15m_{1/2}^2$

Table B.1: Representative SUSY spectrum.

Appendix C

Group Theory for the MDM

In this appendix we summarize the method adopted to deal with tensors in $SU(5)$. It is convenient to write large representations like 75, 50 and $\bar{50}$ as products of smaller representations, like 10 and $\bar{10}$. In this way, it is not difficult to recognize how the multiplets of the SM are embedded in those large representations, thus staying in touch with the physical role of the various superpotential terms. In particular, when considering the interaction terms, it is compelling to accurately take into account the numerical coefficients which follow directly from the tensorial structure of the representations involved. Since these coefficients can be large, it is not possible to omit them when performing a quantitative analysis.

C.1 Basic Conventions for the Tensors in $SU(5)$

In $SU(5)$ the matter fields are contained in just the 10 and $\bar{5}$ representations. We write them according to

$$\Psi_{10} = \frac{1}{\sqrt{2}} \begin{bmatrix} 0 & U_3^c & -U_2^c & U^1 & D^1 \\ -U_3^c & 0 & U_1^c & U^2 & D^2 \\ U_2^c & -U_1^c & 0 & U^3 & D^3 \\ -U^1 & -U^2 & -U^3 & 0 & E^c \\ -D^1 & -D^2 & -D^3 & -E^c & 0 \end{bmatrix}, \quad (\text{C.1})$$

$$\Psi_{\bar{5}} = \begin{bmatrix} D_1^c \\ D_2^c \\ D_3^c \\ E \\ -\nu \end{bmatrix}. \quad (\text{C.2})$$

The 10 representation Ψ_{10}^{AB} is thus represented by a tensor whose two indices $A, B = 1, \dots, 5$ are antisymmetric. Note that the $SU(3)$ indices correspond to $A, B = 1, 2, 3$ while the $SU(2)$ ones are selected for $A, B = 4, 5$. To remember about this in an explicit way, we will sometimes write α (a) instead of A if $A = 1, 2, 3$ ($4, 5$). Due to the antisymmetry of Ψ_{10}^{AB} , it turns out that it is convenient to deal just with $\Psi_{10}^{AB} |_{A < B}$. The $\bar{5}$ representation is instead written as a tensor with one low index Ψ_{5A} .

In our conventions, the Higgs fields $\bar{H}_A, H^A$ belonging respectively to the $\bar{5}$ and 5 representations are written as

$$\bar{H} = \begin{bmatrix} H_{3d1} \\ H_{3d2} \\ H_{3d3} \\ H_{2d}^- \\ -H_{2d}^0 \end{bmatrix} \quad H = \begin{bmatrix} H_{3u}^1 \\ H_{3u}^2 \\ H_{3u}^3 \\ H_{2u}^+ \\ H_{2u}^0 \end{bmatrix} , \quad (C.3)$$

where $\langle H_{2d}^0 \rangle = v_d/\sqrt{2}$, $\langle H_{2u}^0 \rangle = v_u/\sqrt{2}$.

Let us underline a few points.

- The factor $1/\sqrt{2}$ is introduced in the definition of Ψ_{10} in order to have the kinetic terms in the canonical form. In fact, the Kahler potential contains

$$K \ni \Psi_{10}^{AB} \Psi_{10}^{AB*} , \quad (C.4)$$

where summation over repeated indices has to be understood. Due to the antisymmetry of Ψ_{10} under $A \leftrightarrow B$, one immediately realizes that:

$$\begin{aligned} \Psi_{10}^{AB} \Psi_{10}^{AB*} &= 2 \Psi_{10}^{AB} \Psi_{10}^{AB*} |_{A < B} = \\ &= U_\alpha^c U_\alpha^{c*} + U^\alpha U^{\alpha*} + D^\alpha D^{\alpha*} + E^c E^{c*} , \end{aligned} \quad (C.5)$$

where α is the colour index. Thus, the fields U^c, U, D, E^c have a canonical kinetic term. On the other hand the pentaplet Ψ_{5A} has no problems of multiplicity, so that there is no need of introducing any coefficient in the definition of (C.2) to get the canonical form for the kinetic terms of D^c, E, ν .

- At the contrary of what happens for all the other $\Psi_{10}^{AB} |_{A < B}$, we have introduced a minus sign in Ψ_{10}^{13} . The reason is the following. The mass term for up quarks (considering for simplicity one generation only) arises from the superpotential term

$$\begin{aligned} w &\ni \frac{1}{4} \Psi_{10}^{AB} \Psi_{10}^{CD} H^E \epsilon_{ABCDE} = \\ &= \Psi_{10}^{AB} \Psi_{10}^{CD} H^E \epsilon_{ABCDE} |_{A < B, C < D} \end{aligned} \quad (C.6)$$

with $E = 5$. When one picks the indices corresponding to $U_2^c U^2$, that is $(A, B, C, D) = (1, 3, 2, 4)$, a minus sign arises from ϵ_{13245} , at the contrary of what happens for $U_1^c U^1$ and $U_3^c U^3$. The minus sign in Ψ_{10}^{13} is put to eliminate the minus arising from the ϵ . Clearly there is nothing restrictive in doing this, since we are only exploiting the freedom of choosing a convenient phase in the definition of our fields.

C.2 The 75 and 50 representations

How to write the 75

The 75 can be conveniently written through the product

$$10 \times \overline{10} = 1 + 24 + 75 \quad . \quad (\text{C.7})$$

Let us clarify how to isolate the 75 from the 24 and the 1. We define $10^{AB} \times \overline{10}_{CD} \equiv T_{CD}^{AB}$, so that T_{CD}^{AB} is antisymmetric in the upper and lower pair of indices and thus corresponds to 100 independent fields. It can be split according to

$$T_{CD}^{AB} = \underbrace{S_{CD}^{AB}}_1 + \underbrace{\Sigma_{CD}^{AB}}_{24} + \underbrace{Y_{CD}^{AB}}_{75} \quad (\text{C.8})$$

where we identify the 1 with S , the 24 with Σ and the 75 with Y .

Consider the $SU(5)$ singlet first. One can easily guess it to be given by the combination ¹

$$s \equiv T_{MN}^{MN} \quad . \quad (\text{C.9})$$

Thus, S has to be

$$S_{CD}^{AB} = \frac{1}{20}(\delta_C^A \delta_D^B - \delta_D^A \delta_C^B) T_{MN}^{MN} \quad . \quad (\text{C.10})$$

In this case, in fact, substituting S_{CD}^{AB} into eq. (C.8) it turns out that $S_{AB}^{AB} = T_{AB}^{AB} \equiv s$ and that Σ and Y must be such that $\Sigma_{AB}^{AB} = Y_{AB}^{AB} = 0$. This indeed means that the singlet has been completely projected out by S .

The 24 can be written as a tensor σ_C^A satisfying the constraint $\sigma_A^A = 0$. One can guess

$$\sigma_C^A \equiv T_{MC}^{MA} - \frac{1}{5} \delta_C^A T_{MN}^{MN} \quad (\text{C.11})$$

¹Here and in the following summation over repeated indices has to be understood.

and

$$\begin{aligned} \Sigma_{CD}^{AB} = & \frac{1}{3} \left[\delta_C^A (T_{MD}^{MB} - \frac{1}{5} \delta_D^B T_{MN}^{MN}) - \delta_D^A (T_{MC}^{MB} - \frac{1}{5} \delta_C^B T_{MN}^{MN}) \right. \\ & \left. - \delta_C^B (T_{MD}^{MA} - \frac{1}{5} \delta_D^A T_{MN}^{MN}) + \delta_D^B (T_{MC}^{MA} - \frac{1}{5} \delta_C^A T_{MN}^{MN}) \right] . \end{aligned} \quad (C.12)$$

Notice that this form for Σ satisfies $\Sigma_{AB}^{AB} = 0$, that is Σ doesn't contain the combination corresponding to the singlet s . Substituting this expression in eq. (C.8), we get $\Sigma_{MC}^{MA} = \sigma_C^A$, $S_{MC}^{MA} = 0$ and the requirement $Y_{MC}^{MA} = 0$. This means that the 24 is completely projected out by Σ .

Finally, with the expressions of S_{CD}^{AB} and Σ_{CD}^{AB} at our disposal, we know how to write the 75:

$$\begin{aligned} Y_{CD}^{AB} = & T_{CD}^{AB} \\ & - \frac{1}{3} \left[\delta_C^A (T_{MD}^{MB} - \frac{1}{5} \delta_D^B T_{MN}^{MN}) - \delta_D^A (T_{MC}^{MB} - \frac{1}{5} \delta_C^B T_{MN}^{MN}) \right. \\ & \left. - \delta_C^B (T_{MD}^{MA} - \frac{1}{5} \delta_D^A T_{MN}^{MN}) + \delta_D^B (T_{MC}^{MA} - \frac{1}{5} \delta_C^A T_{MN}^{MN}) \right] \\ & - \frac{1}{20} (\delta_C^A \delta_D^B - \delta_D^A \delta_C^B) T_{MN}^{MN} . \end{aligned} \quad (C.13)$$

Note that Y is antisymmetric in both the upper and lower indices. This would give 100 independent fields but, since it satisfies the 25 constraints $Y_{MC}^{MA} = 0$, we are correctly left with only 75 independent fields.

It is important to realize that the fields in Y_{CD}^{AB} don't have canonical kinetic terms. This can be recognized by the following reasoning. Let us suppose to put the 100 elements of $Y_{CD}^{AB}|_{A<B, C<D}$ into a column vector, say $\mathcal{Y}_{\mathcal{A}}$, with $\mathcal{A} = 1, 2, \dots, 100$. Clearly only 75 of these 100 elements correspond to independent fields. The Kahler potential is then

$$\begin{aligned} K & \ni Y_{CD}^{AB*} Y_{CD}^{AB} = 4 Y_{CD}^{AB*} Y_{CD}^{AB} |_{A<B, C<D} = \\ & = 4 \mathcal{Y}^+ \mathcal{Y} . \end{aligned} \quad (C.14)$$

We can always exploit the possibility of rotating the fields by means of a 100 by 100 unitary matrix U . For instance, we can get rid of the unpleasant redundancy by going into a basis where $\mathcal{Y}$ has only 75 non-zero elements. Then, it is also feasible to find a unitary transformation of the 75 fields which carry us into a SM-like basis, that is a basis where the fields are split into multiplets of the SM. In a similar basis it is evident that the kinetic terms are not canonical because of the factor 4 appearing

in eq. (C.14). The remedy is to define $Y_{CD}^{AB} \equiv Y'_{CD}^{AB}/2$ (and $\mathcal{Y} \equiv \mathcal{Y}'/2$), so that the fields associated to Y'_{CD}^{AB} are those actually possessing the correct coefficient 1 in the kinetic term.

How to write the $50, \bar{50}$

By similar considerations, the 50 can be written from the product of 10×10 . In fact

$$10 \times 10 = \bar{5}_s + \bar{45}_a + 50_s \quad . \quad (\text{C.15})$$

We denote the 10×10 with $T_1^{AB}T_2^{CD}$, where $T_{1,2}$ are antisymmetric under exchange of the two upper indices. Again we want to separate $T_1^{AB}T_2^{CD}$ as

$$T_1^{AB}T_2^{CD} = \underbrace{C^{ABCD}}_{\bar{5}_s} + \underbrace{Q^{ABCD}}_{\bar{45}_a} + \underbrace{F^{ABCD}}_{50_s} \quad . \quad (\text{C.16})$$

It is useful to split $T_1^{AB}T_2^{CD}$ into its symmetric and antisymmetric parts under the exchange $1 \leftrightarrow 2$, so that

$$\begin{aligned} T_1^{AB}T_2^{CD} &= \underbrace{\frac{T_1^{AB}T_2^{CD} + T_2^{AB}T_1^{CD}}{2}}_{\bar{5}_s, 50_s} + \underbrace{\frac{T_1^{AB}T_2^{CD} - T_2^{AB}T_1^{CD}}{2}}_{\bar{45}_a} \\ &\equiv S^{ABCD} + A^{ABCD} \quad . \end{aligned} \quad (\text{C.17})$$

It follows that Q is antisymmetric while C, F are symmetric under the exchange $(AB) \leftrightarrow (CD)$ (or equivalently $1 \leftrightarrow 2$). Thus, clearly the $\bar{45}_a$, q^{ABCD} , is given by

$$q^{ABCD} = A^{ABCD} \quad , \quad (\text{C.18})$$

and the simple identification $Q^{ABCD} = A^{ABCD}$ holds because $(Q^{ABCD} - Q^{CDAB})/2 = q^{ABCD}$ while $(C^{ABCD} - C^{CDAB})/2 = (F^{ABCD} - F^{CDAB})/2 = 0$. Then, we have only to separate the $\bar{5}_s$ from the 50_s . One can guess that the $\bar{5}_s$ is given by

$$c_E = \epsilon_{ABCDE} S^{ABCD} \quad (\text{C.19})$$

and verify that the following expression for C^{ABCD}

$$C^{ABCD} = \frac{1}{24} \epsilon^{ABCDE} \epsilon_{FGHIE} S^{FGHI} \quad (\text{C.20})$$

satisfies $\epsilon_{ABCDE} C^{ABCD} = c_E$. Since $\epsilon_{ABCDE} Q^{ABCD} = 0$, C completely projects out the $\bar{5}$ if also $\epsilon_{ABCDE} F^{ABCD} = 0$ holds. This property is automatically satisfied by

$$\begin{aligned} F^{ABCD} &= T_1^{AB}T_2^{CD} \\ &- A^{ABCD} \\ &- \frac{1}{24} \epsilon^{ABCDE} \epsilon_{FGHIE} S^{FGHI} \quad . \end{aligned} \quad (\text{C.21})$$

Note that F is antisymmetric in $[AB]$ and $[CD]$ but is symmetric for the exchange $(AB) \leftrightarrow (CD)$.

By writing the Kahler potential as $K \ni F^{ABCD*} F^{ABCD}$, it turns out that F^{ABCD} hasn't canonical kinetic terms. The proof is very similar to the one already discussed for the case of the 75. Also in the present case, defining $F^{ABCD} \equiv F'^{ABCD}/2$, one realizes that the fields F'^{ABCD} are those with the kinetic terms in the canonical form.

The $\overline{50}$ is obtained following the same procedure but starting from the product of $\overline{10} \times \overline{10} = \overline{T}_{1AB} \overline{T}_{2CD}$.

C.3 The Method for Catching SM Multiplets

The construction of the 75, 50 and $\overline{50}$ in terms of products of 10 and/or $\overline{10}$ is particularly useful because one can apply a simple recipe to recognize how the SM multiplets are included in these large representations. The starting point is the identification of the SM multiplets in the 10 and $\overline{10}$ themselves. Keeping $A < B$:

$$\begin{aligned} T^{AB} &= \underbrace{10^{\alpha\beta}}_{(\overline{3},1,-2/3)} + \underbrace{10^{\alpha b}}_{(3,2,1/6)} + \underbrace{10^{ab}}_{(1,1,1)} \\ \overline{T}_{AB} &= \underbrace{10_{\alpha\beta}}_{(3,1,2/3)} + \underbrace{10_{ab}}_{(\overline{3},2,-1/6)} + \underbrace{10_{ab}}_{(1,1,-1)} . \end{aligned} \quad (C.22)$$

Let us consider the case of the 75 (50) in order to be specific. One starts by writing the SM multiplets obtained by multiplying each SM multiplet of T with each SM multiplet of $\overline{T}$ (T). All the 100 fields obtained are thus separated in SM multiplets. Next we have to further correctly separate the various multiplets into the 1, 24 and 75 ($\overline{5}_s$, $\overline{45}_a$ and 50_s). This is easy to do for such multiplets which are obtained only once, because from Table C.1 we immediately understand in which $SU(5)$ representation they lie. But for the multiplets which are obtained more than once, like e.g. the singlet (the color antitriplet), we have to understand which linear combination of them is the one actually embedded in the 75 (50). As an example, we show the procedure which permits to single out the SM singlet contained in the 75 and the SM antitriplet contained in the 50.

The SM singlet of the 75

There are three singlets $(1, 1, 0)$ arising from the product $T \times \overline{T}$.

$$SU(5) \supset SU(3) \times SU(2) \times U(1)$$

$$1 = (1, 1, 0)$$

$$24 = (1, 1, 0) + (1, 3, 0) + (3, 2, -5/6) + (\bar{3}, 2, 5/6) + (8, 1, 0)$$

$$50 = (1, 1, 2) + (\bar{3}, 1, 1/3) + (3, 2, 7/6) + (6, 3, 1/3) + (\bar{6}, 1, -4/3) + \\ +(8, 2, -1/2)$$

$$\bar{50} = (1, 1, -2) + (3, 1, -1/3) + (\bar{3}, 2, -7/6) + (\bar{6}, 3, -1/3) + (6, 1, 4/3) + \\ +(8, 2, 1/2)$$

$$75 = (1, 1, 0) + (3, 1, 5/3) + (\bar{3}, 1, -5/3) + (3, 2, -5/6) + (\bar{3}, 2, 5/6) + \\ +(\bar{6}, 2, -5/6) + (6, 2, 5/6) + (8, 1, 0) + (8, 3, 0)$$

Table C.1: The SM multiplets contained in some of the $SU(5)$ representations.

We call s_1 the singlet which is obtained by multiplying

$$T_{\gamma\delta}^{\alpha\beta} \equiv \underbrace{T^{\alpha\beta}}_{(\bar{3},1,-2/3)} \underbrace{\bar{T}_{\gamma\delta}}_{(3,1,2/3)} = \underbrace{\frac{1}{6}(\delta_\gamma^\alpha \delta_\delta^\beta - \delta_\delta^\alpha \delta_\gamma^\beta) T_{\mu\nu}^{\mu\nu}}_{(1,1,0)} + \underbrace{T_{\gamma\delta}^{\alpha\beta} - \frac{1}{6}(\delta_\gamma^\alpha \delta_\delta^\beta - \delta_\delta^\alpha \delta_\gamma^\beta) T_{\mu\nu}^{\mu\nu}}_{(8,1,0)} . \quad (\text{C.23})$$

It is then evident that $s_1 \propto T_{\mu\nu}^{\mu\nu} = 2T_{\mu\nu}^{\mu\nu}|_{\mu<\nu}$. Thus, choosing to normalize to 1 each combination of independent fields:

$$s_1 = \frac{1}{\sqrt{3}}(T_{12}^{12} + T_{13}^{13} + T_{23}^{23}) = \frac{1}{\sqrt{3}}T_{\mu\nu}^{\mu\nu}|_{\mu<\nu} . \quad (\text{C.24})$$

The second singlet s_2 comes from the product

$$\begin{aligned} T_{\gamma d}^{\alpha b} \equiv \underbrace{T^{\alpha b}}_{(3,2,1/6)} \underbrace{\bar{T}_{\gamma d}}_{(\bar{3},2,-1/6)} &= \underbrace{\frac{1}{6}\delta_\gamma^\alpha \delta_d^b T_{\mu n}^{\mu n}}_{(1,1,0)} \\ &+ \underbrace{\frac{1}{3}\delta_\gamma^\alpha T_{\mu d}^{\mu b} - \frac{1}{6}\delta_\gamma^\alpha \delta_d^b T_{\mu n}^{\mu n}}_{(1,3,0)} \\ &+ \underbrace{\frac{1}{2}\delta_d^b T_{\gamma n}^{\alpha n} - \frac{1}{6}\delta_\gamma^\alpha \delta_d^b T_{\mu n}^{\mu n}}_{(8,1,0)} \\ &+ \underbrace{T_{\gamma d}^{\alpha b} - \frac{1}{3}\delta_\gamma^\alpha T_{\mu d}^{\mu b} - \frac{1}{2}\delta_d^b T_{\gamma n}^{\alpha n} + \frac{1}{6}\delta_\gamma^\alpha \delta_d^b T_{\mu n}^{\mu n}}_{(8,3,0)} . \end{aligned} \quad (\text{C.25})$$

Thus, $s_2 \propto T_{\mu n}^{\mu n}$. Precisely:

$$s_2 = \frac{1}{\sqrt{6}}(T_{14}^{14} + T_{24}^{24} + T_{34}^{34} + T_{15}^{15} + T_{25}^{25} + T_{35}^{35}) = \frac{1}{\sqrt{6}}T_{\mu n}^{\mu n} . \quad (\text{C.26})$$

Finally the last singlet s_3 is obtained from

$$T_{cd}^{ab} \equiv \underbrace{T^{ab}}_{(1,1,1)} \underbrace{\bar{T}_{cd}}_{(1,1,-1)} = \underbrace{\frac{1}{2}(\delta_c^a \delta_d^b - \delta_d^a \delta_c^b) T_{mn}^{mn}}_{(1,1,0)} \quad (\text{C.27})$$

and is simply

$$s_3 = T_{45}^{45} = T_{mn}^{mn}|_{m<n} . \quad (\text{C.28})$$

The next step is to understand which linear combination of $s_{1,2,3}$ is contained in the 75. This can be accomplished by using eq. (C.8) and the expressions for S, Σ, Y , that is eqs. (C.10, C.12, C.13). Indeed, by constructing on the l.h.s. of eq. (C.8) the combination relative to $s_{1,2,3}$, we obtain respectively:

$$\begin{aligned} s_1 &\equiv \frac{T_{\alpha\beta}^{\alpha\beta}}{2\sqrt{3}} = \left[\sqrt{\frac{3}{10}} S_1 \right]_S + \left[\sqrt{\frac{8}{15}} S_{24} \right]_\Sigma + \left[\sqrt{\frac{1}{6}} S_{75} \right]_Y \\ s_2 &\equiv \frac{T_{ab}^{ab}}{\sqrt{6}} = \left[\sqrt{\frac{6}{10}} S_1 \right]_S + \left[-\sqrt{\frac{1}{15}} S_{24} \right]_\Sigma + \left[-\sqrt{\frac{2}{6}} S_{75} \right]_Y \\ s_3 &\equiv \frac{T_{ab}^{ab}}{2} = \left[\sqrt{\frac{1}{10}} S_1 \right]_S + \left[-\sqrt{\frac{6}{15}} S_{24} \right]_\Sigma + \left[\sqrt{\frac{3}{6}} S_{75} \right]_Y \end{aligned} \quad (C.29)$$

where

$$\begin{aligned} S_1 &\equiv \frac{\sqrt{3}s_1 + \sqrt{6}s_2 + s_3}{\sqrt{10}} \\ S_{24} &\equiv \frac{\sqrt{8}s_1 - s_2 - \sqrt{6}s_3}{\sqrt{15}} \\ S_{75} &\equiv \frac{s_1 - \sqrt{2}s_2 + \sqrt{3}s_3}{\sqrt{6}} . \end{aligned} \quad (C.30)$$

It is clear that by taking in the l.h.s. of eq. (C.8) the combination S_{75} , one obtains $S_{75} = [0]_S + [0]_\Sigma + [S_{75}]_Y$. The SM siglet lying in the 75 is then S_{75} . One can check that the combinations S_1, S_{24} are instead respectively the singlets of the 1 and 24 representations. Note that S_{75} can be written as

$$S_{75} = \frac{1}{3\sqrt{2}} \left[T_{\alpha\beta}^{\alpha\beta} + 2T_{ab}^{ab} - \frac{1}{2}T_{AB}^{AB} \right] . \quad (C.31)$$

The antitriplet of the 50

There are three multiplets in the product 10×10 having the quantum numbers of the antitriplets, that is $(\bar{3}, 1, 1/3)$. Two arise from the products

$$\underbrace{T_1^{ab}}_{(1,1,1)} \underbrace{T_2^{\gamma\delta}}_{(\bar{3},1,-2/3)} \equiv t_1^{ab\gamma\delta} \quad (C.32)$$

and

$$\underbrace{T_1^{\alpha\beta}}_{(\bar{3},1,-2/3)} \underbrace{T_2^{cd}}_{(1,1,1)} \equiv t_2^{\alpha\beta cd} . \quad (C.33)$$

The other one is contained in

$$\underbrace{T_1^{\alpha b}}_{(3,2,1/6)} \underbrace{T_2^{\gamma d}}_{(3,2,1/6)} \quad (C.34)$$

and can be written as

$$\frac{1}{2}(T_1^{\alpha b}T_2^{\gamma d} + T_1^{\gamma d}T_2^{\alpha b} - T_1^{\alpha d}T_2^{\gamma b} - T_1^{\gamma b}T_2^{\alpha d}) \equiv t_3^{\alpha b \gamma d} \quad (C.35)$$

Following the same procedure already described in some detail for the extraction of the SM singlet of the 75, we take on the l.h.s. of eq. (C.16) the combination corresponding to each antitriplet:

$$\begin{aligned} t_1^{ab\gamma\delta} &= \left[\sqrt{\frac{1}{6}} T_{(5)}^{ab\gamma\delta} \right]_C + \left[\sqrt{\frac{1}{2}} T_{(45)}^{ab\gamma\delta} \right]_Q + \left[\sqrt{\frac{1}{3}} T_{(50)}^{ab\gamma\delta} \right]_F \\ t_2^{\gamma\delta ab} &= \left[\sqrt{\frac{1}{6}} T_{(5)}^{ab\gamma\delta} \right]_C + \left[-\sqrt{\frac{1}{2}} T_{(45)}^{ab\gamma\delta} \right]_Q + \left[\sqrt{\frac{1}{3}} T_{(50)}^{ab\gamma\delta} \right]_F \\ t_3^{\gamma a \delta b} &= \left[-\sqrt{\frac{2}{3}} T_{(5)}^{ab\gamma\delta} \right]_C + \left[\sqrt{\frac{1}{3}} T_{(50)}^{ab\gamma\delta} \right]_F \end{aligned} \quad (C.36)$$

where

$$\begin{aligned} T_{(5)}^{ab\gamma\delta} &= \frac{t_1^{ab\gamma\delta} + t_2^{\gamma\delta ab} - 2 t_3^{\gamma a \delta b}}{\sqrt{6}} \\ T_{(45)}^{ab\gamma\delta} &= \frac{t_1^{ab\gamma\delta} - t_2^{\gamma\delta ab}}{\sqrt{2}} \\ T_{(50)}^{ab\gamma\delta} &= \frac{t_1^{ab\gamma\delta} + t_2^{\gamma\delta ab} + t_3^{\gamma a \delta b}}{\sqrt{3}} \end{aligned} \quad (C.37)$$

Thus, one realizes that the linear combination of the above antitriplets which is contained in the 50 is $T_{(50)}^{ab\gamma\delta}$. Note that, as required, it is symmetric under the exchange $T_1 \leftrightarrow T_2$.

To get rid of the indices it is useful to define $t_{(50)\alpha} = 1/4 \epsilon_{\alpha\gamma\delta} \epsilon_{ab} T_{(50)}^{ab\gamma\delta}$. Note that, as already discussed, $t_{(50)\alpha}$ hasn't canonical kinetic terms. It is $H'_{3d\alpha}$ (introduced in section 4.5), defined as $t_{(50)\alpha} \equiv H'_{3d\alpha}/2$, that possesses the correct coefficient 1: $K \ni F^{ABCD*} F^{ABCD} \ni 4 t_{(50)\alpha}^* t_{(50)\alpha} = H'_{3d\alpha}^* H'_{3d\alpha}$.

C.4 Basic Formulae for the MDM

$SU(5)$ breaking

The part of the superpotential responsible for the breaking of $SU(5)$ reads

$$w_{Higgs} = c_1 Y_{CD}^{AB} Y_{EF}^{CD} Y_{AB}^{EF} + M_Y Y_{CD}^{AB} Y_{AB}^{CD} \quad , \quad (C.38)$$

where, as usual, $A, B, \dots = 1, 2, \dots, 5$. This is actually the most general renormalizable superpotential that can be written by means of a $\mathbf{75}$, the cubic term picking the combination $(\mathbf{75} \times \mathbf{75})_{\mathbf{75}} \times \mathbf{75}$. The other cubic term that one could write, namely $Y_{ED}^{AC} Y_{FC}^{BD} Y_{AB}^{EF}$, turns out to be proportional to $Y_{CD}^{AB} Y_{EF}^{CD} Y_{AB}^{EF}$.

The SUSY vacuum of the $\mathbf{75}$ breaks $SU(5)$ uniquely to $SU(3) \otimes SU(2) \otimes U(1)$:

$$\langle Y_{CD}^{AB} \rangle = -\langle Y \rangle \left[\left(\delta_\gamma^\alpha \delta_\delta^\beta + 2\delta_c^a \delta_d^b - \frac{1}{2} \delta_C^A \delta_D^B \right) - (A \leftrightarrow B) \right] \quad , \quad (C.39)$$

where, as usual, Greek indices assume values from 1 to 3, Latin indices from 4 to 5 and

$$\langle Y \rangle \equiv M_Y / (2c_1) \quad . \quad (C.40)$$

The above superpotential gives a mass to all physical components of Y , i.e. those that are not absorbed by the Higgs mechanism. The calculation is not difficult and leads to Table 4.2.

Gauge vector bosons instead acquire a mass m_V :

$$m_V^2 = 24g_5^2 \langle Y \rangle^2 \quad . \quad (C.41)$$

The triplet mass matrix

The doublet-triplet splitting is realized by the following superpotential terms,

$$w_{MDM} = c_2 H^A Y_{BC}^{DE} H_{50}^{BCFG} \epsilon_{ADEF G} + c_3 \bar{H}_A Y_{DE}^{BC} H_{\bar{50}BCFG} \epsilon^{ADEF G} + M_{50} H_{50}^{ABCD} H_{\bar{50}ABCD} \quad . \quad (C.42)$$

The multiplet in Y which gets a vev is the SM singlet. This means that in the term

$$c_2 H^A Y_{BC}^{DE} H_{50}^{BCFG} \epsilon_{ADEF G} \quad (C.43)$$

only the interaction between the triplet contained in H^A and the antitriplet contained in H_{50} is allowed. Since H_{50}^{ABCD} can be identified with F^{ABCD} , using eqs. (C.39) and (C.36), one obtains

$$c_2 H^A \langle Y_{BC}^{DE} \rangle H_{50}^{BCFG} \epsilon_{ADEF G} = -4\sqrt{3} c_2 \langle Y \rangle H'_{3d} H_{3u} \quad . \quad (C.44)$$

Finally,

$$H_{50}{}^{ABCD} H_{\overline{50}ABCD} \ni H'_{3d} H'_{3u} \quad (\text{C.45})$$

so that the coefficients appearing in eq. (3.9) are obtained.

Interaction of $10 \ 10 \ \overline{50}$

We are interested in the interactions of the colour triplet of the $\overline{50}$ with the antitriplet combination of $10_i \times 10_j$, 10 being matter fields and i, j family indices. It is convenient to recognize that

$$I \equiv \Psi_{10 \ i}^{AB} G_{\overline{50}ij} \Psi_{10 \ j}^{CD} H_{\overline{50}ABCD} \ni 4 G_{\overline{50}ij} \tilde{t}_{(50)\alpha ij} t_{(\overline{50})}^{\alpha} \quad , \quad (\text{C.46})$$

where by $\tilde{t}_{(50)\alpha ij}$ we denote the antitriplet combination of $10_i \times 10_j$, that is $\tilde{t}_{(50)\alpha ij} = 1/4 \epsilon_{\alpha\gamma\delta} \epsilon_{ab} \tilde{T}_{(50)ij}^{ab\gamma\delta}$:

$$\tilde{T}_{(50)ij}^{ab\gamma\delta} = \sqrt{\frac{1}{3}} \left[\Psi_{10i}^{ab} \Psi_{10j}^{\gamma\delta} + \Psi_{10i}^{\gamma\delta} \Psi_{10j}^{ab} + \frac{\Psi_{10i}^{\gamma a} \Psi_{10j}^{\delta b} + \Psi_{10i}^{\delta b} \Psi_{10j}^{\gamma a} - \Psi_{10i}^{\gamma b} \Psi_{10j}^{\delta a} - \Psi_{10i}^{\delta a} \Psi_{10j}^{\gamma b}}{2} \right] \quad . \quad (\text{C.47})$$

Substituting the above expression in eq. (C.46) and exploiting the symmetry of $G_{\overline{50}}$, one obtains

$$I = \left[U_{\alpha i}^c \left(\frac{2G_{\overline{50} \ ij}}{\sqrt{3}} \right) E_j^c - \frac{1}{2} Q_i^{\beta} \left(-\frac{G_{\overline{50} \ ij}}{\sqrt{3}} \right) Q_j^{\gamma} \epsilon_{\alpha\beta\gamma} \right] H'_{3u} \quad . \quad (\text{C.48})$$

This is the cause of the introduction of the coefficient $\sqrt{3}/2$ in eq. (4.24).